REWRITING TECHNIQUES FOR RELATIVE COHERENCE

SAMUEL MIMRAM

LIX, CNRS, École polytechnique, Institut Polytechnique de Paris, 91120 Palaiseau, France *e-mail address*: samuel.mimram@polytechnique.edu

ABSTRACT. A series of works has established rewriting as an essential tool in order to prove coherence properties of algebraic structures, such as MacLane's coherence theorem for monoidal categories, based on the observation that, under reasonable assumptions, confluence diagrams for critical pairs provide the required coherence axioms. We are interested here in extending this approach simultaneously in two directions. Firstly, we want to take into account situations where coherence is partial, in the sense that it only applies to a subset of the structural morphisms. Secondly, we are interested in structures which are cartesian in the sense that variables can be duplicated or erased. We develop theorems and rewriting techniques in order to achieve this, first in the setting of abstract rewriting systems, and then extend them to term rewriting systems, suitably generalized to take coherence into account. As an illustration of our results, we explain how to recover the coherence theorems for monoidal and symmetric monoidal categories.

Contents

2
2
2
3
4
5
6
6
7
7
7
12
14
14
18
22
24
24

Key words and phrases: coherence, rewriting system, Lawvere theory.

4.2. Algebras for Lawvere 2-theories	25
5. Coherent term rewriting systems	28
5.1. Extended rewriting systems	28
5.2. Rewriting properties	32
6. Coherence for symmetric monoidal categories	35
6.1. A theory for symmetric monoidal categories	35
6.2. Every affine diagram commutes	36
6.3. General coherence	43
7. Future works	45
References	45

1. INTRODUCTION

1.1. Coherence results. Coherence results are fundamental in category theory. They can be seen both as a way of formally simplifying computations, by ensuring that we can consider strict algebraic structures without loss of generality, and as a guide for generalizing computations, by ensuring that we have correctly generalized algebraic structures in higher dimensions and taken higher-dimensional cells in account. Such results have been obtained for a wide variety of algebraic structures on categories, including monoidal categories [46, 47], symmetric monoidal categories [46, 47], braided monoidal categories [28], tortile monoidal categories [59], symmetric monoidal closed categories [26, 32, 65], biclosed monoidal (multi)categories [40], compact closed categories [31], cartesian closed categories [5, 52], rig categories [41], weakly distributive categories [9], bicategories and pseudo-functors [44], cartesian closed bicategories [17], tricategories [19, 24], etc.

The coherence results are often quickly summarized as "all diagrams commute". However, this is quite misleading [30]: firstly, we only want to consider diagrams made of structural morphisms, and secondly, we actually usually want to consider only a subset of those diagrams. In the general case, the goal is thus to identify a class of diagrams which commute or, better, decide equality between structural morphisms and provide explicit and combinatorial descriptions of those. One of the goals of this article is to clarify the situation and the relationships between the various approaches.

1.2. Coherence from rewriting theory. A field which provides many computational techniques to show that diagrams commutes is rewriting [4, 63]. Namely, when a rewriting system is terminating and locally confluent, which can be verified algorithmically by computing its critical pairs, it is confluent and thus has the Church-Rosser property, which implies that any two zig-zags can be filled by local confluence diagrams. By properly extending the notion of rewriting system with higher-dimensional cells in order to take coherence into account (those cells specifying which confluence diagrams commute), one is then able to show coherence results of the form "all diagrams commute" up to the coherence laws. This idea of extending rewriting theory in order to handle coherence dates back to pioneering work from people such as Newman [53] (who observed that the 1-complex associated to a convergent abstract rewriting system can be made into a simply connected 2-complex by adding 2-cells corresponding to confluence diagrams), Squier [37, 60] (who observed that the

confluence diagrams for critical pairs of a convergent monoid presentation generate the full congruence on the monoidal category associated to the rewriting system), or MacLane [47] (whose original proof for the coherence theorem for monoidal categories is implicitly based on rewriting techniques). Furthermore, the work of Power [54] and Street [62] have reformulated rewriting in categorical terms, and paved the way for higher-dimensional generalizations of rewriting in the context of computads [61], which are also known as polygraphs [3, 14], and have been used to recover various coherence theorems [22], that can often be interpreted as computing (polygraphic) resolutions in suitable settings [3]. More recently, these generalizations were also adapted to homotopy type theory [33]. We would also like to mention here that this trend of work has motivated extensions of rewriting to various 2-dimensional settings, which should certainly be helpful at some point in relation to coherence: monoidal categories [38], and hypergraph categories which allow rewriting modulo symmetries or Frobenius structures [11, 12, 13].

1.3. Coherence for monoidal categories. One of the first and most important instance of a coherence theorem is the one for monoidal categories, originally due to Mac Lane. Since it will be used in the following as one of the main illustrations, we begin by recalling it here, and discuss its various possible formulations.

A monoidal category consists of a category C equipped with a tensor bifunctor and unit object respectively noted

 $\otimes: C \times C \to C \qquad \qquad e: 1 \to C$

together with natural isomorphisms

$$lpha_{x,y,z}: (x\otimes y)\otimes z o x\otimes (y\otimes z) \qquad \qquad \lambda_x: e\otimes x o x \qquad \qquad
ho_x: x\otimes e o x$$

called *associator* and *left* and *right unitors*, satisfying two well-known axioms stating that the diagrams



commute for any objects x, y and z of C.

and

Thanks to these axioms, the way tensor expressions are bracketed does not really matter: we can always rebracket expressions using the structural morphisms α , λ and ρ , and any two ways of rebracketing an expression into the other are equal. In fact, and this is an important point in this article, there are various ways to formalize this [1]:

(M1) Every diagram in a free monoidal category made up of α , λ and ρ commutes [28, Corollary 1.6], [46, Theorem VI.2.1].

- (M2) Every diagram in a monoidal category made up of α , λ and ρ commutes [47, Theorem 3.1], [46, Theorem XI.3.2].
- (M3) Every monoidal category is monoidally equivalent to a strict monoidal category [28, Corollary 1.4], [46, Theorem XI.3.1].
- (M4) The forgetful 2-functor from strict monoidal categories to monoidal categories has a left adjoint and the components of the unit are equivalences.

Condition (M2) implies (M1) as a particular case and the converse implication can also be shown, so that the two are equivalent. Condition (M4) implies (M3) as a particular case, and it can be shown that (M3) in turn implies (M2), see [46, Theorem XI.3.2].

1.4. Coherence for symmetric monoidal categories. Although fundamental, taking the previous example of a coherence theorem as a guiding example can be misleading as it hides the fact that the coherence results are in general more subtle: usually, we do not want all the diagrams made of structural morphisms to commute. In order to illustrate this, let us consider the following variant of monoidal categories.

A symmetric monoidal category is a monoidal category equipped with a natural transformation

$$\gamma_{x,y}: x \otimes y \to y \otimes x$$

called *symmetry* such that the diagrams



and

commute for every objects x, y and z of C.

Analogous coherence theorems as above hold and can be formulated as follows:

- (S1) Every "generic" diagram in a (free) symmetric monoidal category made up of α , λ , ρ and γ commutes.
- (S2) Every diagram in a (free) symmetric monoidal category made up of α, λ ρ and γ commutes precisely when the two sides have the same underlying symmetry [28, Corollary 2.6], [46, Theorem XI.1.1].
- (S3) Every (free) symmetric monoidal category is symmetric monoidally equivalent to a strict symmetric monoidal category [49, Proposition 4.2], [28, Theorem 2.5].
- (S4) The forgetful 2-functor from strict symmetric monoidal categories to symmetric monoidal categories has a left adjoint and the components of the unit are equivalences.

We can see above that the formulations do not anymore require that "all diagrams commute". In order to illustrate why it has to be so, observe that the diagram

$$x \otimes x \underbrace{\stackrel{\gamma_{x,x}}{\underset{\mathrm{id}_{x\otimes x}}{\overset{\gamma_{x,x}}}{\overset{\gamma_{x,x}}{\overset{\gamma_{x,x}}}{\overset{\gamma_{x,x}}{\overset{\gamma_{x,x}}}{\overset{\gamma_{x,x}}}{\overset{\gamma_{x,x}}}{\overset{\gamma_{x,x}}}{\overset{\gamma_{x,x}}}}}}}}}}}}}}}}}}}}}}$$

does *not* commute in general in monoidal categories, although its morphisms are structural ones. For a concrete example, consider the category of sets and functions equipped with cartesian product as tensor product and x to be any set with at least two distinct elements a and b. We namely have

$$\gamma_{x,x}(a,b) = (b,a) \neq (a,b) = \mathrm{id}_{x \otimes x}(a,b).$$

However, note that the two morphisms do not have the same "underlying symmetry" ($\gamma_{x,x}$ corresponds to a transposition, whereas $id_{x\otimes x}$ to an identity on a 2-element set). In fact, as stated in (S2), restricting to diagrams where the two morphisms induce the same symmetry is enough to have them always commute. Another way to ensure that the diagrams should commute is to require them to be generic (or linear) as in (S1), by which we roughly mean that all the objects occurring in the source (or target) object should be distinct: this is not the case in (1.1) since the source object is $x \otimes x$, in which x occurs twice. Intuitively, this condition ensures that the underlying symmetry of the morphisms is uniquely determined by the positions of the variables, and thus that the diagram commutes as a particular instance of (S1). The same subtlety is implicitly present in the condition (S3): for a strict symmetric monoidal category, we do not require that we the symmetric should be strict (only the associator and unitors, such a category is sometimes also called a permutative category [49]).

1.5. Coherence for more general theories. In order to work in a framework which is able to handle many algebraic structures at once (monoidal categories, symmetric monoidal categories, etc.), we axiomatize the notion of structure using Lawvere 2-theories, which are Lawvere theories enriched in groupoids. In such a theory \mathcal{T} , the 0-cells encode the arities, the 1-cells the operations, and the 2-cells the coherence morphisms, so that a product preserving 2-functor $\mathcal{T} \to \mathbf{Cat}$, i.e. an algebra for the theory, corresponds to an actual algebraic structure. In order to perform computations on such a theory, it is often convenient to use a description of it by the means of generators and relations, possibly with good properties, and we use here the notion of term rewriting system, extended in order to account for relations between coherences morphisms (which correspond to zig-zags of rewriting steps).

For "fully coherent" structures (e.g. monoidal categories) one wants to show that all diagrams made of 2-cells commute, i.e. the Lawvere 2-theory contains at most one 2-cell between two given 1-cells. For more general situations (e.g. symmetric monoidal categories), one wants to show coherence results relatively to a subtheory $\mathcal{W} \subseteq \mathcal{T}$ (for symmetric monoidal categories, this would be the one generated by α , λ and ρ). Writing \mathcal{T}/\mathcal{W} for the quotient theory (the theory obtained from \mathcal{T} by turning all the 2-cells of \mathcal{W} into identities), previous conditions can then formulated in this setting in the following way:

- (C1) Identify a class of pairs of 1-cells (f_i, g_i) such that the hom-categories $\mathcal{T}/\mathcal{W}(f_i, g_i)$ contain at most one morphism.
- (C2) Provide an "explicit" description of the quotient theory \mathcal{T}/\mathcal{W} .
- (C3) Show that every algebra of \mathcal{T} is equivalent to an algebra of \mathcal{T}/\mathcal{W} .

(C4) Show that the forgetful 2-functor from algebras of \mathcal{T}/\mathcal{W} to algebras of \mathcal{T} has a left adjoint and the components of the unit are equivalences.

Variants of the condition (C2) have been considered in the literature such as the problem of deciding the equality of 2-cells in \mathcal{T} (or, equivalently, in free algebras), or provide explicit descriptions of free algebras. An important point is that, in order for the above conditions to make sense, one should first make sure that the quotient is a faithful representation of the original theory (in the sense that the canonical 2-functor $\mathcal{T} \to \mathcal{T}/\mathcal{W}$ is a local equivalence) which, as we show, is the case if and only if \mathcal{W} is 2-rigid (theorem 55), i.e. has at most one 2-cell between any pair of 1-cells. We illustrate here coherence conditions (C1) and (C2) in the case of symmetric monoidal categories, where they are respectively proved in theorems 97 and 101. Coherence condition (C3) and (C4) are respectively formulated as conjectures 60 and 61 and left for future work.

1.6. Other forms of coherence. There are other possible formulations of coherence, involving what are called *unbiased* variants of the structures. In the case of monoidal categories, an *unbiased monoidal category* is a category equipped with *n*-ary tensor products for every natural number n, satisfying suitable axioms [43, Section 3.1]. The following variant of (M4) can then be shown:

(M4') The forgetful 2-functor from strict monoidal categories to unbiased monoidal categories has a left adjoint and the components of the unit are equivalences.

This result is in fact a particular instance of a very general coherence theorem due to Power [55], see also [35, 58], which originates in the following observation: there is a 2-monad T on **Cat** whose strict algebras are strict monoidal categories and whose pseudoalgebras are unbiased monoidal categories. Given a 2-monad T on a 2-category, under suitable assumptions (which are satisfied in the case of the monad of monoidal categories), it can be shown that the inclusion T-**StrAlg** \rightarrow T-**PsAlg** of 2-categories, from the 2-category of strict T-algebras (and strict morphisms) to the 2-category of pseudo-2-algebras (and pseudo-morphisms) admits a left 2-adjoint (which can be interpreted as a strictification 2-functor) such that the components of the unit of the adjunction are internal equivalences in T-pseudo-algebras.

We do not insist much on this general route, as our main concern here is the relationship with rewriting, which provides ways of handling biased notions of algebras.

1.7. Contents of the paper. We first investigate, in section 2, an abstract version of the situation and formally compare the various coherence theorems: we show that quotienting a theory by a subtheory \mathcal{W} gives rise to an equivalent theory if and only if \mathcal{W} is coherent (or *rigid*), in the sense that all diagrams commute (theorem 16). Moreover, this is the case if and only if they give rise to equivalent categories of algebras (proposition 21), which can be thought of as a strengthened version of (C4). We then provide, in section 3, rewriting conditions which allow showing coherence in practice (proposition 42).

Those results are extended, in section 4, to the setting of Lawvere 2-theories, where we are able to axiomatize (symmetric) monoidal categories. One of the main novelties here consists in allowing for coherence with respect to a subtheory \mathcal{W} , which is required to handle coherence for symmetric monoidal categories. This leads us to conjecture that, when the subtheory is rigid, we always have coherence for algebras (conjectures 60 and 61). The associated rewriting tools are developed in section 5, based on a coherent extension of term

rewriting systems (definition 68), following [8, 16, 48]. In particular, we provide rewriting tools to show that this theory is rigid (proposition 83). We also show in section 6 that these tools apply in the case of the theory of symmetric monoidal categories, and use those to recover one of the classical coherence theorems in this setting (theorem 97).

Sections 2 and 4 develop the general categorical setting and can be read independently from sections 3, 5 and 6, which specifically develop rewriting tools and applications.

This article is an extended version of [50], and also corrects a few mistakes unfortunately present there.

1.8. Acknowledgments. The author would like to thank the anonymous reviewers for their comments, which improved the paper.

2. Relative coherence for categories

2.1. Quotient of categories. Fix a category C together with a set W of isomorphisms of C. Although the situation is very generic, and the following explanation is only vague for now, it can be helpful to think of C as a theory describing a structure a category can possess, and W as the morphisms we are interested in strictifying. For instance, if we are interested in the coherence theorem for symmetric monoidal categories, we can think of the objects of C as formal iterated tensor products, the morphisms of C as the structural morphisms (the composites of α , λ , ρ and γ), and we would typically take W as consisting of all instances of α , λ and ρ (but not γ). This will be made formal in section 4.

Definition 1. A functor $F : \mathcal{C} \to \mathcal{D}$ is *W*-strict when it sends every morphism of *W* to an identity.

Definition 2. The quotient \mathcal{C}/W of \mathcal{C} under W is the category equipped with a W-strict functor $\mathcal{C} \to \mathcal{C}/W$, such that every W-strict functor $F : \mathcal{C} \to \mathcal{D}$ extends uniquely as a functor $\tilde{F} : \mathcal{C}/W \to \mathcal{D}$ making the following diagram commute:



Such quotient categories always exist [7], and we provide below an explicit construction in nicely behaved cases (proposition 10). We write \mathcal{W} for the subcategory of \mathcal{C} generated by W. This subcategory will be assimilated to the smallest subset of morphisms of \mathcal{C} which contains W is closed under compositions and identities. The category \mathcal{W} is a groupoid and it can shown that passing from W to \mathcal{W} does not change the quotient.

Lemma 3. The categories C/W and C/W are isomorphic.

Proof. By definition of quotient categories (definition 2), it is enough to show that the category \mathcal{C}/W is a quotient of \mathcal{C} by \mathcal{W} . It follows easily from the fact that a functor $\mathcal{C} \to \mathcal{D}$ is W-strict if and only if it is \mathcal{W} -strict. Namely, the left-to-right implication follows from functoriality and the right-to-left implication from the inclusion $W \subseteq \mathcal{W}$.

Thanks to the above lemma, we will be able to assume, without loss of generality, that we always quotient categories by a subgroupoid which has the same objects as C.

We will see that quotients are much better behaved when the groupoid we quotient by satisfies the following property.

Definition 4. A groupoid \mathcal{W} is *rigid* when any two morphisms $f, g : x \to y$ which are parallel (i.e. have the same source, and have the same target) are necessarily equal.

Such a groupoid can be thought of as a "coherent" sub-theory of C: it does not have non-trivial geometric structure in the sense of proposition 7 below.

We will need to use the following properties of categories.

Definition 5. A category is

- *discrete* when its only morphisms are identities,
- *contractible* when it is equivalent to the terminal category,
- connected when there is a morphism between any two objects,
- propositional when it is a rigid and connected groupoid.

Lemma 6. A propositional category with an object is contractible.

Proof. Given a propositional category C, the terminal functor $C \to 1$ is full (because C is connected), faithful (because C is rigid) and surjective (because C has an object).

Proposition 7. Given a groupoid \mathcal{W} , the following are equivalent

- (i) \mathcal{W} is rigid,
- (ii) W has identities as only automorphisms,
- (iii) \mathcal{W} is equivalent to a discrete category,
- (iv) \mathcal{W} is a coproduct of contractible categories.

Proof. (i) implies (ii). Given a rigid category, any automorphism $f: x \to x$ is parallel with the identity and thus has to be equal to it.

(ii) implies (i). Given two parallel morphisms $f, g: x \to y$, we have $g^{-1} \circ f = \mathrm{id}_x$ and thus f = g.

(i) implies (iii). Write \mathcal{D} for category of connected components of \mathcal{W} : this is the discrete category whose objects are the equivalence classes [x] of objects x of \mathcal{W} under the equivalence relation identifying x and y whenever there is a morphism $f: x \to y$ in \mathcal{W} . The quotient functor $Q: \mathcal{W} \to \mathcal{D}$ is full because \mathcal{D} is discrete, faithful because \mathcal{W} is rigid, and surjective on objects by construction of \mathcal{D} . It is thus an equivalence of categories.

(iii) implies (i). Given an equivalence $F : \mathcal{W} \to \mathcal{D}$ to a discrete category \mathcal{D} , any two parallel morphisms $f, g : x \to y$ have the same image Ff = Fg (which is an identity) because \mathcal{D} is discrete, and are thus equal because F is faithful.

(iii) implies (iv). Consider a functor $F: \mathcal{W} \to \mathcal{D}$ which is an equivalence with \mathcal{D} discrete. Given $x \in \mathcal{D}$, we write $F^{-1}x$ for the full subcategory of \mathcal{W} whose objects are sent to x by F. Since \mathcal{D} is discrete, for any morphism $f: x \to y$ in \mathcal{D} , we have Fx = Fy, from which it follows that $\mathcal{W} \cong \bigsqcup_{x \in \mathcal{D}} F^{-1}x$. Since \mathcal{D} is discrete, each $F^{-1}x$ is non-empty, connected and rigid and thus contractible by lemma 6.

(iv) implies (iii). If $\mathcal{W} \cong \bigsqcup_{i \in I} \mathcal{W}_i$ with \mathcal{W}_i contractible, i.e. $\mathcal{W}_i \simeq 1$, then $\mathcal{W} \simeq \bigsqcup_{i \in I} 1$ because equivalences are closed under coproducts and thus \mathcal{W} is equivalent to a discrete category.

The fact that $\mathcal{W} \subseteq \mathcal{C}$ is rigid can be thought of here as the fact that coherence condition (C1) holds for \mathcal{C} , relatively to \mathcal{W} : any two parallel structural morphisms are equal. Conditions (iii) and (iv) can also be interpreted as stating that \mathcal{W} is a set, up to equivalence.

General notions of quotients (with respect to a subcategory, or to a general notion of congruence both on objects and morphisms) have been developed in [7], and are non-trivial to study and construct. However, when quotienting a category \mathcal{C} by a rigid subgroupoid \mathcal{W} , we have the following simple description. In this case, we define the two following equivalence relations $\sim_{\mathcal{W}}$, that we often simply write \sim .

- We write $\sim_{\mathcal{W}}$ for the equivalence relation on objects of \mathcal{C} such that $x \sim y$ whenever there is a morphism $f: x \to y$ in \mathcal{W} . When it exists, such a morphism is unique by rigidity of \mathcal{W} and noted $w_{x,y}: x \to y$.
- We also write $\sim_{\mathcal{W}}$ for the equivalence relation on morphisms of \mathcal{C} such that for $f: x \to y$ and $f': x' \to y'$ we have $f \sim f'$ whenever there exists morphisms $v: x \to x'$ and $w: y \to y'$ in \mathcal{W} making the following diagram commute:



Definition 8. Let C be a category equipped with a rigid subgroupoid W. We write C/\sim_W for the category where

- an object [x] is an equivalence class of an object x of C under \sim ,
- a morphism $[f]: [x] \to [y]$ is the equivalence class of a morphism $f: x \to y$ in \mathcal{C} under \sim , - given morphisms $f: x \to y$ and $g: y' \to z$ with [y] = [y'], their composition is the
 - morphism $[g] \circ [f] = [g \circ w_{y,y'} \circ f]$:

$$x \xrightarrow{f} y \xrightarrow{w_{y,y'}} y' \xrightarrow{g} z$$

- the identity on an object [x] is $[id_x]$.

Note that there is a canonical functor $[-]: \mathcal{C} \to \mathcal{C}/\sim_{\mathcal{W}}$ sending (resp. a morphism) to its equivalence class.

Lemma 9. The above category is well-defined.

Proof. We can check that the operations are well-defined and axioms of categories are satisfied.

- Composition is compatible with the equivalence relation. Given $f_1: x_1 \to y_1, f_2: x_2 \to y_2, g_1: y'_1 \to z_1, g_2: y'_2 \to z_2$ such that $f_1 \sim f_2$ and $g_1 \sim g_2$ (and thus $x_1 \sim x_2, y_1 \sim y_2, y'_1 \sim y'_2$ and $z_1 \sim z_2$) which are composable (i.e. $y_1 \sim y'_1$ and $y_2 \sim y'_2$), the following diagram shows that $[g_1] \circ [f_1] = [g_2] \circ [f_2]$:

where the squares on the left and right respectively commute because $f_1 \sim f_2$ and $g_1 \sim g_2$ and the one in the middle does by rigidity of W.

- Identities are compatible with the equivalence relations. Given objects x and y of C such that $x \sim y$, the diagram

$$\begin{array}{cccc} x & \stackrel{\operatorname{id}_x}{\longrightarrow} & x \\ w_{x,y} & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & y \end{array} \xrightarrow{\operatorname{id}_y} & y \end{array}$$

commutes showing that we have $\mathrm{id}_x \sim \mathrm{id}_y$.

Associativity of composition and the fact that identities are neutral element for composition follow immediately from the fact that those properties are satisfied in C.

Proposition 10. The category C/W is isomorphic to C/\sim_W .

Proof. We show that the category has the universal property of definition 2. The quotient functor $[-]: \mathcal{C} \to \mathcal{C}/\mathcal{W}$ is \mathcal{W} -strict, i.e. for any morphism $w: x \to y$, we have $[w] = [\mathrm{id}_y]$:

$$\begin{array}{ccc} x & \stackrel{w}{\longrightarrow} y \\ & \stackrel{i}{\downarrow} & \stackrel{i}{\downarrow} \operatorname{id}_{y} \\ & \stackrel{i}{\downarrow} & \stackrel{i}{\downarrow} \operatorname{id}_{y} \end{array}$$

Moreover, a \mathcal{W} -strict functor $F : \mathcal{C} \to \mathcal{D}$ induces a unique functor $\tilde{F} : \mathcal{C}/\mathcal{W} \to \mathcal{D}$. Namely, by \mathcal{W} -strictness, two objects (resp. morphisms) which are equivalent have the same image by F.

When $\mathcal{W} \subseteq \mathcal{C}$ is not rigid, we can have a similar description of the quotient, but the description is more complicated. Namely, if we are trying to compose two morphisms [f] and [g] in the quotient with $f: x \to y$ and $g: y' \to z$, we might have multiple morphisms $y \to y'$ in \mathcal{W} (say v and w),

$$x \xrightarrow{f} y \xrightarrow{z} y' \xrightarrow{g} z$$

In such a situation, the compositions $g \circ v \circ f$ and $g \circ w \circ f$ should be identified in the quotient. This observation suggests that the construction of the quotient category C/W, when W is not rigid, is better described in two steps: we first formally make W rigid, and then apply proposition 10.

Definition 11. A functor $F : \mathcal{C} \to \mathcal{D}$ is \mathcal{W} -rigid when for any parallel morphisms $f, g : x \to y$ of \mathcal{C} we have Ff = Fg.

Definition 12. The *W*-rigidification $\mathcal{C}/\!\!/\mathcal{W}$ of \mathcal{C} is the category equipped with a *W*-rigid functor $\mathcal{C} \to \mathcal{C}/\!\!/\mathcal{W}$, such that any *W*-rigid functor $F : \mathcal{C} \to \mathcal{D}$ extends uniquely as a functor $\tilde{F} : \mathcal{C}/\!\!/\mathcal{W} \to \mathcal{D}$ making the following diagram commute:



Lemma 13. The category $C/\!\!/W$ is the category obtained from C by quotienting morphisms under the smallest congruence (wrt composition) identifying any two parallel morphisms of W.

Proposition 14. The quotient C/W is isomorphic to $(C/W)/\tilde{W}$ where \tilde{W} is the set of equivalence classes of morphisms in W under the equivalence relation of lemma 13.

Proof. Since any \mathcal{W} -strict functors are \mathcal{W} -rigid, we have that any \mathcal{W} -strict functor extends as a unique functor $\tilde{F} : \mathcal{C}/\!\!/\mathcal{W} \to \mathcal{D}$ which is \mathcal{W} -rigid, and thus as a unique functor $\mathcal{C}/\tilde{\mathcal{W}} \to \mathcal{D}$:



A consequence of the preceding explicit description of the quotient is the following:

Lemma 15. The quotient functor $\mathcal{C} \to \mathcal{C}/\mathcal{W}$ is surjective on objects and full.

Proof. By proposition 14, the quotient functor is the composite of the quotient functors $\mathcal{C} \to \mathcal{C}/\!\!/\mathcal{W} \to \mathcal{C}/\mathcal{W}$. The first one is surjective on objects and full by lemma 13 and the second one is surjective on objects and full by proposition 10.

This entails the following theorem, which is the main result of the section. Its meaning can be explained by taking the point of view given above: thinking of the category C as describing a structure, and of W as the part of the structure we want to strictify, the structure is equivalent to its strict variant if and only if the quotiented structure does not itself bear non-trivial geometry, in the sense of proposition 7.

Theorem 16. Suppose that \mathcal{W} is a subgroupoid of \mathcal{C} . The quotient functor $[-]: \mathcal{C} \to \mathcal{C}/\mathcal{W}$ is an equivalence of categories if and only if \mathcal{W} is rigid.

Proof. Since the quotient functor is always surjective and full by lemma 15, it remains to show that it is faithful if and only if \mathcal{W} is rigid. Suppose that the quotient functor is faithful. Given $w, w' : x \to y$ in \mathcal{W} , by lemma 13 and proposition 14 we have [w] = [w'] and thus w = w' by faithfulness. Suppose that \mathcal{W} is rigid. The category \mathcal{C}/\mathcal{W} then admits the description given in proposition 10. Given $f, g : x \to y$ in \mathcal{C} such that [f] = [g], there is $v : x \to x$ and $w : y \to y$ such that $w \circ f = g \circ v$. By rigidity, both v and w are identities and thus f = g.

Example 17. As a simple example, consider the groupoid \mathcal{C} freely generated by the graph

$$x \xrightarrow{f} y$$

The subgroupoid generated by $W = \{g\}$ is rigid, so that \mathcal{C} is equivalent to the quotient category \mathcal{C}/W , which is the groupoid generated by the graph

However, the groupoid generated by $W = \{f, g\}$ is not rigid (since we do not have f = g). In this case, C is not equivalent to the quotient category C/W, which is the terminal category.

Remark 18. By taking C to be W in theorem 16, we obtain that a category W is rigid if and only if it is equivalent to the discrete category of its connected components. This thus provides an alternative proof of condition (iii) of proposition 7.

2.2. Coherence for algebras. Given two categories C and D, an algebra of C in D is a functor from C to D. In the following, we will mostly be interested in the case where D = Cat: if we think of the category C as describing an algebraic structure (e.g. the one of monoidal categories), an algebra can be thought of as a category actually possessing this structure (an actual monoidal category). We write Alg(C, D) for the category whose objects are algebras and morphisms are natural transformations, and Alg(C) = Alg(C, Cat). Note that any functor

$$F: \mathcal{C} \to \mathcal{C}'$$

induces, by precomposition, a functor

$$\operatorname{Alg}(F, \mathcal{D}) : \operatorname{Alg}(\mathcal{C}', \mathcal{D}) \to \operatorname{Alg}(\mathcal{C}, \mathcal{D}).$$

We can characterize situations where two categories give rise to the same algebras as follows.

Proposition 19. A functor $F : \mathcal{C} \to \mathcal{C}'$ is an equivalence if and only if there is a family of equivalences of categories $\operatorname{Alg}(\mathcal{C}, \mathcal{D}) \simeq \operatorname{Alg}(\mathcal{C}', \mathcal{D})$ which is natural in \mathcal{D} .

Proof. Given a 2-category \mathcal{K} , one can define a Yoneda functor

$$Y_{\mathcal{K}}: \mathcal{K}^{\mathrm{op}} \to [\mathcal{K}, \mathbf{Cat}]$$
$$c \mapsto \mathcal{K}(c, -)$$

where **Cat** is the 2-category of categories, functors and natural transformations, and $[\mathcal{K}, \mathbf{Cat}]$ denotes the 2-category of 2-functors $\mathcal{K} \to \mathbf{Cat}$, transformations and modifications. In particular, given 0-cells $c \in \mathcal{K}^{\text{op}}$ and $d \in \mathcal{K}$, we have $Y_{\mathcal{K}}cd = \mathcal{K}(c, d)$. The Yoneda lemma states that this functor is a local isomorphism (this is a particular case of the Yoneda lemma for bicategories [27, Corollary 8.3.13]): this means that, for every objects $c, d \in \mathcal{K}^{\text{op}}$, we have an isomorphism of categories

$$[\mathcal{K}, \mathbf{Cat}](Y_{\mathcal{K}}c, Y_{\mathcal{K}}d) \cong \mathcal{K}(d, c).$$

In particular, taking $\mathcal{K} = \mathbf{Cat}$ (and ignoring size issues, see below for a way to properly handle this), the Yoneda functor sends a category $\mathcal{C} \in \mathbf{Cat}^{\mathrm{op}}$ to $Y_{\mathbf{Cat}}\mathcal{C}$, i.e. $\mathbf{Cat}(\mathcal{C}, -)$, i.e. $\mathrm{Alg}(\mathcal{C}, -)$. Given category \mathcal{C} and \mathcal{C}' , by the Yoneda lemma, we thus have an isomorphism of categories

$$[\mathbf{Cat},\mathbf{Cat}](\mathrm{Alg}(\mathcal{C},-),\mathrm{Alg}(\mathcal{C}',-))\cong\mathbf{Cat}(\mathcal{C}',\mathcal{C})$$

which is compatible with 0-composition in **Cat**. The categories C and C' are thus equivalent, if and only if the categories $\operatorname{Alg}(\mathcal{C}, -)$ and $\operatorname{Alg}(\mathcal{C}', -)$ are equivalent, which is the case if and only if there is a family of equivalences of categories between $\operatorname{Alg}(\mathcal{C}, \mathcal{D})$ and $\operatorname{Alg}(\mathcal{C}, \mathcal{D})$ natural in \mathcal{D} .

Alternative proof of proposition 19. We provide here an alternative more pedestrian proof, inspired of [18, Lemma 5.3.1], which does not require ignoring size issues. Suppose that F is an equivalence of categories, with pseudo-inverse $G : \mathcal{C}' \to \mathcal{C}$, i.e. we have $G \circ F \cong \mathrm{Id}_{\mathcal{C}}$ and $F \circ G \cong \mathrm{Id}_{\mathcal{C}'}$. We define functors

$$\begin{array}{ll} \operatorname{Alg}(F,\mathcal{D}):\operatorname{Alg}(\mathcal{C}',\mathcal{D})\to\operatorname{Alg}(\mathcal{C},\mathcal{D}) & \operatorname{Alg}(G,\mathcal{D}):\operatorname{Alg}(\mathcal{C},\mathcal{D})\to\operatorname{Alg}(\mathcal{C}',\mathcal{D}) \\ & A\mapsto A\circ F & A\mapsto A\circ G \end{array}$$

Those induce an equivalence between $\operatorname{Alg}(\mathcal{C}, \mathcal{D})$ and $\operatorname{Alg}(\mathcal{C}', \mathcal{D})$ since, for $A \in \operatorname{Alg}(\mathcal{C}, \mathcal{D})$ and $A' \in \operatorname{Alg}(\mathcal{C}', \mathcal{D})$, we have

$$Alg(G, \mathcal{D}) \circ Alg(F, \mathcal{D}) = A' \circ F \circ G \simeq A'$$
$$Alg(F, \mathcal{D}) \circ Alg(G, \mathcal{D}) = A \circ G \circ F \simeq A$$

Moreover, the family of equivalences $\operatorname{Alg}(F, \mathcal{D})$ is natural in \mathcal{D} , and similarly for $\operatorname{Alg}(G, \mathcal{D})$. Namely, given $H : \mathcal{D} \to \mathcal{D}'$ and considering the functor

$$\operatorname{Alg}(\mathcal{C}, H) : \operatorname{Alg}(\mathcal{C}, \mathcal{D}) \to \operatorname{Alg}(\mathcal{C}, \mathcal{D}')$$

 $A \mapsto H \circ A$

as well as the variant with \mathcal{C}' instead of \mathcal{C} , the diagram

commutes: an object $A \in Alg(\mathcal{C}', \mathcal{D})$ is sent to $H \circ A \circ F$ by both sides.

Conversely, suppose given an equivalence of categories

$$\Phi_{\mathcal{D}} : \operatorname{Alg}(\mathcal{C}', \mathcal{D}) \leftrightarrow \operatorname{Alg}(\mathcal{C}, \mathcal{D}) : \Psi_{\mathcal{D}}$$

which is natural in \mathcal{D} . We define

$$F = \Phi_{\mathcal{C}'}(\mathrm{Id}_{\mathcal{C}'}) : \mathcal{C} \to \mathcal{C}' \qquad \qquad G = \Psi_{\mathcal{C}}(\mathrm{Id}_{\mathcal{C}}) : \mathcal{C}' \to \mathcal{C}$$

and we have

$$G \circ F = G \circ \Phi_{\mathcal{C}'}(\mathrm{Id}_{\mathcal{C}'}) = \Phi_{\mathcal{C}'}(G \circ \mathrm{Id}_{\mathcal{C}'}) = \Phi_{\mathcal{C}'}(\Psi_{\mathcal{C}}(\mathrm{Id}_{\mathcal{C}})) \cong \mathrm{Id}_{\mathcal{C}'}$$

(the second equality is naturality), and similarly for $G \circ F \cong \mathrm{Id}_{\mathcal{C}}$.

Remark 20. We would like to point out a subtle point with respect to naturality in the above theorem. Given a functor $F : \mathcal{C} \to \mathcal{C}'$, it is always the case that the induced functors $\operatorname{Alg}(F, \mathcal{D})$ form a family which is natural in \mathcal{D} . Suppose moreover that all the functors $\operatorname{Alg}(F, \mathcal{D})$ are equivalences. We do not see any argument to show that the pseudo-inverse functors form a natural family, which is why we have to additionally impose this condition.

As a particular application, we have the following proposition, which can be interpreted as the equivalence of coherence conditions (C1) and a strengthened variant of (C4):

Proposition 21. Let C be a category together with a subgroupoid W. We have, for any category D, a functor

$$F_{\mathcal{D}} : \operatorname{Alg}(\mathcal{C}/\mathcal{W}, \mathcal{D}) \to \operatorname{Alg}(\mathcal{C}, \mathcal{D})$$
 (2.1)

induced by precomposition with the quotient functor $\mathcal{C} \to \mathcal{C}/\mathcal{W}$. These functors $(F_{\mathcal{D}})_{\mathcal{D} \in \mathbf{Cat}}$ form a family of equivalences of categories, natural in \mathcal{D} , if and only if \mathcal{W} is rigid.

Proof. By theorem 16, \mathcal{W} is rigid if and only if the quotient functor $\mathcal{C} \to \mathcal{C}/\mathcal{W}$ is an equivalence, and we conclude by proposition 19.

Remark 22. It can be wondered whether the case where $\mathcal{D} = \mathbf{Cat}$ is enough, i.e. whether the quotient functor $\mathcal{C} \to \mathcal{C}/\mathcal{W}$ is an equivalence whenever the functor $\operatorname{Alg}(\mathcal{C}/W) \to \operatorname{Alg}(\mathcal{C})$ it induces is an equivalence. We leave it as an open question, but remark here that it cannot follow from general results: it is not the case that two categories \mathcal{C} and \mathcal{C}' are equivalent whenever $\operatorname{Alg}(\mathcal{C})$ and $\operatorname{Alg}(\mathcal{C}')$ are equivalent. It is namely known that two Cauchy equivalent categories give rise to the same algebras in \mathbf{Cat} , see [36], so that the categories

$$\mathcal{C} = x \supseteq e \qquad \qquad \mathcal{C}' = x \xrightarrow{J} y$$

where $e \circ e = e$ have the same algebras.

3. Coherent abstract rewriting systems

Previous sections illustrate the importance of the property of being rigid for a groupoid, and we now provide tools to show this in practice, based on tools originating from rewriting theory. In the same way the theory of rewriting can be studied "abstractly" [4, 25, 63], i.e. without taking in consideration the structure of the objects being rewritten, we first develop the coherence theorems of interest in an abstract setting.

3.1. Extended abstract rewriting systems. The categorical formalization of the notion of rewriting system given here is based on the notion of polygraph [3, 14, 61].

Definition 23. An abstract rewriting system, or ARS, or 1-polygraph is a diagram

$$\mathsf{P}_0 \xleftarrow[t_0]{s_0} \mathsf{P}_1$$

in the category **Set**.

An ARS is simply another name for a directed graph. It consists of a set P_0 whose elements are the *objects* of interest, a set P_1 of *rewriting rules* and two functions s_0 and t_0 respectively associating to a rewriting rule its *source* and *target*. We often write

$$a: x \to y$$

to denote a rewriting rule a with $s_0(a) = x$ and $t_0(a) = y$. We write P_1^* for the set of *rewriting paths* in the ARS: its elements are (possibly empty) finite sequences a_1, \ldots, a_n of rewriting steps, which are composable in the sense that $t_0(a_i) = s_0(a_{i+1})$ for $1 \le i < n$. Writing $x_i = t_0(a_i)$, such a path can thus be represented as

$$x_0 \xrightarrow{a_1} x_1 \xrightarrow{a_2} \cdots \xrightarrow{a_n} x_n.$$

The source and target of such a rewriting path are respectively $x_0 = s_0(a_1)$ and $x_n = t_0(a_n)$, and n is called the *length* of the path. We sometimes write

$$p: x \xrightarrow{*} y$$

to indicate that p is a rewriting path with x as source and y as target. Given two composable paths $p: x \xrightarrow{*} y$ and $q: y \xrightarrow{*} z$, we write $p \cdot q: x \xrightarrow{*} z$ for their concatenation.

A morphism $f : \mathsf{P} \to \mathsf{Q}$ of ARS is a pair of functions $f_0 : \mathsf{P}_0 \to \mathsf{Q}_0$ and $f_1 : \mathsf{P}_1 \to \mathsf{Q}_1$ such that $s_0 \circ f_1 = f_0 \circ s_0$ and $t_0 \circ f_1 = f_0 \circ t_0$:

$$\begin{array}{c|c} \mathsf{P}_0 \rightleftharpoons s_0 \\ f_0 & \downarrow \\ f_0 & \downarrow \\ \mathsf{Q}_0 \rightleftharpoons s_0 \\ \hline t_0 & \mathsf{Q}_1 \end{array} \mathsf{P}_1$$

We write \mathbf{Pol}_1 for the resulting category or ARS and their morphisms. There is a forgetful functor $\mathbf{Cat} \rightarrow \mathbf{Pol}_1$, sending a category C to the ARS whose objects are those of C and whose rewriting steps are the morphisms of C.

Lemma 24. The forgetful functor $\mathbf{Cat} \to \mathbf{Pol}_1$ admits a left adjoint $-^* : \mathbf{Pol}_1 \to \mathbf{Cat}$. It sends an ARS to the category with P_0 as objects and P_1^* as morphisms, where composition is given by concatenation of paths and identities are the empty paths

Proof. This fact is easily checked directly, but an abstract argument for the existence of the left adjoint in such situations is the following: the categories **Cat** and **Pol**₁ are models of projectives sketches and the forgetful functor **Cat** \rightarrow **Pol**₁ is induced by a functor of sketches (the "inclusion" of the sketch of ARS into the sketch of categories) and, as such, it admits a left adjoint [6, Theorem 4.1], [3, Proposition 15.1.3].

As a variant of the preceding situation, we can consider the forgetful functor $\mathbf{Gpd} \to \mathbf{Pol}_1$, from the category of groupoids. It also admits a left adjoint $-^{\sim} : \mathbf{Pol}_1 \to \mathbf{Gpd}$, which can be described as follows. Given an ARS P, we write P^{\pm} for the ARS

$$\mathsf{P}_0 \xleftarrow[t_0]{s_0} (\mathsf{P}_1 \times \{-,+\})$$

Its objects are the same as for P. A rule in P_1^{\pm} is a pair (a, ϵ) consisting of a rewriting rule $a \in \mathsf{P}_1$ and $\epsilon \in \{-, +\}$, which will be noted a^{ϵ} in the following. The source and target maps are given by

$$s_0(a^+) = s_0(a)$$
 $t_0(a^+) = t_0(a)$ $s_0(a^-) = t_0(a)$ $t_0(a^-) = s_0(a)$

We can think of a^+ as corresponding to a and a^- as corresponding to a taken "backward". A *rewriting zig-zag* is a path $a_1^{\epsilon_1}, \ldots, a_n^{\epsilon_n}$ in P^{\pm} . The intuition is that a zig-zag is a "non-directed" rewriting path, consisting of rewriting steps, some of which are taken backward. We write

$$p: x \xrightarrow{\sim} y$$

to indicate that p is a zig-zag from x to y. Two zig-zags are *congruent* when they are related by the smallest congruence \sim such that, for every rewriting rule $a: x \to y$, we have

$$a^+a^- \sim \mathrm{id}_x \qquad \qquad a^-a^+ \sim \mathrm{id}_y \tag{3.1}$$

We write P_1^{\sim} for the set of zig-zags up to congruence.

Lemma 25. The category P^{\sim} with P_0 as objects, P_1^{\sim} as morphisms, where composition is given by concatenation of paths up to congruence, is the free groupoid on P.

We have a canonical function $i_1 : \mathsf{P}_1 \to \mathsf{P}_1^{\sim}$, sending a rewriting step a to a^+ . Writing $s_0^{\sim}, t_0^{\sim} : \mathsf{P}_1^{\sim} \to \mathsf{P}_0$ for the source and target maps, it induces a morphism of ARS by taking

the identity on objects:

$$\begin{array}{c} \mathsf{P}_{0} \xleftarrow{s_{0}}{t_{0}} \mathsf{P}_{1} \\ \mathsf{id} \downarrow & \downarrow^{i_{1}} \\ \mathsf{P}_{0} \xleftarrow{s_{0}^{\sim}}{t_{0}^{\sim}} \mathsf{P}_{1}^{\sim} \end{array}$$

Writing $i : \mathsf{P} \to \mathsf{P}^{\sim}$ for this morphism of ARS, the universal property of P states that any morphism of ARS $F : \mathsf{P} \to \mathcal{C}$, where \mathcal{C} is a groupoid, extends uniquely as a functor $\tilde{F} : \mathsf{P}^{\sim} \to \mathcal{C}$ making the following diagram commute:



In the following, in order to avoid working with equivalence classes when working with elements of P_1^{\sim} , we will instead only implicitly consider zig-zags $a_1^{\epsilon_1}, \ldots, a_n^{\epsilon_n}$ which are *reduced*, in the sense that they satisfy the following property: for every index *i* with $1 \leq i < n$, we have that $a_i = a_{i+1}$ implies $\epsilon_i = \epsilon_{i+1}$. This is justified by the following result.

Lemma 26. The equivalence class under \sim of a zig-zag contains a unique reduced zig-zag.

Proof. Consider the string rewriting system on words over P_1^{\pm} with rules $a^+a^- \Rightarrow \mathrm{id}$ and $a^-a^+ \Rightarrow \mathrm{id}$ for $a \in \mathsf{P}_1$, corresponding to (3.1). It is length-reducing and thus terminating. Its critical pairs (whose sources are $a^-a^+a^-$ and $a^+a^-a^+$) are confluent, it is thus confluent. We deduce that any equivalence class contains a unique normal form, and those are precisely reduced zig-zags.

Given a path $p: x \xrightarrow{*} y$, we write $p^+: x \xrightarrow{\sim} y$ (resp. $p^-: y \xrightarrow{\sim} x$) for the zig-zag obtained by adding a "+" (resp. "-") exponent to every step of the rewriting path. In particular, the first operation induces a canonical inclusion $i_1^*: \mathsf{P}_1^* \to \mathsf{P}_1^\sim$, defined by $i_1^*(p) = p^+$, witnessing for the fact that rewriting paths are particular zig-zags. We will sometimes leave its use implicit in the following. Note that lemma 26 implies that i_1^* is injective.

We think here of ARS as abstractly describing some algebraic structures. It is thus natural to extend this notion in order to take in account the coherence laws that these structures should possess. This can be done as follows.

Definition 27. An extended abstract rewriting system, or 2-ARS, or 2-polygraph, P consists of an ARS as above, together with a set P_2 and two functions $s_1, t_1 : P_2 \to P_1^{\sim}$, such that

$$s_0^{\sim} \circ s_1 = s_0^{\sim} \circ t_1 \qquad \qquad t_0^{\sim} \circ s_1 = t_0^{\sim} \circ t_1$$

This data can be summarized as a diagram



In a 2-ARS, the elements of P_2 are *coherence relations* and the functions s_1 and t_1 respectively describe their source and target, which are rewriting zig-zags. We sometimes write

$$A:p \Rightarrow q$$

to indicate that $A \in \mathsf{P}_2$ is a coherence relation which admits p (resp. q) as source (resp. target), which can be thought of as a 2-cell

$$x \underbrace{A \Downarrow}_{q}^{p} y$$

where x (resp. y) is the common source (resp. target) of p and q.

Definition 28. The groupoid presented by a 2-ARS P, denoted by $\overline{\mathsf{P}}$, is the groupoid obtained from the free groupoid generated by the underlying ARS by quotienting morphisms under the smallest congruence identifying the source and the target of any element of P_2 .

More explicitly, the groupoid $\overline{\mathsf{P}}$ thus has P_0 as set of objects, the set P_1^{\sim} of rewriting zig-zags as morphisms, quotiented by the smallest equivalence relation \Leftrightarrow such that

$$p \cdot q \cdot r \Leftrightarrow^* p \cdot q' \cdot r$$

for every rewriting zig-zags p and r and coherence relation $A: q \Rightarrow q'$, which are suitably composable:

$$x' \xrightarrow{p} x \xrightarrow{q} y \xrightarrow{r} y'$$
(3.2)

We write \Leftrightarrow for the smallest symmetric relation identifying path $p \cdot q \cdot r$ and $p \cdot q' \cdot r$ when there is a coherence relation $A: q \Rightarrow q'$ as pictured above, so that $\stackrel{*}{\Leftrightarrow}$ is the reflexive transitive closure of \Leftrightarrow . Given a rewriting zig-zag $p \in \mathsf{P}_1^{\sim}$, we write \overline{p} for the corresponding morphism in $\overline{\mathsf{P}}$, i.e. its equivalence class under $\stackrel{*}{\Leftrightarrow}$. Given a zig-zag $p: x \to y$ in P^{\sim} , we write $\overline{p}: x \to y$ for its equivalence class.

Remark 29. A more categorical approach to the equivalences between zig-zags can be developed as follows, see [3] for details. A 2-groupoid is a 2-category whose 1- and 2-cells are invertible. A 2-ARS freely generates a 2-groupoid, whose underlying 1-groupoid is the one freely generated by the underlying 1-ARS of P, and containing the coherence relations as 2-cells. Given zig-zags $p, q: x \to y$, we then have $p \Leftrightarrow^* q$ if and only if there is a 2-cell $p \Rightarrow q$ in the free 2-groupoid: the 2-cells can thus be thought of as witnesses for the equivalences of zig-zags. We do not further detail this approach here, since it is not required, but it would be for instance needed if we were interested in higher coherence laws.

There are many 2-ARS presenting a given groupoid. In particular, one can always perform the following transformations on 2-ARS, while preserving the presented groupoid. Those are analogous to the transformations that one can perform on group presentations (while preserving the presented group) first studied by Tietze [64, 45].

Definition 30. The *Tietze transformations* are the following possible transformations on a 2-ARS P:

- (T1) given a zig-zag $p: x \xrightarrow{\sim} y$, add a new rewriting rule $a: x \to y$ in P_1 together with a new coherence relation $A: a \Rightarrow p$ in P_2 ,
- (T2) given zig-zags $p, q: x \xrightarrow{\sim} y$ such that $p \stackrel{*}{\Leftrightarrow} q$, add a new coherence relation $A: p \Rightarrow q$ in P_2 .

The *Tietze equivalence* is the smallest equivalence relation on 2-ARS identifying P and Q whenever Q can be obtained from P by a Tietze transformation (T1) or (T2).

It is easy to see that the Tietze transformations are "correct", in the sense that they preserve the presented groupoid. With more work [3, Chapter 5], it is even possible to show that those transformations are "complete", in the sense that any two 2-ARS presenting the same groupoid are Tietze equivalent. We only state the first direction here since this is the only one we will need:

Proposition 31. Any two Tietze equivalent 2-ARS present isomorphic groupoids.

From the transformations of definition 30, it is possible to derive other transformations, which also preserve the presented groupoid.

Lemma 32. Suppose that P is a 2-ARS containing a rewriting rule $a : x \to y$ and a relation $A : p \Rightarrow q$ such that a occurs exactly once in the source p, i.e. $p = p_1 \cdot a \cdot p_2$, and does not occur in the target:



Then P is Tietze equivalent to the 2-ARS where

- we have removed the rewriting rule a,
- we have removed the coherence relation A,
- we have replaced every occurrence of a in the source or target of a coherence relation by $p_1^- \cdot q \cdot p_2^-$.

3.2. Rewriting properties. Let P be a 2-ARS. For simplicity, we suppose that for every coherence relation $A: p \Rightarrow q$ in P₂, we have that p and q are rewriting paths (as opposed to zig-zags). We also suppose fixed a set $W \subseteq P_1$. We can think of W as inducing a rewriting subsystem W of P, with P₀ as objects, W as rewriting steps and

$$W_2 = \{A \in P_2 \mid s_1(A) \in W^* \text{ and } t_1(A) \in W^*\}$$

as coherence relations, and formulate the various traditional rewriting concepts with respect to it. In such a situation, consider the presented groupoid $C = \overline{\mathsf{P}}$. The set W of rewriting rules, induces a set of morphisms of C, namely $\{\overline{w} \mid w \in W\}$ that we still write W, which generates a subgroupoid W of C. Our aim here is to provide rewriting tools in order to show that W is rigid, so that C is equivalent to the quotient C/W by theorem 16, and moreover provide a concrete description of the quotient category.

Definition 33. The 2-ARS P is *W*-terminating if there is no infinite sequence a_1, a_2, \ldots of elements of W such that every finite prefix is a rewriting path in W^* .

Definition 34. An element $x \in \mathsf{P}_0$ is a *W*-normal form when there is no rewriting step in *W* with *x* as source. We say that P is weakly *W*-normalizing when for every $x \in \mathsf{P}_0$ there exists a normal form \hat{x} and a rewriting path $x \stackrel{*}{\to} \hat{x}$. In this case, we write $n_x : x \stackrel{*}{\to} \hat{x}$ for an arbitrary choice of such a path, which is however supposed to be the identity when *x* is a normal form.

Lemma 35. If P is W-terminating then it is weakly W-normalizing.

Proof. Consider a maximal rewriting path a_1, a_2, \ldots in W^* starting from x. Because P is W-terminating, this path is necessarily finite, and its target is a normal form by maximality.

Definition 36. A *W*-branching is a pair of rewriting paths

$$p_1: x \xrightarrow{*} y_1 \qquad \qquad q_2: x \xrightarrow{*} y_2$$

in W^* which are coinitial, i.e. have the same source x, which is called the *source* of the branching. A *W*-branching is *local* when both p_1 and p_2 are rewriting steps. A *W*-branching as above is *confluent* when there is a pair of cofinal (with the same target) rewriting paths $q_1: y_1 \xrightarrow{*} z$ and $q_2: y_2 \xrightarrow{*} z$ in W^* such that $p_1 \cdot q_1 \stackrel{*}{\Leftrightarrow} p_2 \cdot q_2$:



Note that, in the above definitions, not only we require that we can close a span of rewriting steps by a cospan of rewriting paths (as in the traditional definition of confluence), but also that the confluence square can be filled coherence relations.

Definition 37. The ARS P is *locally* W-confluent when W-branching is confluent. It is W-confluent when for every $p_1 : x \xrightarrow{*} y_1$ and $p_2 : x \xrightarrow{*} y_2$ in W*, there exist $q_1 : y_1 \xrightarrow{*} z$ and $q_2 : y_2 \xrightarrow{*} z$ in W* such that $p_1 \cdot q_1 \rightleftharpoons p_2 \cdot q_2$. We say that P is W-convergent when it is both W-terminating and W-confluent.

The celebrated Newman's lemma [53] (also sometimes called the diamond lemma) along with its traditional proof [63, Theorem 1.2.1 (ii)] easily generalizes to our setting:

Proposition 38. If P is W-terminating and locally W-confluent then it is W-confluent.

Proof. The relation on objects defined by $x \ge y$ whenever there exists a rewriting path $p: x \xrightarrow{*} y$ in W^* is a well-founded partial order because P is W-terminating. We say that P is W-confluent at x when every W-branching with source x confluent. We are going to show that P is locally W-confluent at x for every object x, by well-founded induction on x. In the base case, x is a W-normal form and the result is immediate. Otherwise, consider a W-branching consisting of paths $a_1 \cdot p_1$ and $a_2 \cdot p_2$ for some rewriting steps a_1 and a_2 and rewriting paths p_1 and p_2 (we suppose that the paths are non-empty, otherwise the result is immediate). The following diagram shows the W-confluence at x:



Above, the diagram LC is W-confluent by local confluence, and the two diagrams IH are by induction hypothesis.

Definition 39. The 2-ARS P is *W*-coherent if for any parallel zig-zags $p, q : x \xrightarrow{\sim} y$ in W^{\sim} , we have $p \stackrel{*}{\Leftrightarrow} q$.

The following is immediate:

Lemma 40. A 2-ARS P is W-coherent precisely when \overline{W} is a rigid subgroupoid of \overline{P} .

The traditional Church-Rosser property [63, Theorem 1.2.2] generalizes as follows in our setting:

Proposition 41. If P is weakly W-normalizing and W-confluent then for any zig-zag $p: x \xrightarrow{\sim} y$ in W^{\sim} , we have $\hat{x} = \hat{y}$ and $p \cdot n_y \stackrel{*}{\Leftrightarrow} n_x$, i.e. the diagram



commutes in $\overline{\mathsf{P}}$.

Proof. By confluence, given a rewriting path $p: x \xrightarrow{*} y$ in W^* , we have $\hat{x} = \hat{y}$ and $p \cdot n_y \stackrel{*}{\Leftrightarrow} n_x$, and thus $p^+ \cdot n_y \stackrel{*}{\Leftrightarrow} n_x$ and $n_x \cdot p^- \stackrel{*}{\Leftrightarrow} n_y$, i.e. the following diagrams commute in $\overline{\mathsf{P}}$:

Any zig-zag $p: x \xrightarrow{\sim} y$ in W^{\sim} decomposes as $p = p_1^- q_1^+ p_2^- p_2^+ \dots p_n^- p_n^+$ for some $n \in \mathbb{N}$ and paths p_i and q_i in W^* . We thus have $p \cdot n_y \stackrel{*}{\Leftrightarrow} n_x$, since all the squares of the following diagram commute in \overline{W} by the preceding remark:

$$\begin{array}{c} x \xrightarrow{p_1^-} y_1 \xrightarrow{q_1^+} x_2 \longrightarrow \cdots \longrightarrow x_n \xrightarrow{p_n^-} y_n \xrightarrow{q_n^-} y \\ n_x \downarrow \qquad n_{y_1}^{\downarrow} \qquad n_{x_2}^{\downarrow} \qquad n_{x_n}^{\downarrow} \qquad n_{y_n}^{\downarrow} \qquad n_{y_n}^{\downarrow} \qquad n_{y_n}^{\downarrow} \\ \hat{x} = \hat{x} = \hat{x} = \cdots = \hat{x} = \cdots = \hat{x} = \hat{x} \end{array}$$

which allows us to conclude.

This implies the following "abstract" variant of Squier's homotopical theorem [23, 37, 60]:

Proposition 42. If P is weakly W-normalizing and W-confluent then it is W-coherent.

Proof. Given two parallel zig-zags $p, q: x \xrightarrow{\sim} y$ in W^{\sim} , we have $p \stackrel{*}{\Leftrightarrow} q$, since the following diagram commutes in $\overline{\mathsf{P}}$:



Namely, we have $\hat{x} = \hat{y}$ by confluence, the two triangles above commute by proposition 41, and the two triangles below do because n_y^- is an inverse for n_y .

Example 43. As a variant of example 17, consider the 2-ARS P with $P_0 = \{x, y\}$, $P_1 = \{a, b : x \to y\}$ and $P_2 = \emptyset$, i.e. $x \xrightarrow[b]{a} y$. With $W = \{a\}$, we have that P is W-terminating and locally W-confluent, thus W-confluent by proposition 38, and thus W-coherent by lemma 35 and proposition 42. With $W = \{a, b\}$, we have seen in example 17 that the groupoid \overline{W} is not rigid and, indeed, P is not W-confluent because $\overline{a} \neq \overline{b}$ (because $P_2 = \emptyset$).

Definition 44. We write $N(\overline{P})$ for the *category of normal forms* of \overline{P} , defined as the full subcategory of \overline{P} whose objects are those in *W*-normal form.

Lemma 45. If P is weakly W-normalizing, then the inclusion functor $N(\overline{P}) \to \overline{P}$ is an equivalence of categories.

Proof. An object x of P admits a normal form \hat{x} , by lemma 35. Writing $n_x : x \xrightarrow{*} \hat{x}$ for a normalization path, we have an isomorphism $\overline{n_x} : x \to \hat{x}$ in $\overline{\mathsf{P}}$. The inclusion functor is thus full and faithful (by definition), and every object of $\overline{\mathsf{P}}$ is isomorphic to an object in the image, it is thus an equivalence of categories.

When P is W-convergent, the equivalence given in the above lemma is precisely the one with the quotient category:

Proposition 46. If P is W-convergent, then the quotient category \overline{P}/W is isomorphic to the category of normal forms $N(\overline{P})$.

Proof. Since P is W-convergent, by proposition 42 and lemma 40, the groupoid generated by W is rigid and we thus have the description of the quotient $\overline{\mathsf{P}}/W$ given by proposition 10. We have a canonical functor $N(\overline{\mathsf{P}}) \to \overline{\mathsf{P}}/W$, obtained as the composite of the inclusion functor $N(\overline{\mathsf{P}}) \to \overline{\mathsf{P}}$ with the quotient functor $\overline{\mathsf{P}} \to \overline{\mathsf{P}}/W$. An object of $\overline{\mathsf{P}}/W$ is an equivalence class [x] of objects which, by convergence, contains a unique normal form, namely \hat{x} . The functor is bijective on objects. By weak normalization (lemma 35), any morphism $f: x \to y$ is equivalent to one with both normal source and target, namely $\hat{f} = n_y \circ f \circ n_x^- : \hat{x} \to \hat{y}$, hence the functor is full. Consider two morphisms $f, g: \hat{x} \to \hat{y}$ in N(P) with the same image [f] = [g]: by definition of the equivalence on morphisms, there exist morphisms $v: \hat{x} \to \hat{x}$ and $w: \hat{y} \to \hat{y}$ in W^{\sim} making the diagram

$$\begin{array}{ccc} \hat{x} & \stackrel{f}{\longrightarrow} \hat{y} \\ v \downarrow & & \downarrow^{u} \\ \hat{x} & \stackrel{g}{\longrightarrow} \hat{y} \end{array}$$

commute. By the Church-Rosser property (proposition 42), we have $v = n_{\hat{x}} \circ n_{\hat{x}}^-$ and thus $v = \mathrm{id}_x$ (since $n_{\hat{x}} = \mathrm{id}_{\hat{x}}$ by hypothesis), and similarly $w = \mathrm{id}_y$. Hence f = g and the functor is faithful. The functor is thus an isomorphism as being full, faithful and bijective on objects.

We can finally summarize the results obtained in this section as follows. Given a 2-ARS P and a set $W \subseteq P_1$, we have the following possible reasonable definitions of the fact that P is *coherent* wrt W:

- (1) Every parallel zig-zags with edges in W are equal
- (i.e. the subgroupoid of \overline{P} generated by W is rigid).
- (2) The quotient map $\overline{\mathsf{P}} \to \overline{\mathsf{P}}/W$ is an equivalence of categories.
- (3) The inclusion $N(\overline{P}) \to \overline{P}$ is an equivalence.
- (4) The inclusion $\operatorname{Alg}(\overline{\mathsf{P}}/W, \mathcal{D}) \to \operatorname{Alg}(\overline{\mathsf{P}}, \mathcal{D})$ is a natural equivalence of categories.

Theorem 47. If P is W-convergent then all the above coherence properties hold.

Proof. (1) is given by proposition 42, (2) is given by (1) and theorem 16, (3) is given by proposition 46, and (4) is given by (1) and proposition 21.

3.3. Presenting the groupoid of normal forms. We would now like to provide an explicit description of $N(\overline{P})$, by a 2-ARS. A good candidate is the following 2-ARS $P \setminus W$ obtained by "restricting P to normal forms". More precisely, it consists of

- $(\mathsf{P} \setminus W)_0$: the objects of $\mathsf{P} \setminus W$ are those of P in W-normal form,
- $(\mathsf{P} \setminus W)_1$: the rewriting rules of $\mathsf{P} \setminus W$ are those of P whose source and target are both in $(\mathsf{P} \setminus W)_0$ (in particular, it does not contain any element of W, thus the notation),
- $(P \setminus W)_2$: the coherence relations are those of P_2 whose source and target both belong to $(\mathsf{P} \setminus W)_1^{\sim}$.

It is not the case in general that this 2-ARS presents the category $N(\overline{P})$, but we will provide here conditions which ensure that this holds, see also [15, 51] for alternative conditions. For simplicity, we suppose here that the source and target of every rewriting step in P₂ is a path (as opposed to a zig-zag).

Proposition 48. Suppose that

- (1) P is W-convergent,
- (2) every rule $a: x \to y$ in P_1 whose source x is W-normal also has a W-normal target y,
- (3) for every coinitial rule $a: x \to y$ in P_1 and path $w: x \xrightarrow{*} x'$ in W^* , there are paths $p: x' \xrightarrow{*} y'$ in P_1^* and $w': y \xrightarrow{*} y' \in W^*$ such that $a \cdot w' \stackrel{*}{\Leftrightarrow} w \cdot p$:

(4) for every coherence relation $A: p \Rightarrow q: x \stackrel{*}{\rightarrow} y$, and for every path $w: x \stackrel{*}{\rightarrow} x'$, the paths $p', q': x' \stackrel{*}{\rightarrow} y'$ in P_1^* and $w': y \stackrel{*}{\rightarrow} y'$ in W^* such that $p \cdot w' \stackrel{*}{\Leftrightarrow} w \cdot p'$ and $q \cdot w' \stackrel{*}{\Leftrightarrow} w \cdot q'$ induced by (3) satisfy $p' \stackrel{*}{\Leftrightarrow} q'$:



Then $N(\overline{P})$ is isomorphic to $\overline{P \setminus W}$.

Proof. We write $\mathbf{Q} = \mathbf{P} \setminus W$. Since \mathbf{Q} is, by definition, a sub-2-ARS of \mathbf{P} there is a canonical functor $\overline{\mathbf{Q}} \to \overline{\mathbf{P}}$. Moreover, since the objects of \mathbf{P} are, by definition, in *W*-normal form, this functor corestricts to a functor $F : \overline{\mathbf{Q}} \to \mathbf{N}(\overline{\mathbf{P}})$ which is the identity on objects.

First, note that condition (2) implies that for any path p of the form

$$x_0 \xrightarrow{a_1} x_1 \xrightarrow{a_2} \cdots \xrightarrow{a_n} x_n$$

with x_0 in W-normal form, we have that all the x_i are in W-normal form and thus p belongs to \mathbb{Q}_1^* . Similarly, every coherence relation $A: p \Rightarrow q: x \stackrel{*}{\to} y$ with x in W-normal form belongs to \mathbb{Q}_2 .

We claim that for every zig-zag $p: x \xrightarrow{\sim} y$ in P_1^{\sim} there is zig-zag $q \in \mathsf{Q}_1^{\sim}$ such that $p \Leftrightarrow n_x \cdot q \cdot n_y^{-}$. We have that p is of the form $p = w_0 \cdot a_1 \cdot w_1 \cdot a_2 \cdot w_2 \cdot \ldots \cdot a_n \cdot w_n$ where the a_i are rules in P_1 which are not in W (possibly taken backward) and the w_i are in W^{\sim} . For instance, consider the case n = 1 and a path p of the form $p = v \cdot a \cdot w$ with $a \in \mathsf{Q}_1$ and $v, w \in W^{\sim}$ (the case where a is reversed is similar, and the general case follows by induction):



By hypothesis (1) and proposition 41, we have $v \Leftrightarrow n_x \cdot n_{x'}^-$ and $w \Leftrightarrow n_{y'} \cdot n_y^-$. By hypothesis (3), there exist paths $q : \hat{x} \Leftrightarrow y''$ in P_1^* and $w' : y' \Leftrightarrow y''$ in W^* such that $a \cdot w' \Leftrightarrow n_{x'} \cdot q$. By hypothesis (2) and the remark at the beginning of this proof, we have that $q \in \mathsf{Q}_1^*$ and y'' is a normal form. By (1), we thus have $y'' = \hat{y}$ and $w' \Leftrightarrow n_{y'}$, and we conclude. As a particular case of the property we have just shown, for any zig-zag $p : x \to y$ whose source and target are in *W*-normal form, we have that that p is equivalent to a zig-zag $q : x \to y$ (since in this case both n_x and n_y are identities). The functor $F : \overline{\mathsf{Q}} \to \mathsf{N}(\overline{\mathsf{P}})$ is thus full.

In the following, given a path $p: x \to y$ in P_1^* , we write $\hat{p}: \hat{x} \to \hat{y}$ in Q_1^* for a path such that $p = n_x \cdot \hat{p} \cdot n_y^-$. Such a path always exists by the previous reasoning and can be constructed in a functorial way (i.e. $\hat{p_1} \cdot \hat{p_2} = \hat{p_1} \cdot \hat{p_2}$). Now, suppose given two zig-zags $p, p': x \to y$ in P_1^- such that $p \Leftrightarrow p'$. The relation $p \Leftrightarrow p'$ means that there is a sequence p_1, \ldots, p_n of zig-zags in P_1^- such that $p_1 = p, p_n = p'$ and each p_i is related to p_{i+1} by taking a relation in context, as in (3.2):

$$p = p_1 \Leftrightarrow p_2 \Leftrightarrow \ldots \Leftrightarrow p_n = p'$$

More formally, for $1 \leq i < n$, there is a decomposition

$$p_i = q_i \cdot r_i \cdot s_i \qquad \qquad p_{i+1} = q_i \cdot r'_i \cdot s_i$$

such that there is a relation $A: r_i \Rightarrow r'_i$ or $A: r'_i \Rightarrow r_i$ (another approach would consist in reasoning by induction on the 2-cells of the freely generated 2-groupoid of remark 29). By hypothesis (4), there is a relation $\hat{r}_i \stackrel{*}{\Leftrightarrow} \hat{r}'_i$, and thus $\hat{p}_i \stackrel{*}{\Leftrightarrow} \hat{p}_{i+1}$ by functoriality. By recurrence on n, we thus have $\hat{p} \stackrel{*}{\Leftrightarrow} \hat{p}'$. From this, we deduce that that the functor F is also faithful. In practice, condition (1) can be shown using traditional rewriting techniques (e.g. proposition 38) and condition (2) is easily checked by direct inspection of the rewriting rules. We provide below sufficient conditions in order to show the two remaining conditions:

Proposition 49. We have the following.

(3) Suppose that for every coinitial rules $a : x \to y$ in P_1 and $w : x \to x'$ in W, there are paths $p : x' \stackrel{*}{\to} y'$ in P_1^* and $w' : y \stackrel{*}{\to} y'$ in W^* such that w is of length at most one and $a \cdot w' \stackrel{*}{\Leftrightarrow} w \cdot p$:

The condition (3) of proposition 48 is satisfied.

(4) Suppose that condition (3) is satisfied and for every coherence relation A : p ⇒ q : x → y in P₂ and rule w : x → x' in W, the paths p', q' : x' → y' in P₁^{*} and w' : y → y' in W^{*} of length at most one such that p · w' ⇔ w · p' and q · w' ⇔ w · q' induced by (3) are such that p' ⇔ q':

Then condition (4) of proposition 48 is satisfied.

Proof. Both properties are easily shown by recurrence on the length of w.

4. Relative coherence for Lawvere theories

In order to use the developments of section 3 in concrete situations, such as (symmetric) monoidal categories, we need to consider a more structured notion of theory. For this reason, we consider here Lawvere 2-theories, as well as the adapted notion of rewriting, which is a coherent extension of the traditional notion of term rewriting systems.

4.1. Lawvere 2-theories. We begin by recalling the traditional notion due to Lawvere [42]:

Definition 50. A Lawvere theory \mathcal{T} is a cartesian category, with \mathbb{N} as set of objects, and cartesian product given on objects by addition. A morphism between Lawvere theories is a product-preserving functor and we write \mathbf{Law}_1 for the category of Lawvere theories.

For simplicity, we restrict here to unsorted theories, but the developments performed here could easily adapted to the multi-sorted case. In such a theory, we usually restrict our attention to morphisms with 1 as codomain, since $\mathcal{T}(n,m) \cong \mathcal{T}(n,1)^m$ by cartesianness.

A (2, 1)-category is a 2-category in which every 2-cell is invertible, i.e. a category enriched in groupoids. The following generalization of Lawvere theories was introduced in various places, see [20, 66, 67], as well as [56] for the enriched point of view: **Definition 51.** A Lawvere 2-theory \mathcal{T} is a cartesian (2, 1)-category with \mathbb{N} as objects, and cartesian product given on objects by addition. A morphism $F : \mathcal{T} \to \mathcal{U}$ between 2-theories is a 2-functor which preserves products. We write \mathbf{Law}_2 for the resulting category (which can be extended to a 3-category by respectively taking natural transformations and modifications as 2- and 3-cells).

We can reuse the properties developed in sections 2 and 3 by working "hom-wise" as follows. Let \mathcal{T} be a 2-theory together with a subset W of the 2-cells. We write \mathcal{W} for the sub-2-theory of \mathcal{T} , with the same 0- and 1-cells, and whose 2-cells contain W (we often assimilate this 2-theory to its set of 2-cells). A morphism $F: \mathcal{T} \to \mathcal{U}$ of Lawvere 2-theories is W-strict when it sends every 2-cell in W to an identity.

Definition 52. The quotient 2-theory \mathcal{T}/W is the theory equipped with a W-strict morphism $\mathcal{T} \to \mathcal{T}/W$ such that every W-strict morphism $F : \mathcal{T} \to \mathcal{U}$ extends uniquely as a morphism $\mathcal{T}/W \to \mathcal{U}$:



Such a quotient 2-theory always exists by general arguments: 2-theories form a locally presentable category, which therefore has colimits, and thus coequalizers [57]. We have $\mathcal{T}/W \cong \mathcal{T}/W$, so that we can always assume that we are quotienting by a sub-2-theory. On hom-categories, the quotient corresponds to the one introduced in section 2.1:

Lemma 53. For every $m, n \in \mathbb{N}$, we have

$$(\mathcal{T}/\mathcal{W})(m,n) = \mathcal{T}(m,n)/\mathcal{W}(m,n)$$

We say that a morphism

 $F:\mathcal{T}\to\mathcal{U}$

is a *local equivalence* when for every objects $m, n \in \mathcal{T}$, the induced functor

$$F_{m,n}: \mathcal{T}(m,n) \to \mathcal{U}(m,n)$$

between hom-categories is an equivalence.

Definition 54. A theory \mathcal{W} is 2-*rigid* when any two parallel 2-cells are equal.

Note that a theory \mathcal{W} is 2-rigid if and only if the category $\mathcal{W}(m, n)$ is rigid for every 0-cells m and n. By direct application of theorem 16, we have

Theorem 55. The quotient 2-functor $\mathcal{T} \to \mathcal{T}/\mathcal{W}$ is a local equivalence iff \mathcal{W} is 2-rigid.

4.2. Algebras for Lawvere 2-theories. The notion of algebra for 2-theories was extensively studied by Yanofsky [66, 67], we refer to his work for details.

Definition 56. An algebra for a Lawvere 2-theory \mathcal{T} is a 2-functor $C : \mathcal{T} \to \mathbf{Cat}$ which preserves products. By abuse of notation, we often write C instead of C1 and suppose that products are strictly preserved, so that $Cn = C^n$.

Definition 57. A pseudo-natural transformation $F : C \Rightarrow D$ between algebras C and D consists in a functor $F : C \to D$ together with a family $\phi_f : Df \circ F^n \Rightarrow F \circ Cf$ of natural transformations indexed by 1-cells $f : n \to 1$ in \mathcal{T} ,

$$\begin{array}{ccc} C^n & \xrightarrow{Cf} & C \\ F^n & & & \downarrow F \\ D^n & \xrightarrow{Df} & D \end{array}$$

which is compatible with products, composition and 2-cells of \mathcal{T} .

Definition 58. A modification $\mu : F \Rightarrow G : C \Rightarrow D$ between two pseudo-natural transformations is a natural transformation $\mu : F \Rightarrow G$ which is compatible with 2-cells of \mathcal{T} . We write $\operatorname{Alg}(\mathcal{T})$ for the 2-category of algebras of a 2-theory \mathcal{T} , pseudo-natural transformations and modifications.

Example 59. Consider the 2-TRS Mon of example 70 below. The 2-category Alg(Mon) of algebras of the presented 2-theory is isomorphic to the 2-category **MonCat** of monoidal categories, strong monoidal functors and monoidal natural transformations. It might be surprising that Mon has five coherence relations whereas the traditional definition of monoidal categories only features two axioms, which correspond to the coherence relations A and C. There is no contradiction here: the commutation of the two axioms can be shown to imply the one of the three other [22, 29].

Similarly, writing $\mathcal{W} = \overline{\text{Mon}}$, the 2-category Alg($\overline{\text{Mon}}/\mathcal{W}$) is the sub-2-category of Alg($\overline{\text{Mon}}$) whose algebras are monoidal categories where the coherence natural transformations are identities, i.e. the 2-category **MonCat**_{str} of strict monoidal categories.

We conjecture that one can generalize the classical proof that any monoidal category is monoidally equivalent to a strict one [46, Theorem XI.3.1] to show the following general (C3) coherence theorem, as well as its (C4) generalization:

Conjecture 60 (C3). When \mathcal{W} is 2-rigid, every \mathcal{T} -algebra is equivalent to a \mathcal{T}/\mathcal{W} algebra.

Conjecture 61 (C4). When \mathcal{W} is 2-rigid, the 2-functor $\operatorname{Alg}(\mathcal{T}/\mathcal{W}) \to \operatorname{Alg}(\mathcal{T})$ induced by precomposition with the quotient 2-functor $\mathcal{T} \to \mathcal{T}/\mathcal{W}$ has a left adjoint such that the components of the unit are equivalences.

By example 59, the conjectures would allow recovering the coherence theorem for monoidal categories by taking $\mathcal{W} = \mathcal{T} = \overline{\text{Mon}}$. We will see in section 6 that the conjectures would also allow recovering the coherence theorem for symmetric monoidal categories by taking $\mathcal{T} = \overline{\text{SMon}}$ to be the 2-theory presented by a suitable 2-TRS, and \mathcal{W} the subtheory generated by the generating 2-cells corresponding to α , λ and ρ . One should also be able to obtain the coherence theorems for braided monoidal categories in a similar way.

A detailed study of those conjectures is left for future works, since it would require introducing some more categorical material, and our aim in this article is to focus on the rewriting techniques. Note that, apart from informal explanations, we could not find a proof of conjectures 60 and 61 for symmetric or braided monoidal categories in the literature, e.g. in [28, 46, 47] (in [28, Theorem 2.5] the result is only shown for free braided monoidal categories). Once proved, it would be interesting to investigate whether the converse implications hold, i.e. whether the conditions imply \mathcal{W} being 2-rigid. Remark 62. One could hope for the following alternative generalization of proposition 19: a 2-functor $F : \mathcal{T} \to \mathcal{T}'$ between theories is a biequivalence if and only if the functor $\operatorname{Alg}(F) : \operatorname{Alg}(\mathcal{T}') \to \operatorname{Alg}(\mathcal{T})$ induced by precomposition is an equivalence. This is claimed in [67, Proposition 7], along with the corollary that the categories $\operatorname{Alg}(\mathcal{T}/\mathcal{W})$ and $\operatorname{Alg}(\mathcal{T})$ are equivalent when \mathcal{W} is 2-rigid (at least in the particular case where \mathcal{T} is the theory of monoids and $\mathcal{W} = \mathcal{T}$, as in example 59). However, both proofs are based on the wrong claim that biessentially surjective local equivalences coincide with biequivalences [67, Proposition 6].

We recall that a 2-functor $F : \mathcal{C} \to \mathcal{D}$ between 2-categories is

- biessentially surjective when for every 0-cell x of C there is a 0-cell y of \mathcal{D} together with an equivalence $Fx \simeq y$,
- a local equivalence when for every 0-cells x, y of C, the induced functor $C(x, y) \to C(Fx, Fy)$ is an equivalence,
- a biequivalence when there is a 2-functor $G : \mathcal{D} \to \mathcal{C}$ together with natural equivalences $G \circ F \simeq \mathrm{Id}_{\mathcal{C}}$ and $F \circ G \simeq \mathrm{Id}_{\mathcal{D}}$.

Any biequivalence is a biessentially surjective local equivalence, but the converse is not true: from a biessentially surjective local equivalence $F : \mathcal{C} \to \mathcal{D}$, one can in general only construct a pseudofunctor (as opposed to a 2-functor) $G : \mathcal{D} \to \mathcal{C}$ satisfying the desired properties. A concrete counter-example is given in [34, Example 3.1], as we now recall. Consider the 2-categories

- \mathcal{C} with one 0-cell \star , with $(\mathbb{N}, +, 0)$ as monoid of endomorphisms on \star , and one 2-cell between 1-cells $m, n \in \mathbb{N}$ whenever m and n have the same parity,
- \mathcal{D} with one 0-cell \star , with $(\mathbb{N}/2\mathbb{N}, +, 0)$ as monoid of endomorphisms on \star , and only trivial 2-cells.

In other terms, the 2-category C is freely generated by one 0-cell \star , one 1-cell 1 and 2-cells are the congruence generated by 1 + 1 = 0:



and the category \mathcal{D} is obtained from \mathcal{C} by formally turning 2-cells into identities. We thus have a quotient 2-functor $F : \mathcal{C} \to \mathcal{D}$ sending a 1-cell $n \in \mathbb{N}$ to 0 or 1 depending on whether nis even or odd, i.e. F(2n) = 0 and F(2n+1) = 1. Conversely, the functor $G : \mathcal{D} \to \mathcal{C}$ should associate to every 1-cell of \mathcal{D} a representative, i.e. G(0) = 2m and G(1) = 2n + 1 for some $m, n \in \mathbb{N}$. Since we require that G is functorial, we should have

$$2m = G(0) = G(0+0) = G(0) + G(0) = 4m$$

so that m = 0, and

$$0 = G(0) = G(1+1) = G(1) + G(1) = 2n + 2$$

thus reaching a contradiction. The morale is that a choice of representatives is usually not strictly functorial, but only pseudo-functorial (above, 0 and 2n + 2 are not equal, but they are in the same equivalence class).

Whether the above generalization of proposition 19 holds is left open. However, we cannot use it to conclude that, when $\mathcal{W} \subseteq \mathcal{T}$ is 2-rigid, we have that the canonical functor $\operatorname{Alg}(\mathcal{T}/\mathcal{W}) \to \operatorname{Alg}(\mathcal{T})$ is an equivalence, because the functor $\mathcal{T} \to \mathcal{T}/\mathcal{W}$ is in general a biessentially surjective local equivalence (theorem 55), but not a biequivalence.

5. Coherent term rewriting systems

5.1. Extended rewriting systems. We now recall the categorical setting for term rewriting systems, as well as their extension in order to handle coherence. A more detailed presentation can be found in [3, 8, 16, 48].

Definition 63. A signature consists in a set S_1 of symbols together with a function $s_0 : S_1 \to \mathbb{N}$ associating to each symbol an arity, and we write $a : n \to 1$ for a symbol a of arity n. A morphism of signatures is a function between the corresponding sets of symbols which preserves arity, and we write \mathbf{Pol}_1^{\times} for the corresponding category.

Remark 64. If we were interested in the multi-sorted case, our signature would rather consist in a set S_1 together with a set S_0 of *sorts*, along with functions $s_0 : S_1 \to S_0^*$ (where S_0^* is the free monoid on S_0) and $t_0 : S_1 \to S_0$, respectively indicating the sorts of the inputs and of the output. The above definition can be recovered as the particular case where S_0 is the terminal set. This point of view explains why the index fo S_1 is 1.

There is a forgetful functor $\mathbf{Law}_1 \to \mathbf{Pol}_1^{\times}$, sending a theory \mathcal{T} to the set $\bigsqcup_{n \in \mathbb{N}} \mathcal{T}(n, 1)$ with first projection as arity. This functor admits a left adjoint $-^* : \mathbf{Pol}_1^{\times} \to \mathbf{Law}_1$, which we now describe. Given a signature S_1 , and $n \in \mathbb{N}$, $S_1^*(n, 1)$ is the set of *terms* of arity n: those are formed using operations, with variables in $\{x_1^n, x_2^n, \ldots, x_n^n\}$. Note that the superscript for variables is necessary to unambiguously recover the type of a variable, i.e. $x_i^n : n \to 1$, but for simplicity we will often omit it in the following. More explicitly, the family of sets $S_1^*(n, 1)$ is the smallest one such that

- for $1 \leq i \leq n$, we have

$$x_i^n \in \mathsf{S}_1^*(n,1)$$

- given $m, n \in \mathbb{N}$, a symbol $a: n \to 1$ and terms $t_1, \ldots, t_n \in S_1^*(m, 1)$, we have

$$a(t_1,\ldots,t_m) \in \mathsf{S}_1^*(m,1)$$

More generally, a morphism f in $S_1^*(n,m)$ is an *m*-uple

$$f = \langle t_1, \ldots, t_m \rangle$$

of terms t_i with variables in $\{x_1^n, \ldots, x_n^n\}$, which can be thought of as a formal substitution. Given such a substitution f and a term t, we write

$$t[f]$$
 or $t[t_1/x_1,\ldots,t_n/x_n]$

for the term obtained from t by formally replacing each variable x_i^n by t_i . This operation is thus defined inductively by

$$x_i^n[f] = t_i$$
 $a(u_1, \dots, u_k)[f] = a(u_1[f], \dots, u_k[f])$

The composition of two morphisms $\langle t_1, \ldots, t_m \rangle : S_1^*(n, m)$ and $\langle u_1, \ldots, u_k \rangle : S_1(m, k)$ is given by parallel substitution:

$$\langle u_1, \dots, u_k \rangle \circ \langle t_1, \dots, t_m \rangle = \langle u_1[t_1/x_1, \dots, t_n/x_n], \dots, u_k[t_1/x_1, \dots, t_m/x_m] \rangle$$

and the identity in $S_1^*(n, n)$ is $\langle x_1^n, \ldots, x_n^n \rangle$. The resulting category S_1^* is easily checked to be a Lawvere theory, which satisfies the following universal property:

Lemma 65. The Lawvere theory S_1^* is the free Lawvere theory on the signature S_1 .

By abuse of notation, we sometimes write

$$\mathsf{S}_1^* = \bigsqcup_{m,n \in \mathbb{N}} \mathsf{S}_1^*(m,n)$$

for the set of all substitutions and $s_0^*, t_0^* : S_1^* \to \mathbb{N}$ for the source and target maps, and $i_1 : S_1 \to S_1^*$ for the map sending an operation $a : n \to 1$ to the substitution consisting of one term $\langle a(x_1^n, \ldots, x_n^n) \rangle$, so that we have $s_0^* \circ i_1 = s_0$ and $t_0^* \circ i_1 = 1$.

Definition 66. A term rewriting system, or TRS, S consists of a signature S_1 together with a set S_2 of rewriting rules and functions $s_1, t_1 : S_2 \to S_1^*$ which indicate the source and target of each rewriting rule, and are supposed to satisfy

$$s_0^* \circ s_1 = s_0^* \circ t_1$$
 $t_0^* \circ s_1 = t_0^* \circ t_1 = 1$

This data can be summarized in the following diagram:



We sometimes write

$$\rho: t \Rightarrow u$$

for a rule ρ with t as source and u as target. The relations satisfied by any TRS ensure that both t and u have the same arity.

We now need to introduce some notions in order to be able to define rewriting in this setting. In case it helps, those are illustrated in example 70 below. A *context* C of arity n is a term with variables in $\{x_1, \ldots, x_n, \Box\}$ where the variable \Box is a particular variable, the *hole*, occurring exactly once. Here, we define the number $|t|_i$ of occurrences of a variable x_i (and similarly for \Box) in a term t by induction by

$$|x_i|_i = 1$$
 $|a(t_1, \dots, t_n)|_i = \sum_{k=1}^n |t_k|_i$

We write S_n^{\Box} for the set of contexts of arity n. Given a context C and a term t, both of same arity n, we write C[t] for the term obtained from C by replacing \Box by t. The composition of contexts C and D is given by substitution

$$D \circ C = D[C]$$

This composition is associative and admits the identity context \Box as neutral element. A *bicontext* from *n* to *k*, is a pair (C, f) consisting of a context *C* of arity *n* and a substitution $f \in S_1^*(n, k)$. This data can be thought of as the specification of a function on terms

$$\begin{aligned} \mathsf{S}_1^*(n,1) &\to \mathsf{S}_1^*(k,1) \\ &\langle t \rangle \mapsto C[\langle t \rangle \circ f] \end{aligned}$$

In the following, for simplicity, we will omit the brackets and simply write t instead of $\langle t \rangle$ in such an expression, so that the image of the function can also be denoted $C[t \circ f]$. This function will be referred as the *action* of a bicontext on terms. The composition

of bicontexts (C, f) and (D, g) of suitable types is given by $(D \circ C, f \circ g)$. The action is compatible with this composition, in the sense that we have

$$D[C[-\circ f] \circ g] = (D \circ C[f])[-\circ (f \circ g)]$$

A rewriting step of arity n

$$C[\rho \circ f] : C[t \circ f] \Rightarrow C[u \circ f]$$

is a triple consisting of

- a rewriting rule $\rho: t \Rightarrow u$, with t and u of arity k,
- a context C of arity n,
- a substitution $f: n \to k$ in S_1^* .

A rewriting step can thus be thought of as a rewriting rule in a bicontext. Its source is the term $C[t \circ f]$ and its target is the term $C[u \circ f]$. We write $S_2^{[]}$ for the set of rewriting steps. A rewriting path π is a composable sequence

$$C_1[t_1 \circ f_1] \xrightarrow{C_1[\rho_1 \circ f_1]} C_1[u_1 \circ f_1] = C_2[t_2 \circ f_2] \xrightarrow{C_2[\rho_2 \circ f_2]} \cdots \xrightarrow{C_n[\rho_n \circ f_n]} C_n[u_n \circ f_n]$$

of rewriting steps. We write S_2^* for the set of rewriting paths and adopt the previous notation, e.g. we write $\pi \cdot \pi'$ for the concatenation of two composable rewriting paths π and π' . As in section 3, we can also define a notion of *rewriting zig-zag* which is similar to rewriting paths excepting that some rewriting steps may be taken backwards, and write S_2^{\sim} for the corresponding set. We sometimes write $\pi : t \stackrel{*}{\Rightarrow} u$ (resp. $\pi : t \stackrel{\sim}{\Rightarrow} u$) to indicate the source and target or a rewriting path π .

Given a signature S_1 , there is a forgetful functor from the category of Lawvere 2-theories with S_1^* as underlying Lawvere theory to the category rewriting systems with S_1 as signature (with the expected notion of morphism).

Lemma 67. Given a TRS S, the Lawvere 2-theory S^{\sim} with S_1^* as 1-cells and S_2^{\sim} as 2-cells is free on S.

The action of bicontexts on terms extend to rewriting steps as follows. Given a rewriting step

$$C[\rho \circ f]: C[t \circ f] \Rightarrow C[u \circ f]$$

a context D and a substitution g of suitable types, we define $D[C[\rho \circ f] \circ g]$ to be the rewriting step

$$(D \circ C[g])[\rho \circ (f \circ g)] : (D \circ C[g])[t \circ (f \circ g)] \Rightarrow (D \circ C[g])[u \circ (f \circ g)]$$

Moreover, we extend this action to rewriting paths and zig-zags by functoriality, i.e.

$$C[(p \cdot q) \circ f] = C[p \circ f] \cdot C[q \circ f]$$

Definition 68. An extended term rewriting system, or 2-TRS, consists of a term rewriting system as above, together with a set S_3 of coherence relations and functions $s_2, t_2 : S_3 \to S_2^{\sim}$, indicating their source and target, satisfying

$$s_1^{\sim} \circ s_2 = s_1^{\sim} \circ t_2 \qquad \qquad t_1^{\sim} \circ s_2 = t_1^{\sim} \circ t_2$$

Diagrammatically,



Given a 2-TRS as above, we sometimes write

 $A:\pi \Rrightarrow \pi'$

to indicate that A is a coherence relation in S_3 with π as source and π' as target. Given two rewriting paths π and π' , we write $\pi \Leftrightarrow \pi'$ when they are related by the smallest congruence identifying the source and target of any coherence relation.

Definition 69. The Lawvere 2-theory *presented* by a 2-TRS S is the (2, 1)-category noted \overline{S} , with \mathbb{N} as 0-cells, S_1^* as 1-cells and the quotient of S_2^{\sim} under the congruence $\stackrel{*}{\Leftrightarrow}$ as 2-cells.

Example 70. The extended rewriting system Mon for monoids has symbols and rules

$$\mathsf{Mon}_1 = \{m : 2 \to 1, e : 0 \to 1\}$$
$$\mathsf{Mon}_2 = \left\{\begin{array}{l} \alpha : m(m(x_1, x_2), x_3) \Rightarrow m(x_1, m(x_2, x_3))\\ \lambda : m(e, x_1) \Rightarrow x_1\\ \rho : m(x_1, e) \Rightarrow x_1 \end{array}\right\}$$

There are coherence relations A, B, C, D and E, respectively corresponding to a confluence for the five critical pairs of the rewriting system (as defined below), whose 0-sources are respectively

$$m(m(m(x_1, x_2), x_3), x_4) \quad m(m(e, x_1), x_2) \quad m(m(x_1, e), x_2) \quad m(m(x_1, x_2), e) \quad m(e, e)$$

Those coherence relations can be pictured as follows:

For concision, for each arrow, we did not indicate the proper rewriting step, but only the rewriting rule of the rewriting step (hopefully, the reader will easily be able to reconstruct the missing bicontext). For instance, the coherence relation C has type

$$C: m(\rho(x_1), x_2) \Rightarrow \alpha(x_1, e, x_2) \cdot m(x_1, \lambda(x_2))$$

so that the missing bicontexts for the rules labeled by ρ , α and λ are respectively

$$m(\Box, x_2)[-\circ \langle x_1 \rangle] \qquad \qquad \Box[-\circ \langle x_1, e, x_2 \rangle] \qquad \qquad m(x_1, \Box)[-\circ \langle x_2 \rangle]$$

This coherent term rewriting system has been considered in various places in literature [8, 16].

We mention here that the notion of Tietze transformation can be defined for 2-TRS in a similar way as for 2-ARS (definition 30):

Definition 71. The *Tietze transformations* are the following possible transformations on a 2-TRS P:

- (T1) given a zig-zag $\pi : t \stackrel{\sim}{\Rightarrow} u$, add a new rewriting rule $\alpha : t \Rightarrow u$ in P₂ together with a new coherence relation $A : \alpha \Rightarrow \pi$ in P₃,
- (T2) given zig-zags $\pi, \rho: t \stackrel{\sim}{\Rightarrow} u$ such that $\pi \stackrel{*}{\Leftrightarrow} \rho$, add a new coherence relation $A: \pi \Rightarrow \rho$ in P_3 .

The *Tietze equivalence* is the smallest equivalence relation on 2-TRS identifying P and Q whenever Q can be obtained from P by a Tietze transformation (T1) or (T2).

Proposition 72. Any two Tietze equivalent 2-TRS present isomorphic groupoids.

5.2. Rewriting properties. Let S be a 2-TRS together with $W \subseteq S_2$. The 2-TRS S induces an 2-ARS in each hom-set: this point of view will allow reusing the work done on 2-ARS on section 3.

Definition 73. Given a 2-TRS S and $n \in \mathbb{N}$, we write S(n, 1) for the 2-ARS whose

- objects are the *n*-ary terms:

$$S(n,1)_0 = S_1^*(n,1)$$

– morphisms are the *n*-ary rewriting steps:

$$\mathsf{S}(n,1)_1 = \mathsf{S}_2^{||}(n,1)$$

 $S(n, 1)_1$ where $S_2^{[]}(n, 1)$ is the set of rewriting steps

$$C[\rho \circ f]: C[t \circ f] \Rightarrow C[u \circ f]$$

with both $C[t \circ f]$ and $C[u \circ f]$ of arity n,

- coherence relations are triples (C, A, f), written $C[A \circ f]$, for some context C, coherence relation $A \in S_3$ and substitution f, of suitable type, of the form

$$C[A \circ f] : C[\pi \circ f] \Longrightarrow C[\pi' \circ f] : C[t \circ f] \Rightarrow C[u \circ f]$$

such that both $C[t \circ f]$ and $C[u \circ f]$ of arity n.

Similarly, a set W induces a set $W(m, 1) \subseteq S(m, 1)_1$, where W(m, 1) is the set of W-rewriting steps, i.e. rewriting steps of the form $C[\alpha \circ f]$ with $\alpha \in W$. We say that a 2-TRS S is W-terminating / locally W-confluent / W-confluent / W-coherent when each S(m, n) is with respect to W(m, n). We say that S is confluent when it is W-confluent for $W = S_2$ (and similarly for other properties). More explicitly,

Definition 74. A *W*-branching (α_1, α_2) is a pair of rewriting steps $\alpha_1 : t \Rightarrow u_1$ and $\alpha_2 : t \Rightarrow u_2$ in $\mathcal{W}^{[]}$ with the same source:

$$u_1 \xleftarrow{\alpha_1} t \xrightarrow{\alpha_2} u_2$$

Such a W-branching is W-confluent when there are cofinal rewriting paths $\pi_1 : u_1 \Rightarrow v$ and $\pi_2 : u_2 \Rightarrow v$ in W^* such that $\overline{\alpha_1 \cdot \pi_1} = \overline{\alpha_2 \cdot \pi_2}$, which is depicted on the left

By extension of proposition 38, we have

Proposition 75. If S is W-terminating and locally W-confluent then it is W-confluent.

In practice, termination can be shown as follows [4, Section 5.2].

Definition 76. A reduction order \geq is a well-founded preorder on terms in S_1^* which is compatible with context extension: given terms $t, u \in S_1^*, t > u$ implies $C[t \circ f] > C[u \circ f]$ for every context C and substitution $f \in S_1^*$ (whose types are such that the expressions make sense).

Proposition 77. A 2-TRS S equipped with a reduction order such that t > u for any rule $\alpha : t \Rightarrow u$ in W is W-terminating.

Proof. For any rewriting step $C[\rho \circ f] : C[t \circ f] \Rightarrow C[u \circ f]$ we have $C[t \circ f] > C[u \circ f]$ and we conclude by well-foundedness.

Moreover, in order to construct a reduction order one can use the following "interpretation method" [4, Section 5.3].

Proposition 78. Let (X, \leq) be a well-founded poset together with an interpretation

$$\llbracket a \rrbracket : X^n \to X$$

of each symbol $a \in S_1$ of arity n as a function which is strictly decreasing in each argument. This induces an interpretation $\llbracket t \rrbracket : X^n \to X$ of every term t of arbitrary arity n defined by induction by

$$\llbracket x_i^n \rrbracket = \pi_i^n \qquad \qquad \llbracket a(t_1, \dots, t_n) \rrbracket = \llbracket a \rrbracket \circ \langle \llbracket t_1 \rrbracket, \dots, \llbracket t_n \rrbracket \rangle$$

where $\pi_i^n : X^n \to X$ is the projection on the *i*-th coordinate. We define an order on functions $f, g : X^n \to X$ by

 $f \succ g$ iff $f(x_1, \dots, x_n) \succ g(x_1, \dots, x_n)$ for every $x_1, \dots, x_n \in X$

and we still write \succeq for the order on terms such that $t \succeq u$ whenever $\llbracket t \rrbracket \succeq \llbracket u \rrbracket$. This order is always a reduction order.

Note that given a reduction order \succeq defined as above, by proposition 77, if we have $t \succ u$ for every rule $\alpha : t \Rightarrow u$ the 2-TRS is W-terminating.

Example 79. Consider the 2-TRS Mon of example 70. We consider the set $X = \mathbb{N} \setminus 0$ and interpret the symbols as

$$\llbracket m(x_1, x_2) \rrbracket = 2x_1 + x_2 \qquad \llbracket e \rrbracket = 1$$

All the rules are decreasing since we have

$$\llbracket m(m(x_1, x_2), x_3) \rrbracket = 4x_1 + 2x_2 + x_3 > 2x_1 + 2x_2 + x_3 = \llbracket m(x_1, m(x_2, x_3)) \rrbracket$$
$$\llbracket m(e, x_1) \rrbracket = 2 + x_1 > x_1 = \llbracket x_1 \rrbracket$$
$$\llbracket m(x_1, e) \rrbracket = 2x_1 + 1 > x_1 = \llbracket x_1 \rrbracket$$

and the rewriting system is terminating.

We now briefly recall the notion of critical pair, see [48] for a more detailed presentation. We say that a branching (α_1, α_2) is smaller than a branching (β_1, β_2) when the second can be obtained from the first by "extending the context", i.e. when there exists a context Cand a morphism f of suitable types such that $\beta_i = C[\alpha_i \circ f]$ for i = 1, 2. In this case, the confluence of the first branching implies the confluence of the second one (see the diagram on the right above). The notion of context can be generalized to define the notion of a binary context C, with two holes, each of which occurs exactly once: we write C[t, u] for the context where the holes have respectively been substituted with terms t and u. A branching is orthogonal when it consists of two rewriting steps at disjoint positions, i.e. when it is of the form

$$C[u_1 \circ f_1, t_2 \circ f_2] \xleftarrow{C[\alpha_1 \circ f_1, t_2 \circ f_2]} C[t_1 \circ f_1, t_2 \circ f_2] \xrightarrow{C[t_1 \circ f_1, \alpha_2 \circ f_2]} C[t_1 \circ f_1, u_2 \circ f_2]$$

for some binary context C, rewriting rules $\alpha_i : t_i \Rightarrow u_i$ in S_2 and morphisms f_i in S_1^* of suitable types. A branching forms a *critical pair* when it is not orthogonal and minimal (wrt the above order). A TRS with a finite number of rewriting rules always have a finite number of critical pairs and those can be computed efficiently [4].

Lemma 80. A 2-TRS S is locally W-confluent when all its critical W-branchings are W-confluent.

Proof. Suppose that all critical W-branchings are confluent. A non-overlapping W-branching is easily shown to be W-confluent. A non-minimal W-branching is greater than a minimal one, which is W-confluent by hypothesis, and is thus itself also W-confluent.

We write $W_3 \subseteq S_3$ for the set of coherence relations $A : \pi \Rightarrow \rho$ such that both π and ρ belong to W^{\sim} . As a useful particular case, we have the following variant of the Squier theorem:

Lemma 81. If 2-TRS S has a coherence relation in W_3 corresponding to a choice of confluence for every critical W-branching then it is locally W-confluent.

Example 82. The 2-TRS Mon of example 70. By definition, every critical pair is confluent and Mon is thus locally confluent. From example 79 and proposition 75, we deduce that it is confluent.

As a direct consequence of proposition 42 and lemma 40, we have

Proposition 83. If S is W-terminating and locally W-confluent then it is W-coherent, and the subgroupoid W of \overline{S} generated by W is thus rigid.

For instance, from examples 79 and 82, we deduce that the 2-TRS Mon is coherent, thus showing the coherence property (C1) for monoidal categories.

Fix a W-convergent 2-TRS S. By proposition 83, \overline{S} is W-coherent, by theorem 55, the quotient functor $\overline{S} \to \overline{S}/W$ is a local equivalence, and by proposition 46, \overline{S}/W is obtained from \overline{P} by restricting to 1-cells in normal form. Moreover, in good situations, we can provide a description of the quotient category \overline{S}/W by applying proposition 48 hom-wise.

6. Coherence for symmetric monoidal categories

In this section, we illustrate the use of the methods developed in the article, by applying them in order to recover the coherence theorems for symmetric monoidal categories [28], which requires quotienting the Lawvere 2-theory of symmetric monoids by a subtheory W, generated by the associator and the unitors. Related results using rewriting in polygraphs where obtained earlier [2, 22, 38]. They however require heavier computations since manipulations of variables (duplication, erasure and commutation) need to be implemented as explicit rules in this context.

6.1. A theory for symmetric monoidal categories. A symmetric monoidal category is a monoidal category equipped with a natural isomorphism $\gamma_{x,y} : x \otimes y \to y \otimes x$, called the symmetry, satisfying the three axioms recalled in section 1.4. A symmetric monoidal category is strict when the structural isomorphisms α , λ and ρ are identities (but we do not require γ to be an identity). We write **SMonCat** (resp. **SMonCat**_{str}) for the category of symmetric monoidal categories (resp. strict ones).

We write SMon for the 2-TRS obtained from Mon, see example 70, by adding a rewriting rule

$$\gamma: m(x_1, x_2) \Rightarrow m(x_2, x_1)$$

corresponding to symmetry, together with a coherence relation

$$F: \gamma(x_1, x_2) \cdot \gamma(x_2, x_1) \Rightarrow \mathrm{id}_{m(x_1, x_2)}$$

which can be pictured as

$$\begin{array}{c} m(x_1, x_2) & \xrightarrow{\gamma} & m(x_2, x_1) \\ \| & \stackrel{F}{\Rightarrow} & \bigvee^{\gamma} \\ m(x_1, x_2) & = & m(x_1, x_2) \end{array}$$

as well as the coherence relations

$$\begin{array}{cccc} m(m(x_1, x_2), x_3) \xrightarrow{\gamma} m(m(x_2, x_1), x_3) \xrightarrow{\alpha} m(x_2, m(x_1, x_3)) & m(e, x_1) \xrightarrow{\gamma} m(x_1, e) \\ & \alpha & & \downarrow \gamma & & \downarrow \gamma \\ m(x_1, m(x_2, x_3)) \xrightarrow{\gamma} m(m(x_2, x_3), x_1) \xrightarrow{\alpha} m(x_2, m(x_3, x_1)) & & x_1 \end{array}$$

It is immediate to see that the algebras of SMon are precisely symmetric monoidal categories:

Proposition 84. The category $Alg(\overline{SMon})$ is isomorphic to the category **SMonCat**.

Since our aim is to study the relationship between symmetric monoidal categories and their strict version, it is natural to consider the set of rewriting rules

$$W = \{\alpha, \lambda, \rho\}$$

i.e. all the rules excepting γ . Namely,

Lemma 85. The category Alg(SMon/W) is isomorphic to the category $SMonCat_{str}$ of strict monoidal categories.

Lemma 86. The 2-TRS SMon is W-coherent.

Proof. Since W consists in α , λ and ρ only, this can be deduced as in the case of monoids: the 2-TRS is W-terminating by example 79 and W-locally confluent by definition (example 70), it is thus W-coherent by proposition 83.

Provided that conjecture 60 holds, we could deduce that any symmetric monoidal category is monoidally equivalent to a strict one. Note that the above reasoning only depends on the convergence of the subsystem induced by W, i.e. on the fact that every diagram made of α , λ and ρ commutes, but it does not require anything on diagrams containing γ 's. In particular, if we removed the compatibility relations G, H, I and J, the strictification theorem would still hold. The resulting notion of strict symmetric monoidal category would however be worrying since, for instance, in absence of I, the 2-cell

$$\gamma_{e,x_1}: m(e,x_1) \Rightarrow m(x_1,e)$$

would induce, in the quotient, a non-trivial automorphism

$$\gamma_{e,x_1}: x_1 \Rightarrow x_1$$

of each variable x_1 . We prove below (theorem 97) a variant of the coherence theorem that is "stronger" in the sense that it requires these axioms to hold and implies that the identity is the only automorphism of x_1 .

6.2. Every affine diagram commutes. We have seen that, for the theory of monoidal categories, "every diagram commutes", in the sense that \overline{Mon} is a 2-rigid (2, 1)-category. For symmetric monoidal categories, we do not expect this to hold since we have two rewriting paths

$$\gamma_{x_1,x_1}: m(x_1,x_1) \Rightarrow m(x_1,x_1) \qquad \text{id}_{m(x_1,x_1)}: m(x_1,x_1) \Rightarrow m(x_1,x_1)$$

which are both from $m(x_1, x_1)$ to itself, and are not equal in general as explained in the introduction. It can however be shown that it holds for the subclass of 2-cells whose source and target are affine terms:

Definition 87. A term t is affine if no variable occurs twice, i.e. $|t|_i \leq 1$ for every index i.

We now explain this, thus recovering a well-known property [47, Theorem 4.1] using rewriting techniques. In order to use those, it will be convenient to work with a variant SMon' of the 2-TRS SMon, obtained by adding a new generator δ as well as coherence relations corresponding to the local confluence diagrams: this variant will allow proving lemma 90 below. A similar completion has been investigated by Lafont [38, Figure 9], although working in a monoidal setting whereas we are working in a cartesian one. We write SMon' for the 2-TRS obtained from SMon by adding a rewriting rule

$$\delta: m(x_1, m(x_2, x_3)) \Rightarrow m(x_2, m(x_1, x_3))$$

removing the coherence relation G and adding coherence relations

$$\begin{split} m(x_1, m(x_2, x_3)) &\stackrel{\delta}{\longrightarrow} m(x_2, m(x_1, x_3)) & m(m(x_1, x_2), x_3) \xrightarrow{\gamma} m(m(x_2, x_1), x_3) \\ \parallel & \stackrel{F'}{\Rightarrow} & \parallel^{\delta} & a \parallel & \stackrel{G'}{\Rightarrow} & \parallel^{\alpha} \\ m(x_1, m(x_2, x_3)) &\longrightarrow m(x_1, m(x_2, x_3)) & m(x_1, m(x_2, x_3)) \xrightarrow{\delta} m(x_2, m(x_1, x_3))) \\ m(m(x_1, x_2), x_3) &\xrightarrow{\gamma} m(x_1, m(x_3, x_2)) \xrightarrow{\delta} m(x_3, m(x_1, x_2)) & m(x_1, e) \xrightarrow{\gamma} m(e, x_1) \\ a \parallel & \stackrel{H}{\Rightarrow} & \parallel \\ m(x_1, m(x_2, x_3)) \xrightarrow{\gamma} m(x_1, m(x_3, x_2)) \xrightarrow{\delta} m(x_3, m(x_1, x_2)) & x_1 \\ m(m(x_1, x_2), m(x_3, x_4)) &\xrightarrow{\delta} m(x_1, m(x_3, m(x_2, x_4))) & \xrightarrow{\delta} m(x_3, m(m(x_1, x_2), x_4)) \\ a \parallel & \stackrel{K}{\Rightarrow} & \parallel^{\alpha} & \parallel^{\alpha} \\ m(x_1, m(x_2, m(x_3, x_4))) \xrightarrow{\delta} m(x_1, m(x_3, m(x_2, x_4))) \xrightarrow{\delta} m(x_3, m(x_1, m(x_2, x_4))) \\ m(x_1, m(x_2, m(x_3, x_4))) \xrightarrow{\delta} m(x_2, m(x_1, m(x_3, x_4))) \xrightarrow{\delta} m(x_2, m(x_3, m(x_1, x_4))) \\ m(x_1, m(x_2, m(x_3, x_4))) \xrightarrow{\delta} m(x_2, m(x_1, x_3, x_4))) \xrightarrow{\delta} m(x_2, m(x_3, m(x_1, x_4))) \\ m(x_1, m(x_2, x_3), x_4)) \xrightarrow{\delta} m(x_2, m(x_1, x_3)) \xrightarrow{\delta} m(x_2, m(x_3, x_1))) \\ m(x_1, m(x_2, x_3), x_4)) \xrightarrow{\delta} m(x_2, m(x_1, x_3)) \xrightarrow{\delta} m(x_2, m(x_3, x_4))) \\ m(x_1, m(x_2, m(x_3, x_4))) \xrightarrow{\delta} m(x_2, m(x_1, x_3)) \xrightarrow{\delta} m(x_2, m(x_3, x_4))) \\ m(x_1, m(x_2, m(x_3, x_4))) \xrightarrow{\delta} m(x_2, m(x_1, x_3)) \xrightarrow{\delta} m(x_2, m(x_3, m(x_1, x_4)))) \\ m(x_1, m(x_3, m(x_2, x_4))) \xrightarrow{\delta} m(x_3, m(x_1, x_2)) \xrightarrow{\gamma} m(x_3, m(x_2, m(x_1, x_4))) \\ m(x_1, m(x_3, m(x_2, x_4))) \xrightarrow{\delta} m(x_3, m(x_1, m(x_3, x_4))) \xrightarrow{\delta} m(x_3, m(x_2, m(x_3, m(x_1, x_4)))) \\ m(x_1, m(x_3, m(x_2, x_4))) \xrightarrow{\delta} m(x_3, m(x_1, m(x_2, x_4))) \xrightarrow{\delta} m(x_3, m(x_2, m(x_1, x_4))) \\ m(x_1, m(x_3, m(x_2, x_4))) \xrightarrow{\delta} m(x_3, m(x_1, m(x_2, x_4))) \xrightarrow{\delta} m(x_1, m(x_2, x_2)) \xrightarrow{\delta} m(x_1, m(x_2, x_2)) \xrightarrow{\delta} m(x_1, m(x_2, m(x_1, x_2)) \xrightarrow{\delta} m(x_1, m(x_2, x_2)) \xrightarrow{\delta} m(x_1, m(x_2, x_2)) \xrightarrow{\delta} m(x_1, x_2) \xrightarrow{K} m(x_1, x_2) \xrightarrow{K}$$

The following adapts [22, Proposition 3.3.5] from the monoidal setting to our cartesian setting, see also [2]:

Proposition 88. The 2-TRS SMon and SMon' present isomorphic categories.

Proof. By proposition 72, it is enough to show that both 2-TRS are Tietze equivalent. The commutation of H and I is immediate in presence of the other axioms, we can thus add them using Tietze transformations (T2). Namely, we can show the commutation of H by using F and G twice:

and the commutation of J can be obtained from F and I:



Next, by a Tietze transformation (T1), we can add the rule δ together with its definition

$$\delta(x_1, x_2, x_3) = \alpha(x_1, x_2, x_3) \circ m(\gamma(x_1, x_2), x_3) \circ \alpha(x_1, x_2, x_3)^{-1}$$

which is formally given by the relation G'. From this definition, one easily shows that the coherence relations K to R are derivable and can thus be added by Tietze transformations (T2). Finally, the coherence relation G is then superfluous, since it can be derived as

and can thus be removed by a Tietze transformation (T2).

Lemma 89. The 2-TRS SMon' is locally confluent.

Proof. By lemma 80, it is enough to show that all the critical pairs are confluent, which holds by definition of SMon': the critical pairs involving α , λ and ρ are handled in example 70, and those involving γ or δ and another rewriting rule in the above definition of SMon'.

Again, analogous results in a monoidal (as opposed to cartesian) setting predate this work, see [38, Figure 51] and [21, Section 5.3.3].

The 2-TRS SMon' is not terminating (even when restricted to affine terms) because of the rules γ and δ which witnesses for the commutativity of the operation m: for instance,

we have the loop

$$m(x_1, x_2) \xrightarrow{\gamma(x_1, x_2)} m(x_2, x_1) \xrightarrow{\gamma(x_2, x_1)} m(x_1, x_2)$$
(6.1)

In order to circumvent this problem, we are going to formally "remove" the second morphism above and only keep instances of γ (resp. δ) which tend to make variables in decreasing order. Namely, by the coherence relation F, i.e.



we have $\gamma(x_2, x_1) = \gamma(x_1, x_2)^-$ so that $\gamma(x_2, x_1)$ is superfluous and we can remove it, by using Tietze transformations, without changing the presented (2, 1)-category. Note that this operation is clearly not stable under substitution (for instance consider the substitution $\langle x_2, x_1 \rangle$ which exchanges the two variable names), so that this cannot actually be performed at the level of (2, 1)-categories, but it can if we work within the hom-groupoids, which will be enough for our purposes. If we remove all the rewriting steps involving γ which tend to put variables in increasing order as explained above, we still have some loops such as

$$m(e, x_1) \xrightarrow{\gamma(e, x_1)} m(x_1, e) \xrightarrow{\gamma(x_1, e)} m(e, x_1)$$

Intuitively, this is because the above rewriting path involves terms containing a unit e, whereas our previous criterion relies on the order of variables. Fortunately, we can first remove all units by applying the rules λ and ρ , and then apply the above argument.

Fix an arity $n \in \mathbb{N}$, and consider the 2-ARS $\mathsf{P} = \mathsf{SMon}'(n, 1)$ as defined in definition 73. We write P' for the 2-ARS obtained from P by

- removing from P_1 the terms where the unit *e* occurs, excepting *e* itself,
- removing from P_2 the rewriting steps whose source or target terms contain e (in particular, we remove all rewriting steps involving λ or ρ),
- removing from P_3 the coherence relations where a removed step occurs in the source or the target.

Lemma 90. The groupoid presented by P' is equivalent to the one presented by P.

Proof. We write $W \subseteq \mathsf{P}_1$ for the set of rewriting steps involving λ or ρ . By lemma 89, the 2-ARS P' is locally *W*-confluent, and thus, by lemma 45, $\overline{\mathsf{P}}$ is equivalent to $N(\overline{\mathsf{P}})$, the full subcategory on *W*-normal forms. In turn, by proposition 48 (see below for details), the category $N(\overline{\mathsf{P}})$ is isomorphic to the groupoid presented by $\mathsf{P} \setminus W$ and we conclude by observing that the 2-ARS P' is precisely $\mathsf{P} \setminus W$ (terms in normal form are precisely those where *e* does not occur, with the exception of *e* itself).

Let us explain why the conditions of proposition 48 are satisfied.

- (1) We have seen above that P is W-convergent.
- (2) It is immediate to check that no rewriting rule can produce a term containing e from a term which does not have this property.

(3) From proposition 49, in order to show this condition, we have to check that every diagram of the form



can be closed as in (3.3) for arbitrary rewriting steps $\alpha \in \mathsf{P}_1$ and $\omega \in W$. It is enough to show this when they form a critical pair. There are five of them, which correspond to the five coherence relations B, C, D, I and J, from which we conclude.

(4) From proposition 49, in order to show this condition, we have to check that every diagram of the form



can be closed as in (3.4) for ω in W. Again, it is enough to show this in situations which are not orthogonal and minimal, in a similar sense as for critical pairs, which we call a "critical pair between a rewriting rule and a coherence relation". For instance, one such critical pair between λ and A can be closed as follows:

where the vertical dotted arrows are the obvious rewriting steps involving λ (this is the only critical pair between λ and A and there are three critical pairs between ρ and A).

Lemma 91. The 2-ARS P' is locally confluent.

Proof. We can deduce local confluence of P' from the one of P: given a local pair with t as source, it is confluent in P by lemma 89 and thus in P'. Namely, since t lies in P', the whole diagram does by property (2) shown in the proof of lemma 90 above.

Given a term t, we write ||t|| the list of variables occurring in it, from left to right, e.g. $||m(m(x_2, e), x_1)|| = x_2x_1$. We order variables by $x_i \succeq x_j$ whenever $i \le j$ and extend it to lists of variables by lexicographic ordering. Given terms t and u, we write $t \succeq u$ when ||t|| is greater than u according to the preceding order.

Lemma 92. The preorder \succeq is well-founded on affine terms with fixed arity.

Proof. Any infinite decreasing sequence $t_1 \succ t_2 \succ \ldots$ of terms, would induce an infinite decreasing sequence $||t||_1 \succ ||t||_2 \succ \ldots$ of lists of variables, but there is only a finite number

of those since we consider affine terms (so that there are no repetitions of variables) of fixed arity (so that there is a finite number of variables). \Box

A rewriting step $\rho : t \Rightarrow u$ in P'_1 is *decreasing* when $t \succ u$. We write P'' for the 2-ARS obtained from P' by

- removing from P'_1 all the rewriting steps of the form

$$C[\gamma(t_1, t_2)] : C[m(t_1, t_2)] \Rightarrow C[m(t_2, t_1)]$$

which are not decreasing,

- replacing in the source or target of a relation in P'_2 all the non-decreasing steps $C[\gamma(t_1, t_2)]$ by $C[\gamma(t_2, t_1)^-]$.

Lemma 93. The 2-ARS P' and P" present isomorphic groupoids.

Proof. This is a direct application of lemma 32, because P'_3 contains the coherence rules

which allow us to conclude.

In the following, for a 2-ARS whose objects are terms such as P'' , we say that it is "terminating on affine terms" when there is no infinite sequence of rewriting steps $t_0 \to t_1 \to \ldots$ where the term t_0 is affine. Note that all the rules of P'' rewrite affine terms into affine terms, so that all the t_i are also necessarily affine in such a sequence of rewriting steps.

Lemma 94. The 2-ARS P'' is terminating on affine terms and locally confluent.

Proof. In order to show termination, we can take the lexicographic product of the orders \succeq of lemma 92 and the one of example 79. This order is well-founded as a lexicographic product of well-founded orders. The rewriting steps involving γ are strictly decreasing wrt \succeq , by definition of P'' . The rewriting steps involving α are left invariant by \succeq but are strictly decreasing wrt the second order. We deduce that P'' is terminating.

We have seen in lemma 91 that P' is locally confluent. We thus have that P'' is also locally confluent because the confluence diagrams involving γ (namely G, H, I and J) only require decreasing instances of rewriting rules involving γ .

As a direct consequence, we have:

Lemma 95. Given two rewriting zig-zags $p, q : t \xrightarrow{\sim} u$ in P'' such that t is affine, we have that u is also affine and $p \Leftrightarrow^* q$.

Proof. By lemma 94, the restriction of P'' to affine terms is terminating and locally confluent, thus confluent by proposition 38 and thus coherent by proposition 42.

From the properties shown in section 3.2, we deduce that P'' is coherent, which implies the following:

Lemma 96. Given two terms $t, u : n \to 1$, with t affine, there is at most one 2-cell $t \Rightarrow u$ in SMon.

Proof. We write $\mathsf{P} = \mathsf{SMon}'(n, 1)$. Given two rewriting zig-zags $p, q: t \stackrel{\sim}{\Rightarrow} u$, we have

$\overline{SMon}(p,q)\cong\overline{P}(p,q)$	by proposition 88
$\cong \overline{P}'(p,q)$	by lemma 90,
$\cong \overline{P}''(p,q)$	by lemma 93,
$\cong 1$	by lemma 95,

from which we conclude.

Finally, we thus conclude to the following coherence theorem (S1):

Theorem 97. In a symmetric monoidal category, every diagram whose 0-source is a tensor product of distinct objects, and whose morphisms are composites and tensor products of structural morphisms, commutes.

Proof. Fix a symmetric monoidal category C. By proposition 84, C can be seen as a product preserving 2-functor $\overline{\mathsf{SMon}} \to \mathbf{Cat}$. A coherence diagram in C thus corresponds to a pair of rewriting paths $p, q: t \stackrel{*}{\Rightarrow} u$ in SMon_2^* for some terms t and u. When the 0-source of the coherence diagram is a tensor product of distinct objects, t affine, and we have p = q in $\overline{\mathsf{SMon}}$ by lemma 96.

Previous theorem essentially follows from lemma 95, which identifies a class of 1-cells of \overline{SMon} , the affine ones, between which there is at most one 2-cell. The following proposition characterizes, among them, those between which there actually exists a 2-cell.

Proposition 98. Given two affine terms t and u, there exists a rewriting zig-zag $p: t \stackrel{\sim}{\Rightarrow} u$ if and only if t and u have the same variables.

Proof. The left-to-right implication can be proved by checking that all the rewriting rules of SMon preserve the variables of rewritten terms. Now, consider the right-to-left implication and suppose that t and u are two terms with the same variables. By example 79 and lemma 45, we can suppose that t and u are in normal form with respect to the rules $W = \{\alpha, \lambda, \rho\}$, and thus respectively of the form

$$t = m(x_0, m(x_1, m(\dots, m(x_{n-2}, x_{n-1}))))$$

$$u = m(x_{f(0)}, m(x_{f(1)}, m(\dots, m(x_{f(n-2)}, x_{f(n-1)}))))$$

(up to renaming the variables of t) for some bijection $f: [n] \to [n]$ with $[n] = \{0, \ldots, n-1\}$. From the well-known fact that any such bijection can be obtained as a composite of adjacent transpositions, it is easy to construct a 2-cell $t \stackrel{*}{\Rightarrow} u$ as a composite of rewriting steps γ or δ corresponding to those transpositions. For instance, with $t = m(x_0, m(x_1, x_2))$ and $u = m(x_2, m(x_1, x_0))$, the bijection $f: [3] \to [3]$ is defined by f(0) = 2, f(1) = 1 and f(0) = 0, which can be obtained as the composite of adjacent transpositions (01) and (12), and the corresponding 2-cell is

$$m(x_0, m(x_1, x_2)) \xrightarrow{\delta(x_0, x_1, x_2)} m(x_1, m(x_0, x_2)) \xrightarrow{m(x_1, \gamma(x_0, x_2))} m(x_2, m(x_1, x_0))$$

6.3. General coherence. Finally, we explain how to recover the more general coherence theorem for symmetric monoidal categories. Given a natural number n, we write $[n] = \{0, \ldots, n-1\}$ for the cardinal with n elements.

We define a Lawvere 2-theory S as follows. Its 0-cells are natural numbers, as for any Lawvere theory. Given a 0-cell n, a 1-cell $f: n \to 1$ is a list $[f^1, f^2, \ldots, f^{|l|}]$ of elements of [n], sometimes being referred to as *colors*, the natural number |f| being the *length* of the list. Since a Lawvere 2-theory is cartesian, we more generally have that a morphism $n \to m$ is a tuple $\langle f_0, \ldots, f_{m-1} \rangle$ of m 1-cells $f_i: n \to 1$. Given 1-cells $f: m \to n$ and $g: n \to 1$, their composition is the list computed by taking the list g, replacing each color g^i by f_i so that we obtain a list of lists, and then flattening the resulting list, i.e. taking the concatenation of all the lists. For instance, we have the composite

$$4 \xrightarrow{\langle [1,1], [3,3,2], [2,0,3] \rangle} 3 \xrightarrow{\langle [2,0,2] \rangle} 1 = 4 \xrightarrow{\langle 2,0,3,1,1,2,0,3 \rangle} 1$$

The 2-cells $\alpha : f \Rightarrow g : n \to 1$ are color-preserving bijections on n elements, i.e. bijections $\alpha : [n] \to [n]$ such that $f^{\alpha(i)} = g^i$ for every $i \in [n]$. More generally, 2-cells $\alpha : f \Rightarrow g : m \to 1$ are tuples $\langle \alpha^0, \ldots, \alpha^{m-1} \rangle$ of 2-cells $\alpha^i : n \to 1$ for $i \in [m]$. The 1-composition of 2-cells is the usual composition of functions, and the 0-composition is left to the reader.

We claim that this 2-theory provides an explicit description of the quotient category for the theory of symmetric monoids, thus establishing a coherence theorem (S2) or (C2). In order to prove this, we will need to relate affine terms to non-affine ones. Informally, every non-affine term can be described as an affine term in which some variables have been identified, and moreover rewriting between non-affine terms can be uniquely lifted to the affine term they originate from. We say that a 1-cell $f: m \to n$ of SMon^{*}₁ is a *renaming* when it is a tuple of variables, i.e. of the form $\langle x_{i_0}, \ldots, x_{i_{k-1}} \rangle$.

Lemma 99. Consider the 2-TRS SMon. For any term $t : n \to 1$, there is a natural number \tilde{n} and an affine term $\tilde{t} : \tilde{n} \to 1$ together with a renaming $f : n \to \tilde{n}$ such that $t = \tilde{t} \circ f$:

$$\begin{array}{c} \tilde{n} & \overset{t}{\longrightarrow} & 1 \\ f \uparrow & & \parallel \\ n & \overset{t}{\longrightarrow} & 1 \end{array}$$

Moreover, for any rewriting zig-zag $p: t \stackrel{\sim}{\Rightarrow} u$, there is a unique term \tilde{u} and rewriting path $p: \tilde{t} \stackrel{\sim}{\Rightarrow} \tilde{u}$ such that $\tilde{p} \circ f = p$:

$$\begin{array}{c}
\tilde{t} \\
\tilde{p} \\
\tilde{p} \\
\tilde{v} \\
\tilde{t} \\
n \\
u
\end{array}$$

$$\begin{array}{c}
\tilde{t} \\
\tilde{u} \\
\tilde{u} \\
1
\end{array}$$

and \tilde{u} is affine. Moreover, any equivalence $p \Leftrightarrow^* q$ between $p, q : t \xrightarrow{\sim} u$ lifts as an equivalence $\tilde{p} \Leftrightarrow^* \tilde{q}$.

Proof. Given a term t, we can construct \tilde{t} by renaming distinctively every occurrence of each variable. For instance, the term $m(x_1, m(x_0, x_1)) : 2 \to 1$ can be obtained form the affine term $m(x_0, m(x_1, x_2)) : 3 \to 1$ with the renaming $\langle x_1, x_0, x_1 \rangle : 2 \to 3$. The fact that we can

lift rewriting steps follows from the fact that all the rewriting rules in SMon_1 have affine source and targets. And similarly for coherence relations.

The same structure can be identified in S. We say that a 1-cell $f : n \to 1$ is affine, when it consists in a list which does not contain the same element twice. We say that a 1-cell $f : m \to n$ is a *renaming* when it is a tuple of lists of length 1, i.e. of the form $\langle [i_0], \ldots, [i_{k-1}] \rangle$.

Lemma 100. Consider the Lawvere 2-theory S. For every 1-cell $t : n \to 1$ there is an affine 1-cell $\tilde{t} : n \to 1$ together with a renaming $f : n \to \tilde{n}$ such that $t = \tilde{t} \circ f$, moreover any 2-cell $p : t \Rightarrow u$ lifts uniquely as a 2-cell $\tilde{p} : \tilde{t} \Rightarrow \tilde{u}$ such that $\tilde{p} \circ f = f$.

Proof. Given a 1-cell $t: n \to 1$, i.e. a list $[i_0, i_1, \ldots, i_{k-1}]$ with values in [n], we can take $\tilde{n} = k$ and $\tilde{t} = [0, 1, \ldots, k-1]: k \to 1$ together with the renaming $f = \langle [i_0], [i_1], \ldots, [i_{k-1}] \rangle : n \to k$. Given a morphism $p: t \Rightarrow u$, i.e. a color-preserving bijection $p: [k] \to [k]$, it is easily shown that the same bijection provides a suitable 2-cell $t \Rightarrow u$, with $u = [p(0), p(1), \ldots, p(k-1)]$, and that this is the only possible one.

We can finally show the announced coherence theorem (C2) for symmetric monoidal categories:

Theorem 101. We have an isomorphism of Lawvere 2-theories $\overline{\mathsf{SMon}}/W \cong S$.

Proof. The two 2-theories have the same 0-cells: the natural numbers, as any Lawvere theory. Since W forms a rewriting system, the 1-cells of $\overline{\mathsf{SMon}}/W$ can be identified to the 1-cells $n \to 1$ of $\overline{\mathsf{SMon}}$ which are in normal form with respect to the rules of W, i.e. terms of the form

$$m(x_{i_0}, m(x_{i_1}, m(\dots, m(x_{i_{k-2}}, x_{i_{k-1}})))))$$

for some $k \in \mathbb{N}$ with $i_j \in [n]$ for $j \in [k]$, and those are clearly in bijection with lists

$$[i_0, i_2, \ldots, i_{n-2}, i_{k-1}]$$

of elements of [n], and the compositions coincide in both categories. Given two terms $t, u : n \to 1$, the category $\overline{\mathsf{SMon}}/W(t, u)$ can be identified with $\overline{\mathsf{SMon}}(\hat{t}, \hat{u})$ by proposition 46, where \hat{t} is the normal form of t with respect to the rules of W and similarly for \hat{u} . In the following, we implicitly assume that the 1-cells we consider are in normal form and make this identification. Given two terms $t, u : n \to 1$, with t affine, we know by proposition 98 that there is at most one 2-cell $t \Rightarrow u$ in $\overline{\mathsf{SMon}}/W$, and this is the case precisely when t and u have the same variables. If we consider the corresponding lists, still written t and u, in \mathcal{S} , we can observe that there is most one color-preserving bijection between the two lists (because they do not contain the same color twice) and this is the case precisely when they have the same colors. We thus have a bijection between the sets of 2-cells $\overline{\mathsf{SMon}}/W(t, u) \cong \mathcal{S}(t, u)$ when t and u are affine. Finally, given two arbitrary 1-cells $t, u : n \to 1$, by lemma 99 the set of 2-cells $\overline{\mathsf{SMon}}/W(t, u)$ can be described as

$$\overline{\mathsf{SMon}}/W(t,u) = \bigsqcup_{\substack{u':n \to 1 \\ u' \text{ affine}}} \left\{ p: t \Rightarrow u' \mid u' \circ f = u \right\}$$

Similarly, by lemma 100 the set of 2-cells $\mathcal{S}(t, u)$ can be described as

$$\mathcal{S}(t,u) = \bigsqcup_{\substack{u': n \to 1 \\ u' \text{ affine}}} \left\{ p: t \Rightarrow u' \mid u' \circ f = u \right\}$$

45

Those sets are isomorphic as disjoint unions of 2-cells whose source is affine, and the bijection can be checked to be compatible with composition. \Box

Alternatively, the proof could be done by constructing explicitly a presentation of the Lawvere 2-theory S, following previous work in the setting of monoidal categories [22, 38]. The proof scheme used above however has the advantage of being simpler to check, and we believe that it is quite general; in particular, extensions to the study of coherence of fundamental structures such as cartesian closed categories should be developed in subsequent works.

7. FUTURE WORKS

We claim that the framework developed here applies to a wide variety of algebraic structures, which will be explored in subsequent work. In fact, the full generality of the framework was not needed for (symmetric) monoidal categories, since the rules of the corresponding theory never need to duplicate or erase variables (and, in fact, those can be handled by traditional polygraphs [22, 38]). This is however, needed for the case of rig categories [41], which feature two monoidal structures \oplus and \otimes , and natural isomorphisms such as

$$\delta_{x,y,z}: x \otimes (y \oplus z) \to (x \otimes y) \oplus (x \otimes z)$$

where x occurs twice in the target, generalizing the laws for rings. Those were a motivating example for this work, and we will develop elsewhere a proof of coherence of those structures based on our rewriting framework, as well as related approaches on the subject [10, Appendix G].

The importance of the notion of polygraph can be explained by the fact that they are the cofibrant objects in a model structure on ω -categories [3, 39]. It would be interesting to develop a similar point of view for higher-dimensional term rewriting systems: a first step in this direction is the model structure developed in [67].

References

- [1] Coherence theorem for monoidal categories. URL: https://ncatlab.org/nlab/show/coherence+theorem+for+monoidal+categories.
- Matteo Acclavio. A constructive proof of coherence for symmetric monoidal categories using rewriting. Preprint, 2016. arXiv:1606.01722.
- [3] Dimitri Ara, Albert Burroni, Yves Guiraud, Philippe Malbos, François Métayer, and Samuel Mimram. Polygraphs: From rewriting to higher categories. To appear, 2023. arXiv:2312.00429.
- [4] Franz Baader and Tobias Nipkow. Term rewriting and all that. Cambridge university press, 1999.
- [5] A. A. Babaev and S. V. Solov'ev. A coherence theorem for canonical morphisms in cartesian closed categories. *Journal of Soviet Mathematics*, 20:2263–2279, 1982.
- [6] Michael Barr and Charles Wells. Toposes, triples, and theories. Springer-Verlag, 2000.
- Marek A. Bednarczyk, Andrzej M. Borzyszkowski, and Wieslaw Pawlowski. Generalized congruences Epimorphisms in Cat. Theory and Applications of Categories, 5(11):266–280, 1999.
- [8] Tibor Beke. Categorification, term rewriting and the Knuth-Bendix procedure. Journal of Pure and Applied Algebra, 215(5):728–740, 2011.
- [9] Richard F. Blute, J. Robin B. Cockett, Robert A. G. Seely, and Todd H. Trimble. Natural deduction and coherence for weakly distributive categories. *Journal of Pure and Applied Algebra*, 113(3):229–296, 1996.
- [10] Filippo Bonchi, Alessandro Di Giorgio, and Alessio Santamaria. Tape diagrams for rig categories with finite biproducts. Technical report, Università di Pisa, 2022.
- [11] Filippo Bonchi, Fabio Gadducci, Aleks Kissinger, Pawel Sobocinski, and Fabio Zanasi. String diagram rewrite theory I: rewriting with Frobenius structure. *Journale of the ACM*, 69(2):14:1–14:58, 2022.

- [12] Filippo Bonchi, Fabio Gadducci, Aleks Kissinger, Pawel Sobocinski, and Fabio Zanasi. String diagram rewrite theory II: rewriting with symmetric monoidal structure. *Mathematical Structures in Computer Science*, 32(4):511–541, 2022.
- [13] Filippo Bonchi, Fabio Gadducci, Aleks Kissinger, Pawel Sobocinski, and Fabio Zanasi. String diagram rewrite theory III: confluence with and without Frobenius. *Mathematical Structures in Computer Science*, 32(7):829–869, 2022.
- [14] Albert Burroni. Higher-dimensional word problems with applications to equational logic. Theoretical computer science, 115(1):43–62, 1993.
- [15] Florence Clerc and Samuel Mimram. Presenting a category modulo a rewriting system. In 26th International Conference on Rewriting Techniques and Applications (RTA 2015). Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2015.
- [16] Jonathan Asher Cohen. *Coherence for rewriting 2-theories*. PhD thesis, Australian National University, 2008.
- [17] Marcelo Fiore and Philip Saville. Coherence for bicategorical cartesian closed structure. Mathematical Structures in Computer Science, 31(7):822–849, 2021.
- [18] Stéphane Gaussent, Yves Guiraud, and Philippe Malbos. Coherent presentations of Artin monoids. Compositio Mathematica, 151(5):957–998, 2015.
- [19] Robert Gordon, Anthony John Power, and Ross Street. Coherence for tricategories, volume 558. American Mathematical Society, 1995.
- [20] John W. Gray. 2-algebraic theories and triples. Cahiers de Topologie et Géométrie Différentielle, 14:178– 180, 1973.
- [21] Yves Guiraud. Présentations d'opérades et systèmes de réécriture. PhD thesis, Montpellier 2 University, France, 2004.
- [22] Yves Guiraud and Philippe Malbos. Coherence in monoidal track categories. Mathematical Structures in Computer Science, 22(6):931–969, 2012.
- [23] Yves Guiraud and Philippe Malbos. Polygraphs of finite derivation type. Mathematical Structures in Computer Science, 28(2):155–201, 2018.
- [24] Nick Gurski. Coherence in three-dimensional category theory, volume 201. Cambridge University Press, 2013.
- [25] Gérard Huet. Confluent reductions: Abstract properties and applications to term rewriting systems. Journal of the ACM (JACM), 27(4):797–821, 1980.
- [26] C. Barry Jay. The structure of free closed categories. Journal of Pure and Applied Algebra, 66(3):271–285, 1990.
- [27] Niles Johnson and Donald Yau. 2-dimensional categories. Oxford University Press, USA, 2021.
- [28] André Joyal and Ross Street. Braided tensor categories. Advances in Math, 102:20–78, 1993.
- [29] G. Maxwell Kelly. On MacLane's conditions for coherence of natural associativities, commutativities, etc. Journal of Algebra, 1(4):397–402, 1964.
- [30] G. Maxwell Kelly. An abstract approach to coherence. In *Coherence in categories*, pages 106–147. Springer, 1972.
- [31] G. Maxwell Kelly and Miguel L. Laplaza. Coherence for compact closed categories. Journal of pure and applied algebra, 19:193–213, 1980.
- [32] Gregory Maxwell Kelly and Saunders MacLane. Coherence in closed categories. Journal of Pure and Applied Algebra, 1(1):97–140, 1971.
- [33] Nicolai Kraus and Jakob von Raumer. A rewriting coherence theorem with applications in homotopy type theory. *Mathematical Structures in Computer Science*, 32(7):982–1014, 2022.
- [34] Stephen Lack. A Quillen model structure for 2-categories. K-theory, 26(2):171–205, 2002.
- [35] Stephen Lack. Codescent objects and coherence. Journal of Pure and Applied Algebra, 175(1-3):223–241, 2002.
- [36] Stephen Lack and Giacomo Tendas. Flat vs. filtered colimits in the enriched context. Advances in Mathematics, 404:108381, 2022.
- [37] Yves Lafont. A new finiteness condition for monoids presented by complete rewriting systems (after Craig C. Squier). Journal of Pure and Applied Algebra, 98(3):229–244, 1995.
- [38] Yves Lafont. Towards an algebraic theory of boolean circuits. Journal of Pure and Applied Algebra, 184(2-3):257–310, 2003.

- [39] Yves Lafont, François Métayer, and Krzysztof Worytkiewicz. A folk model structure on omega-cat. Advances in Mathematics, 224(3):1183–1231, 2010.
- [40] Jim Lambek. Deductive systems and categories II: Standard constructions and closed categories. Lecture Notes in Mathematics, 86:76–122, 1969.
- [41] Miguel L. Laplaza. Coherence for distributivity. In Coherence in categories, pages 29–65. Springer, 1972.
- [42] F. William Lawvere. Functorial semantics of algebraic theories. Proceedings of the National Academy of Sciences, 50(5):869–872, 1963.
- [43] Tom Leinster. Higher operads, higher categories. Number 298. Cambridge University Press, 2004.
- [44] Maxime Lucas. A coherence theorem for pseudonatural transformations. Journal of Pure and Applied Algebra, 221(5):1146–1217, 2017.
- [45] Roger C. Lyndon and Paul E. Schupp. Combinatorial group theory, volume 188. Springer, 1977.
- [46] Saunders Mac Lane. Categories for the Working Mathematician, volume 5. Springer Science & Business Media, 1998.
- [47] Saunders MacLane. Natural associativity and commutativity. Rice Institute Pamphlet-Rice University Studies, 49(4), 1963.
- [48] Philippe Malbos and Samuel Mimram. Homological computations for term rewriting systems. In 1st International Conference on Formal Structures for Computation and Deduction (FSCD 2016), volume 52, pages 27–1, 2016.
- [49] J. Peter May. E_∞-spaces, group completions and permutative categories. New Developments in Topology, London Mathematical Society Lecture Note Series, 11, 1974.
- [50] Samuel Mimram. Categorical coherence from term rewriting systems. In 8th International Conference on Formal Structures for Computation and Deduction (FSCD 2023), volume 260 of LIPIcs, page 16:1–16:17. Schloss Dagstuhl, 2023.
- [51] Samuel Mimram and Pierre-Louis Curien. Coherent presentations of monoidal categories. Logical Methods in Computer Science, 13, 2017.
- [52] Gregorii E. Mints. A simple proof of the coherence theorem for cartesian closed categories. 1982.
- [53] Maxwell H. A. Newman. On theories with a combinatorial definition of "equivalence". Annals of Mathematics, 43(2):223–243, 1942.
- [54] John Power. An abstract formulation for rewrite systems. In Category Theory and Computer Science, pages 300–312, Berlin, Heidelberg, 1989. Springer Berlin Heidelberg.
- [55] John Power. A general coherence result. Journal of Pure and Applied Algebra, 57(2):165–173, 1989.
- [56] John Power. Enriched Lawvere theories. Theory and Applications of Categories, 6(7):83–93, 1999.
- [57] Jiří Rosickỳ et al. Locally presentable and accessible categories, volume 189. Cambridge University Press, 1994.
- [58] Michael A. Shulman. Not every pseudoalgebra is equivalent to a strict one. Advances in Mathematics, 229(3):2024–2041, 2012.
- [59] Mei Chee Shum. Tortile tensor categories. Journal of Pure and Applied Algebra, 93(1):57–110, 1994.
- [60] Craig C. Squier, Friedrich Otto, and Yuji Kobayashi. A finiteness condition for rewriting systems. *Theoretical Computer Science*, 131(2):271–294, 1994.
- [61] Ross Street. Limits indexed by category-valued 2-functors. Journal of Pure and Applied Algebra, 8(2):149– 181, 1976.
- [62] Ross Street. Categorical structures. Handbook of algebra, 1:529–577, 1996.
- [63] Terese. Term rewriting systems. Cambridge University Press, 2003.
- [64] Heinrich Tietze. Über die topologischen Invarianten mehrdimensionaler Mannigfaltigkeiten. Monatshefte für Mathematik und Physik, 19:1–118, 1908.
- [65] Rodiani Voreadou. Coherence and non-commutative diagrams in closed categories, volume 182. American Mathematical Society, 1977.
- [66] Noson S. Yanofsky. The syntax of coherence. Cahiers de topologie et géométrie différentielle catégoriques, 41(4):255–304, 2000.
- [67] Noson S. Yanofsky. Coherence, Homotopy and 2-Theories. K-Theory, 23(3):203-235, 2001.