

OBSTRUCTIONS TO WEAK DECOMPOSABILITY FOR SIMPLICIAL POLYTOPES

NICOLAI HÄHNLE, STEVEN KLEE, AND VINCENT PILAUD

Abstract. Provan and Billera introduced notions of (weak) decomposability of simplicial complexes as a means of attempting to prove polynomial upper bounds on the diameter of the facet-ridge graph of a simplicial polytope. Recently, De Loera and Klee provided the first examples of simplicial polytopes that are not weakly vertex-decomposable. These polytopes are polar to certain simple transportation polytopes. In this paper, we refine their analysis to prove that these d -dimensional polytopes are not even weakly $O(\sqrt{d})$ -decomposable. As a consequence, (weak) decomposability cannot be used to prove a polynomial version of the Hirsch Conjecture.

1. Introduction

The Hirsch Conjecture, which was originally proposed by Hirsch in a letter to Dantzig in 1957, stated that the graph of a simple d -dimensional polytope with n facets has diameter at most $n-d$. Over the past 55 years, the Hirsch Conjecture has motivated a tremendous amount of research on diameters of polytopes. We do not attempt to recount this work here, but instead we point out excellent surveys of De Loera [DL11], Kim and Santos [KS10], and Todd [Tod02]. The most remarkable result in this realm is Santos's counterexample to the Hirsch Conjecture [San12] and its recent improvement by Matschke, Santos, and Weibel [MSW12].

These counterexamples all have diameter $(1 + \epsilon)(n - d)$. It leaves open the Polynomial Hirsch Conjecture, which states that the diameter of the graph of a simple d -dimensional polytope with n facets is bounded by a polynomial function of n and d . This conjecture was the subject of a recent Polymath project hosted by Kalai [Kal10]. The current best known upper bounds on diameter are quasi-exponential in n and d [KK92] or linear in fixed dimension but exponential in

of a (weakly) k -decomposable simplicial complex is bounded by a polynomial of degree $k + 1$ in its number n of vertices and its dimension d .

One approach to proving the Polynomial Hirsch Conjecture would be to show that there is a fixed integer k such that all simplicial polytopes are (weakly) k -decomposable. In this paper, we show that this approach will not work, according to the following counterexample.

Theorem. *Let a, b be positive integers. There is a simple $2 \times (a + b + 1)$ transportation polytope whose simplicial polar is not weakly k -decomposable for any k satisfying $\frac{k+3}{2} \leq \min\{a, b\}$.*

The remainder of the paper is structured as follows. In Section 2, we give all relevant definitions related to simplicial complexes, their diameters, and (weak) decomposability. Further, we discuss the family of (simple) transportation polytopes and their combinatorial properties. In Section 3, we define a certain family of simple $2 \times n$ transportation polytopes whose polars are not weakly k -decomposable for any fixed integer k .

2. Background and definitions

2.1. Simplicial complexes, decomposability, and diameter bounds. A *simplicial complex* Δ on vertex set V is a collection of subsets $\tau \subseteq V$ (called *faces*) such that if τ is a face of Δ and $\sigma \subseteq \tau$, then σ is also a face of Δ . The *dimension* of a face $\tau \in \Delta$ is $\dim \tau := |\tau| - 1$, and the *dimension* of Δ is $\dim \Delta := \max\{\dim \tau : \tau \in \Delta\}$. We say that Δ is *pure* if all of its *facets* (maximal faces under inclusion) have the same dimension.

We will borrow terminology from matroid complexes to define the *rank* of a subset $S \subseteq V$ to be $\text{rank}(S) := \max\{|\tau| : \tau \subseteq S, \tau \in \Delta\}$, and the *corank* of S to be $\text{corank}(S) := |S| - \text{rank}(S)$. Alternatively, the rank of S is one more than the dimension of the restriction of Δ to the vertices in S .

Let τ be a face of the simplicial complex Δ . The *deletion* of τ in Δ is the simplicial complex $\text{del}_\Delta(\tau) := \{\sigma \in \Delta : \tau \not\subseteq \sigma\}$. The *link* of τ in Δ is the simplicial complex $\text{lk}_\Delta(\tau) := \{\sigma \in \Delta : \sigma \cap \tau = \emptyset, \sigma \cup \tau \in \Delta\}$.

Given a pure simplicial complex Δ and facets $\sigma, \tau \in \Delta$, the *distance* from σ to τ is the length of the shortest path $\sigma = \rho_0, \rho_1, \dots, \rho_t = \tau$ where the ρ_i are facets of Δ and ρ_i intersects ρ_{i+1} along a *ridge* (a codimension-one face) for all $0 \leq i < t$. The *diameter* of a pure simplicial complex, denoted $\text{diam}(\Delta)$, is the maximum distance between any two facets in Δ .

One approach to establishing diameter bounds for pure simplicial complexes is to study decompositions that generalize the more well-known property of a shedding order in a natural way. Provan and Billera [PB80] defined a notion of (weak) k -decomposability for pure simplicial complexes and showed that (weakly) k -decomposable simplicial complexes satisfy natural diameter bounds.

Definition 2.1 ([PB80, Definition 2.1]). Let Δ be a $(d - 1)$ -dimensional simplicial complex and let $0 \leq k \leq d - 1$. We say that Δ is *k -decomposable* if it is pure and if either Δ is a $(d - 1)$ -simplex or there exists a face $\tau \in \Delta$ with $\dim(\tau) \leq k$ (called a *shedding* face) such that

- (i) the deletion $\text{del}_\Delta(\tau)$ is $(d - 1)$ -dimensional and k -decomposable, and
- (ii) the link $\text{lk}_\Delta(\tau)$ is $(d - |\tau| - 1)$ -dimensional and k -decomposable.

Theorem 2.2 ([PB80, Theorem 2.10]). *If Δ is a k -decomposable $(d-1)$ -dimensional simplicial complex, then*

$$\text{diam}(\Delta) \leq f_k(\Delta) - \frac{d}{k+1},$$

where $f_k(\Delta)$ denotes the number of k -dimensional faces in Δ .

Definition 2.3 ([PB80, Definition 4.2.1]). Let Δ be a $(d-1)$ -dimensional simplicial complex and let $0 \leq k \leq d-1$. We say that Δ is *weakly k -decomposable* if it is pure and if either Δ is a $(d-1)$ -simplex or there exists a face $\tau \in \Delta$ of dimension $\dim(\tau) \leq k$ (called a *shedding* face) such that the deletion $\text{del}_\Delta(\tau)$ is $(d-1)$ -dimensional and weakly k -decomposable.

Theorem 2.4 ([PB80, Theorem 4.2.3]). *If Δ is a weakly k -decomposable $(d-1)$ -dimensional simplicial complex, then*

$$\text{diam}(\Delta) \leq 2f_k(\Delta).$$

In particular, any 0-decomposable complex (also called *vertex-decomposable*) satisfies the Hirsch Conjecture while any weakly 0-decomposable complex (also called *weakly vertex-decomposable*) has diameter bounded by a linear function of its number of vertices. Klee and Kleinschmidt [KK87, Proposition 6.3] pointed out that Lockeberg [Loc77] had constructed a 4-dimensional polytope on 12 vertices that was not vertex-decomposable (but still satisfied the Hirsch bound), while the non-Hirsch polytopes of Santos [San12] and Matschke, Santos, and Weibel [MSW12] provide further examples of simplicial polytopes that are not vertex-decomposable. In a recent paper [DLK12], De Loera and the second author provided the first examples of simplicial polytopes that are not even weakly vertex-decomposable. Their smallest counterexample is a 4-dimensional polytope on 10 vertices and 30 facets, which can be seen as the polar of the intersection of the 5-dimensional cube $[-1, 1]^5$ with the linear hyperplane of equation $\sum_i x_i = 0$.

In the hopes of establishing a polynomial diameter bound, it is natural to ask whether there is a constant k such that all simplicial polytopes are (weakly) k -decomposable. We will answer this question in the negative by establishing that for any $k \geq 0$, there exist simplicial polytopes that are not weakly k -decomposable. Our construction is motivated by the counterexamples introduced by De Loera and Klee [DLK12], which arise as the polars of a certain family of simple transportation polytopes. The necessary background on transportation polytopes is briefly presented below.

2.2. Transportation polytopes. Transportation polytopes are classical polytopes from optimization. Here we recall their basic definitions and combinatorial properties and refer to classical surveys on the topic for more details and proofs [KW68, YKK84].

Definition 2.5. Fix two integers $m, n \geq 1$, and two vectors $\mathbf{m} \in \mathbb{R}^m$ and $\mathbf{n} \in \mathbb{R}^n$. The $m \times n$ *transportation polytope* $P(\mathbf{m}, \mathbf{n})$ is the collection of all non-negative matrices $X := (x_{\mu,\nu}) \in \mathbb{R}_{\geq 0}^{m \times n}$ such that

$$\forall \mu \in [m], \quad \sum_{\nu \in [n]} x_{\mu,\nu} = \mathbf{m}_\mu \quad \text{and} \quad \forall \nu \in [n], \quad \sum_{\mu \in [m]} x_{\mu,\nu} = \mathbf{n}_\nu.$$

The vectors \mathbf{m} and \mathbf{n} are called the *margins* of $P(\mathbf{m}, \mathbf{n})$.

Intuitively, a point in the transportation polytope $P(\mathbf{m}, \mathbf{n})$ is an assignment of weights to be transported between m sources and n sinks on the edges of the complete bipartite graph $K_{m,n}$ such that the total quantity that a source μ provides corresponds to its supply \mathbf{m}_μ while the total quantity that a sink ν receives corresponds to its demand \mathbf{n}_ν . The [support](#) of a point $X := (x_{\mu,\nu})$ of $P(\mathbf{m}, \mathbf{n})$ is the subgraph $\text{supp}(X) := \{(\mu, \nu) \in K_{m,n} : x_{\mu,\nu} > 0\}$ of the complete bipartite graph $K_{m,n}$. The following theorem summarizes classical properties of transportation polytopes. See [KW68, YKK84] for proofs.

Theorem 2.6. *The transportation polytope $P(\mathbf{m}, \mathbf{n})$, with margins $\mathbf{m} \in \mathbb{R}^m$ and $\mathbf{n} \in \mathbb{R}^n$, has the following properties.*

Feasibility: $P(\mathbf{m}, \mathbf{n})$ is non-empty if and only if $\sum_{\mu \in [m]} \mathbf{m}_\mu = \sum_{\nu \in [n]} \mathbf{n}_\nu$.

Dimension: The dimension of $P(\mathbf{m}, \mathbf{n})$ is $(m-1)(n-1)$, provided it is nonempty.

Non-degeneracy: $P(\mathbf{m}, \mathbf{n})$ is nondegenerate (hence simple) if and only if there are no proper subsets $\emptyset \subsetneq M \subsetneq [m]$ and $\emptyset \subsetneq N \subsetneq [n]$ such that $\sum_{\mu \in M} \mathbf{m}_\mu = \sum_{\nu \in N} \mathbf{n}_\nu$. The support of the points of $P(\mathbf{m}, \mathbf{n})$ are then connected and spanning.

Vertices: A point of $P(\mathbf{m}, \mathbf{n})$ is a vertex of $P(\mathbf{m}, \mathbf{n})$ if and only if its support is a forest (i.e. contains no cycle). In particular, if $P(\mathbf{m}, \mathbf{n})$ is nondegenerate, a point of $P(\mathbf{m}, \mathbf{n})$ is a vertex if and only if its support is a spanning tree of $K_{m,n}$.

Facets: All facets of $P(\mathbf{m}, \mathbf{n})$ are of the form $F_{\mu,\nu} := \{X \in P(\mathbf{m}, \mathbf{n}) : x_{\mu,\nu} = 0\}$ for some $\mu \in [m]$ and $\nu \in [n]$. Moreover, if $m, n \geq 4$, then the set $F_{\mu,\nu}$ is a facet if and only if $\mathbf{m}_\mu + \mathbf{n}_\nu < \sum_{\mu' \in [m]} \mathbf{m}_{\mu'} = \sum_{\nu' \in [n]} \mathbf{n}_{\nu'}$.

Optimizing transportation costs naturally gives rise to linear optimization problems on transportation polytopes, and thus leads to the question to evaluate the diameter of transportation polytopes. Although this question has been largely studied in the literature, the Hirsch Conjecture for transportation polytopes is still open. On the one hand, Brightwell, van den Heuvel, and Stougie [BvdHS06] showed that the diameter of any $m \times n$ transportation polytope is at most $8(m+n-1)$, i.e. eight times the Hirsch bound. On the other hand, the precise Hirsch bound was only proven for $2 \times n$ transportation polytopes by Kim [Kim10, Section 3.5] and for the family of signature polytopes, defined as follows by Balinski and Rispoli [BR93].

Definition 2.7 ([BR93]). Fix a vector $\mathbf{d} \in \mathbb{R}^m$ such that $\sum_{\mu \in [m]} \mathbf{d}_\mu = m+n-1$. Let $\mathcal{T}(\mathbf{d})$ denote the set of all spanning trees of the complete bipartite graph $K_{m,n}$ where each vertex $\mu \in [m]$ has degree \mathbf{d}_μ . An $m \times n$ transportation polytope $P(\mathbf{m}, \mathbf{n})$ is a [d-signature polytope](#) if any tree of $\mathcal{T}(\mathbf{d})$ is the support of a vertex of $P(\mathbf{m}, \mathbf{n})$.

Controlling the feasible trees of signature polytopes, Balinski and Rispoli [BR93] prove that these polytopes satisfy the Monotone Hirsch Conjecture. In particular, for any vector $\mathbf{d} \in \mathbb{R}^m$ such that $\sum_{\mu \in [m]} \mathbf{d}_\mu = m+n-1$, observe that the $m \times n$ transportation polytope $P(\mathbf{m}, \mathbf{n})$ with margins $\mathbf{m} := m\mathbf{d} - (m+1)\mathbf{1} \in \mathbb{R}^m$ and $\mathbf{n} := m\mathbf{1} \in \mathbb{R}^n$ is a signature polytope and thus satisfies the Monotone Hirsch Conjecture. These particular examples of signature polytopes were previously studied by Balinski [Bal74].

3. Simplicial polytopes that are not weakly k -decomposable

In this section, we study a certain family of simple $2 \times n$ transportation polytopes whose polars are not weakly k -decomposable for sufficiently large n .

Definition 3.1. Fix integers $a, b \geq 1$ and define $\Delta(a, b)$ to be the boundary complex of the polar polytope to the $2 \times (a + b + 1)$ transportation polytope $P(\mathbf{m}, \mathbf{n})$ with margins

$$\mathbf{m} := (2a + 1, 2b + 1) \in \mathbb{R}^2 \quad \text{and} \quad \mathbf{n} := (2, 2, \dots, 2) \in \mathbb{R}^{a+b+1}.$$

According to Theorem 2.6, $P(\mathbf{m}, \mathbf{n})$ is a simple polytope of dimension $a + b$ whose facets are precisely the sets $F_{\mu, \nu} := \{X \in P(\mathbf{m}, \mathbf{n}) : x_{\mu, \nu} = 0\}$, for $\mu \in [2]$ and $\nu \in [a + b + 1]$. For simplicity, we decompose the vertex set of $\Delta(a, b)$ into sets $U := \{u_1, \dots, u_{a+b+1}\}$ and $V := \{v_1, \dots, v_{a+b+1}\}$ where u_ν is the polar vertex to the facet $F_{1, \nu}$ and v_ν is the polar vertex to the facet $F_{2, \nu}$. That is to say, the vertices of U correspond to the coordinates of the top row of the $2 \times n$ contingency table defining $P(\mathbf{m}, \mathbf{n})$ and the vertices of V correspond to the coordinates of the bottom row. Theorem 2.6 yields the following description of the facets of $\Delta(a, b)$.

Lemma 3.2. *The facets of $\Delta(a, b)$ are precisely the sets $A \cup B$ where $A \subseteq V$, $B \subseteq U$, $|A| = a$, $|B| = b$, and $A \cup B$ contains at most one element from each pair $\{u_\nu, v_\nu\}$.*

Proof. Consider a facet F of $\Delta(a, b)$. Let $X := (x_{\mu, \nu})$ be the vertex of $P(a, b)$ polar to F , let $A := F \cap V$, and let $B := F \cap U$. If $v_\nu \in A$, then $x_{2, \nu} = 0$, and thus $x_{1, \nu} = 2$. Consequently, $2a + 1 = \sum_{\nu} x_{1, \nu} \geq 2|A|$. Similarly $2b + 1 \geq 2|B|$. Since $|A| + |B| = a + b$, we obtain that $|A| = a$ and $|B| = b$. Finally, F cannot contain both u_ν and v_ν since X cannot have $x_{1, \nu} = x_{2, \nu} = 0$.

Conversely, let $A \subseteq V$ and $B \subseteq U$ be such that $|A| = a$, $|B| = b$, and $A \cup B$ contains at most one element from each pair $\{u_\nu, v_\nu\}$. Let $\bar{B} := \{\nu : u_\nu \in B\}$ and $\bar{A} := \{\nu : v_\nu \in A\}$. Then \bar{A} and \bar{B} are disjoint subsets of $[a + b + 1]$ and their complement $[a + b + 1] \setminus (\bar{A} \cup \bar{B})$ has a unique element $\bar{\nu}$. Define $X := (x_{\mu, \nu})$ by

$$x_{1, \nu} := \begin{cases} \geq 1 & \text{if } \nu = \bar{\nu} \\ 2 & \text{if } \nu \in \bar{A} \\ 0 & \text{if } \nu \in \bar{B} \end{cases} \quad \text{and} \quad x_{2, \nu} := \begin{cases} \geq 1 & \text{if } \nu = \bar{\nu} \\ 0 & \text{if } \nu \in \bar{A} \\ 2 & \text{if } \nu \in \bar{B} \end{cases}.$$

The point X is a vertex of $P(\mathbf{m}, \mathbf{n})$ since it is a point of $P(\mathbf{m}, \mathbf{n})$ whose support is a spanning tree. Moreover, its polar facet in $\Delta(a, b)$ is $A \cup B$. \square

Remark 3.3. We can alternatively describe the transportation polytope $P(\mathbf{m}, \mathbf{n})$ of Definition 3.1 as the intersection of the $(a+b+1)$ -dimensional cube $C := [0, 2]^{a+b+1}$ with the hyperplane H of equation $\sum_i x_i = 2a + 1$. The $2(a + b + 1)$ facets of P are supported by the hyperplanes supporting the facets of the cube C . The $\frac{(a+b+1)!}{a!b!}$ vertices of P are the intersections of the hyperplane H with the edges of the cube C at distance $2a$ from the origin. Equivalently, P is the Minkowski sum of the hypersimplices $\Delta_a + \Delta_{a+1}$, where the hypersimplex Δ_a is the intersection of the cube $[0, 1]^{a+b+1}$ with the hyperplane $\sum_i x_i = a$.

According to the above-mentioned results on $2 \times n$ transportation polytopes [Kim10, Section 3.5] and on signature polytopes [Bal74, BR93], the transportation polytope $P(\mathbf{m}, \mathbf{n})$ satisfies the (Monotone) Hirsch Conjecture. In contrast, the

results of [DLK12, Theorem 3.1] imply that the boundary complex $\Delta(a, b)$ of the polar of $P(\mathbf{m}, \mathbf{n})$ is not weakly vertex-decomposable when $a, b \geq 2$. We now refine the analysis of the weak decompositions of $\Delta(a, b)$ to derive the following theorem.

Theorem 3.4. *If $\frac{k+3}{2} \leq \min(a, b)$, then the simplicial complex $\Delta(a, b)$ of Definition 3.1 is not weakly k -decomposable.*

The following immediate consequence of Theorem 3.4 shows that there is no hope of using (weak) k -decomposability as a means of proving the Polynomial Hirsch Conjecture.

Corollary 3.5. *There exist d -dimensional simplicial polytopes whose boundary complexes are not (weakly) $O(\sqrt{d})$ -decomposable.*

To prove Theorem 3.4, we will suppose by way of contradiction that $\Delta(a, b)$ is weakly k -decomposable with shedding sequence $\tau_1, \tau_2, \dots, \tau_t$. We set $\Delta_0 := \Delta(a, b)$ and $\Delta_i := \text{del}_{\Delta_{i-1}}(\tau_i)$, for $i \geq 1$. Each Δ_i is a pure simplicial complex of dimension $a+b-1$ and Δ_t consists of a single $(a+b-1)$ -simplex. Our goal now is to understand how shedding each of the faces τ_i affects certain subsets of vertices of $\Delta(a, b)$. With this motivation, we define a family of functions

$$\varphi_i : \begin{array}{c} U \\ \leq b+1 \end{array} \cup \begin{array}{c} V \\ \leq a+1 \end{array} \rightarrow \mathbb{N},$$

by setting $\varphi_i(S)$ to be the corank of the set of vertices S in the complex Δ_i , i.e.

$$\varphi_i(S) := |S| - \max\{|\tau| : \tau \subseteq S, \tau \in \Delta_i\}.$$

Here we use the notation $\overset{W}{\leq w}$ to denote the collection of subsets of the set W of size at most w . We will need the following properties of the functions φ_i .

Lemma 3.6. *The family of functions φ_i satisfy the following properties:*

- (1) $\varphi_0(S) \leq 1$ for all $S \in \overset{U}{\leq b+1} \cup \overset{V}{\leq a+1}$.
- (2) There exists a set S such that $\varphi_t(S) \geq 2$.
- (3) If $\tau_i \cap U \neq \emptyset$ and $\tau_i \cap V \neq \emptyset$, then $\varphi_i(S) = \varphi_{i-1}(S)$ for all S .
- (4) If $\tau_i \subseteq U$, then $\varphi_i(S) = \varphi_{i-1}(S)$ for all $S \in \overset{V}{\leq a+1}$.
- (5) $\varphi_i(S) \leq 1$ if and only if there is some $v \in S$ such that $S \setminus v \in \Delta_i$.

Proof. Since each b -element subset of U and each a -element subset of V spans a face of $\Delta_0 = \Delta(a, b)$, it follows that $\varphi_0(S) \leq 1$ for all $S \in \overset{U}{\leq b+1} \cup \overset{V}{\leq a+1}$. This proves part (1).

For part (2), suppose Δ_t consists of a simplex on vertex set $A \cup B$ with $A \subseteq V$, $B \subseteq U$, $|A| = a$, and $|B| = b$. Note that $|U \setminus B| = a+1$, and let S be a subset of $U \setminus B$ of size $\min\{a+1, b+1\}$ so that $S \in \overset{U}{\leq b+1}$. Since no vertex of $U \setminus B$ belongs to Δ_t , it follows that $\varphi_t(S) = |S| - 0 \geq \min\{a+1, b+1\} \geq 2$.

For part (3), assume that $\tau_i \cap U \neq \emptyset$ and $\tau_i \cap V \neq \emptyset$ and let $S \in \overset{U}{\leq b+1} \cup \overset{V}{\leq a+1}$ be an arbitrary set. We will show that $\{\tau : \tau \subseteq S, \tau \in \Delta_i\} = \{\tau : \tau \subseteq S, \tau \in \Delta_{i-1}\}$. Since Δ_i is a subcomplex of Δ_{i-1} , we only need to show that the former set contains the latter. Let $\tau \subseteq S$ be a face of Δ_{i-1} . Since τ_i contains vertices from both U and V and τ is a subset of either U or V , τ does not contain τ_i . Thus τ is a face of Δ_i as well.

The proof of part (4) is identical to that of part (3). By symmetry, if $\tau_i \subseteq V$, then $\varphi_i(S) = \varphi_{i-1}(S)$ for all $S \in \overset{U}{\leq b+1}$.

Finally, part (5) follows from the fact that we have $\varphi_i(S) \leq 1$ if and only if $\max\{|\tau| : \tau \subseteq S, \tau \in \Delta_i\} \geq |S| - 1$. \square

The following lemma will be essential to the proof of Theorem 3.4.

Lemma 3.7 \dashv Let \mathcal{X} be a collection of sets, each of which has size at most $k + 1$, and suppose $\bigcap_{X \in \mathcal{X}} X = \emptyset$. Then there is a sub-collection $\mathcal{Y} \subseteq \mathcal{X}$ such that

$$\bigcap_{X \in \mathcal{Y}} X = \emptyset \quad \text{and} \quad \left[\bigcap_{X \in \mathcal{Y}} X \leq \frac{k+3}{2} \right]^2.$$

Proof. Let \mathcal{Y} be a minimal collection of sets of \mathcal{X} such that $\bigcap_{X \in \mathcal{Y}} X = \emptyset$ (i.e. any proper sub-collection of sets in \mathcal{Y} has a non-empty intersection).

For each $X \in \mathcal{Y}$, there is an element $f(X) \in \bigcap_{Y \in \mathcal{Y} \setminus \{X\}} Y$ by our assumption that \mathcal{Y} is minimal. Furthermore, $f(X) \notin X$, since otherwise $\bigcap_{Y \in \mathcal{Y}} Y \neq \emptyset$. Hence the elements $\{f(X) : X \in \mathcal{Y}\}$ are all distinct. This means any $X \in \mathcal{Y}$ contains each of the elements $f(Y)$ with $Y \in \mathcal{Y} \setminus \{X\}$ and at most $k + 2 - |\mathcal{Y}|$ other elements. Thus

$$\left[\bigcap_{X \in \mathcal{Y}} X \leq |\mathcal{Y}| + |\mathcal{Y}|(k + 2 - |\mathcal{Y}|) = |\mathcal{Y}|(k + 3 - |\mathcal{Y}|) \leq \frac{k+3}{2} \right]^2,$$

since the quadratic function $y \mapsto y(k + 3 - y)$ is maximized when $y = \frac{k+3}{2}$. \square

Remark 3.8. Observe that the bound in the previous lemma is tight. Consider all $\lfloor \frac{k+1}{2} \rfloor$ -element subsets of a $(\lfloor \frac{k+1}{2} \rfloor + 1)$ -element set. Add to each of these sets $\lceil \frac{k+1}{2} \rceil$ new elements (all additional elements are distinct). The intersection of the resulting $\lfloor \frac{k+1}{2} \rfloor + 1$ sets is empty, but the intersection of any proper sub-collection is non-empty. Moreover, their union has cardinality $(\lfloor \frac{k+1}{2} \rfloor + 1)(\lceil \frac{k+1}{2} \rceil + 1)$.

We are now ready to prove Theorem 3.4.

Proof of Theorem 3.4. By Lemmas 3.6(1) and 3.6(2), we can consider the smallest value $i \in [t]$ for which there exists some S with $\varphi_i(S) \geq 2$. By Lemma 3.6(3), the corresponding shedding face τ_i is a subset of either U or V . Without loss of generality, we may assume that $\tau_i \subseteq U$.

Let \mathcal{X} denote the collection of shedding faces $\tau \in \{\tau_1, \dots, \tau_i\}$ such that $\tau \subseteq S$. Observe that $\bigcap_{\tau \in \mathcal{X}} \tau = \emptyset$. Indeed, otherwise if there is an element $u \in \bigcap_{\tau \in \mathcal{X}} \tau$, then $S \setminus u$ does not contain any set $\tau \in \mathcal{X}$. This would mean that $S \setminus u \in \Delta_i$ and hence $\varphi_i(S) \leq 1$ by Lemma 3.6(5). According to Lemma 3.7, there exists a sub-collection $\mathcal{Y} \subseteq \mathcal{X}$ of shedding faces such that

$$\bigcap_{\tau \in \mathcal{Y}} \tau = \emptyset \quad \text{and} \quad \left[\bigcap_{\tau \in \mathcal{Y}} \tau \leq \frac{k+3}{2} \right]^2 \leq b.$$

Replacing S by $\bigcap_{\tau \in \mathcal{Y}} \tau$, we can thus assume that $\varphi_i(S) \geq 2$ and $|S| \leq b$.

Since $|S| \leq b$, we can choose a subset $T \subseteq V$ of $a + 1$ vertices of $\Delta(a, b)$ such that $S \cup T$ contains at most one element from each pair $\{u_j, v_j\}$. By our choice of i and Lemma 3.6(4), $\varphi_i(T) = \varphi_{i-1}(T) \leq 1$. Thus there is a subset $A \subseteq T$ with $|A| = a$ such that $A \in \Delta_i$.

We claim that A is not contained in any face of size $a + b$ in Δ_i , which will contradict our assumption that Δ_i is pure of dimension $a + b - 1$. Suppose that there is a subset $B \subseteq U$ with $|B| = b$ such that $A \cup B$ is a facet of Δ_i . Then $B \cap S$ is a face of Δ_i and $|B \cap S| \geq |S| - 1$, meaning $\varphi_i(S) \leq 1$, which contradicts our choice of S . \square

Acknowledgments

We are grateful to Jesús De Loera and Paco Santos for a number of helpful and insightful conversations during the cultivation of the ideas that motivated this paper. We discovered the main ideas behind this paper during the Triangulations workshop at the Oberwolfach MFO during May 2012. We would like to extend our sincere thanks to the organizers – William Jaco, Frank Lutz, Paco Santos, and John Sullivan – for the invitation to attend the workshop. Nicolai Hähnle and Steven Klee also thank the Oberwolfach Leibniz Grant for providing travel support for the workshop. We also thank an anonymous referee for helpful suggestions.

References

- [Bal74] M. Balinski. On two special classes of transportation polytopes. *Math. Programming Stud.*, 1:43–58, 1974. Pivoting and extensions.
- [Bar74] D. Barnette. An upper bound for the diameter of a polytope. *Discrete Math.*, 10:9–13, 1974.
- [BR93] M. Balinski and F. Rispoli. Signature classes of transportation polytopes. *Math. Programming*, 60(2, Ser. A):127–144, 1993.
- [BvdHS06] G. Brightwell, J. van den Heuvel, and L. Stougie. A linear bound on the diameter of the transportation polytope. *Combinatorica*, 26(2):133–139, 2006.
- [DL11] J. A. De Loera. New insights into the complexity and geometry of linear optimization. *Optima, newsletter of the Mathematical Optimization Society*, 87:1–13, 2011.
- [DLK12] J. De Loera and S. Klee. Transportation problems and simplicial polytopes that are not weakly vertex-decomposable. To appear in *Math. Oper. Res.*. Preprint [arxiv:1203.1676](https://arxiv.org/abs/1203.1676), 2012.
- [DTZ09] A. Deza, T. Terlaky, and Y. Zinchenko. A continuous d -step conjecture for polytopes. *Discrete Comput. Geom.*, 41(2):318–327, 2009.
- [EHR10] F. Eisenbrand, N. Hähnle, A. Razborov, and T. Rothvoß. Diameter of polyhedra: limits of abstraction. *Math. Oper. Res.*, 35(4):786–794, 2010.
- [Kal10] G. Kalai. Polymath 3: The Polynomial Hirsch Conjecture, 2010. <http://gilkalai.wordpress.com>.
- [Kim10] E. D. Kim. *Geometric Combinatorics of Transportation Polytopes and the Behavior of the Simplex Method*. PhD thesis, University of California, Davis, 2010.
- [KK87] V. Klee and P. Kleinschmidt. The d -step conjecture and its relatives. *Math. Oper. Res.*, 12(4):718–755, 1987.
- [KK92] G. Kalai and D. Kleitman. A quasi-polynomial bound for the diameter of graphs of polyhedra. *Bull. Amer. Math. Soc. (N.S.)*, 26(2):315–316, 1992.
- [KS10] E. D. Kim and F. Santos. An update on the Hirsch conjecture. *Jahresber. Dtsch. Math.-Ver.*, 112(2):73–98, 2010.
- [KW68] V. Klee and C. Witzgall. Facets and vertices of transportation polytopes. In *Mathematics of the Decision Sciences, Part I (Seminar, Stanford, Calif., 1967)*, pages 257–282. Amer. Math. Soc., Providence, R.I., 1968.
- [Lar70] D. Larman. Paths of polytopes. *Proc. London Math. Soc. (3)*, 20:161–178, 1970.
- [Loc77] E. R. Lockeberg. *Refinements in boundary complexes of polytopes*. PhD thesis, University College London, 1977.
- [MSW12] B. Matschke, F. Santos, and C. Weibel. The width of 5-dimensional prisms. Preprint [arxiv:1202.4701](https://arxiv.org/abs/1202.4701), 2012.
- [PB80] S. Provan and L. Billera. Decompositions of simplicial complexes related to diameters of convex polyhedra. *Math. Oper. Res.*, 5(4):576–594, 1980.
- [San12] F. Santos. A counter-example to the Hirsch conjecture. *Ann. of Math. (2)*, 176:383–412, 2012.
- [Tod02] M. J. Todd. The many facets of linear programming. *Math. Program.*, 91(3, Ser. B):417–449(Hr9Pr)55TJ -21iMP499(Hr9Pr0,i.)-354(P)2(Part)-354rtSanB.
- [Kr74]

(NH) Technische Universität Berlin
E-mail address: haehnle@math.tu-berlin.de

(SK) Seattle University
E-mail address: klees@seattleu.edu
URL: <http://fac-staff.seattleu.edu/klees/web/>

(VP) CNRS & LIX, Ecole Polytechnique, Palaiseau
E-mail address: vincent.pilaud@lix.polytechnique.fr
URL: <http://www.lix.polytechnique.fr/~pilaud/>