

Fonctions discrètes harmoniques dans des cônes

KILIAN RASCHEL



Séminaire de Combinatoire du LIX
28 septembre 2016
Polytechnique

Introduction & motivations


Applications in probability theory

Applications in combinatorics

Discrete harmonic functions in the quadrant


Introductory example & definition

Absorption probabilities for the SRW on \mathbb{N}



Introductory example & definition

Absorption probabilities for the SRW on \mathbb{N}




Probability $a(i) := \mathbb{P}_i[\exists n \geq 0 : \text{SRW } S(n) = 0]$ satisfies

▷ $a(0) = 1 \rightsquigarrow$ *initial condition*

▷ $a(i) = p \cdot a(i+1) + (1-p) \cdot a(i-1) \rightsquigarrow$ *recurrence*

Introductory example & definition

Absorption probabilities for the SRW on \mathbb{N}



Probability $a(i) := \mathbb{P}_i[\exists n \geq 0 : \text{SRW } S(n) = 0]$ satisfies


▷ $a(0) = 1 \rightsquigarrow$ *initial condition*

▷ $a(i) = p \cdot a(i+1) + (1-p) \cdot a(i-1) \rightsquigarrow$ *recurrence*

$$\text{Solution } a(i) = \begin{cases} 1 & \text{if } p \leq \frac{1}{2} \\ \left(\frac{1-p}{p}\right)^i & \text{if } p > \frac{1}{2} \end{cases}$$

Introductory example & definition

Absorption probabilities for the SRW on \mathbb{N}



Probability $a(i) := \mathbb{P}_i[\exists n \geq 0 : \text{SRW } S(n) = 0]$ satisfies

▷ $a(0) = 1 \rightsquigarrow$ *initial condition*

▷ $a(i) = p \cdot a(i+1) + (1-p) \cdot a(i-1) \rightsquigarrow$ *recurrence*

$$\text{Solution } a(i) = \begin{cases} 1 & \text{if } p \leq \frac{1}{2} \\ \left(\frac{1-p}{p}\right)^i & \text{if } p > \frac{1}{2} \end{cases}$$


Definition: f harmonic if $L[f](x) = 0$ for all x in a region $\subset \mathbb{Z}^d$

$$L[f](x) = \sum_{y \in N} p(y) \{f(x+y) - f(x)\},$$

with *set of neighbors* $N \subset \mathbb{Z}^d$ and *weights* $p = \{p(y)\}_{y \in \mathbb{Z}^d}$

Introductory example & definition

Absorption probabilities for the SRW on \mathbb{N}



Probability $a(i) := \mathbb{P}_i[\exists n \geq 0 : \text{SRW } S(n) = 0]$ satisfies

▷ $a(0) = 1 \rightsquigarrow$ *initial condition*

▷ $a(i) = p \cdot a(i+1) + (1-p) \cdot a(i-1) \rightsquigarrow$ *recurrence*

$$\text{Solution } a(i) = \begin{cases} 1 & \text{if } p \leq \frac{1}{2} \\ \left(\frac{1-p}{p}\right)^i & \text{if } p > \frac{1}{2} \end{cases}$$

Definition: f harmonic if $L[f](x) = 0$ for all x in a region $C \subset \mathbb{Z}^d$

$$L[f](x) = \sum_{y \in N} p(y) \{f(x+y) - f(x)\},$$

with *set of neighbors* $N \subset \mathbb{Z}^d$ and *weights* $p = \{p(y)\}_{y \in \mathbb{Z}^d}$

▷ Multivariate linear recurrences with constant coefficients

{History of/Questions on} preharmonic functions (1/2)

Classical (continuous) harmonic functions in \mathbb{R}^d

$$\Delta[f](x) = \sum_{i=1}^d \frac{\partial^2 f(x)}{\partial x_i^2} = 0$$

- ▷ Possibility of adding *weights* \leadsto *elliptic operators*
- ▷ *Harmonic functions satisfy various properties*: maximum principle/mean value property/Harnack inequalities/Liouville's theorem/relations with analytic functions/etc.
- ▷ *Examples of application*: Heat equation/Dirichlet problem/Poisson's equation/more general PDEs/etc.





{History of/Questions on} preharmonic functions (1/2)

Classical (continuous) harmonic functions in \mathbb{R}^d

$$\Delta[f](x) = \sum_{i=1}^d \frac{\partial^2 f(x)}{\partial x_i^2} = 0$$




- ▷ Possibility of adding *weights* \leadsto *elliptic operators*
- ▷ *Harmonic functions satisfy various properties*: maximum principle/mean value property/Harnack inequalities/Liouville's theorem/relations with analytic functions/etc.
- ▷ *Examples of application*: Heat equation/Dirichlet problem/Poisson's equation/more general PDEs/etc.

Do preharmonic functions satisfy similar properties?

- ▷ Dirichlet problem  Phillips & Wiener '23; Bouligand '25
- ▷ Harnack inequalities  Lawler & Polaski '92; Varopoulos '99
- ▷ Maximum principle, Liouville's theorem & related topics  Heilbronn '48
- ▷ Cauchy-Riemann equations  Duffin '55; Kiselman '05-'08




{History of/Questions on} preharmonic functions (2/2)

Further properties

- ▷ Rate of growth  Murdoch '63-'65; Ignatiuk-Robert '10
- ▷ Picard's theorem (sign of harmonic functions) & factorization  Murdoch '63-'65
- ▷ Absolute monotonicity  Lippner & Mangoubi '15

{History of/Questions on} preharmonic functions (2/2)

Further properties




- ▷ Rate of growth  Murdoch '63-'65; Ignatiuk-Robert '10
- ▷ Picard's theorem (sign of harmonic functions) & factorization  Murdoch '63-'65
- ▷ Absolute monotonicity  Lippner & Mangoubi '15

Preharmonic & harmonic functions

- ▷ Relations between discrete & continuous harmonic functions  Lusternik '26; Ferrand '44; Kesten '91; Varopoulos '09

{History of/Questions on} preharmonic functions (2/2)



Further properties

- ▷ Rate of growth  Murdoch '63-'65; Ignatiuk-Robert '10
- ▷ Picard's theorem (sign of harmonic functions) & factorization  Murdoch '63-'65
- ▷ Absolute monotonicity  Lippner & Mangoubi '15

Preharmonic & harmonic functions




- ▷ Relations between discrete & continuous harmonic functions  Lusternik '26; Ferrand '44; Kesten '91; Varopoulos '09

Probability theory models

- ▷ Ising models  Mercat '01; Smirnov '10
- ▷ Conformal invariance of lattice models  Duminil-Copin & Smirnov '12

{History of/Questions on} preharmonic functions (2/2)



Further properties

- ▷ Rate of growth  Murdoch '63–'65; Ignatiuk-Robert '10
- ▷ Picard's theorem (sign of harmonic functions) & factorization  Murdoch '63–'65
- ▷ Absolute monotonicity  Lippner & Mangoubi '15

Preharmonic & harmonic functions

- ▷ Relations between discrete & continuous harmonic functions  Lusternik '26; Ferrand '44; Kesten '91; Varopoulos '09

Probability theory models




- ▷ Ising models  Mercat '01; Smirnov '10
- ▷ Conformal invariance of lattice models  Duminil-Copin & Smirnov '12

Special discrete functions

- ▷ Conformal mappings  Ferrand '44; Isaacs '52
- ▷ Discrete harmonic polynomials & discrete exponential functions  Terracini '45–'46; Heilbronn '48; Isaacs '52; Duffin '55; Duffin & Peterson '68

{History of/Questions on} preharmonic functions (2/2)



Further properties

- ▷ Rate of growth  Murdoch '63–'65; Ignatiuk-Robert '10
- ▷ Picard's theorem (sign of harmonic functions) & factorization  Murdoch '63–'65
- ▷ Absolute monotonicity  Lippner & Mangoubi '15

Preharmonic & harmonic functions

- ▷ Relations between discrete & continuous harmonic functions  Lusternik '26; Ferrand '44; Kesten '91; Varopoulos '09

Probability theory models

- ▷ Ising models  Mercat '01; Smirnov '10
- ▷ Conformal invariance of lattice models  Duminil-Copin & Smirnov '12

Special discrete functions


- ▷ Conformal mappings  Ferrand '44; Isaacs '52
- ▷ Discrete harmonic polynomials & discrete exponential functions  Terracini '45–'46; Heilbronn '48; Isaacs '52; Duffin '55; Duffin & Peterson '68

Potential theory

- ▷ Martin boundary  Woess '92; Kurkova & Malyshev '98; Ignatiuk-Robert & Loree '10; Mustapha '15

Warning: lattice walk enum. vs. preharmonic functions


Multivariate recurrence relations in both cases



- ▷ $q(n; i, j) = \#_{\mathbb{N}^2} \{(0, 0) \xrightarrow{n} (i, j)\}$
- ▷ $q(n+1; i, j) =$
 $q(n; i-1, j) + q(n; i+1, j) + q(n; i, j-1) + q(n; i, j+1)$
(Caloric functions)

Warning: lattice walk enum. vs. preharmonic functions

Multivariate recurrence relations in both cases




- ▷ $q(n; i, j) = \#_{\mathbb{N}^2}\{(0, 0) \xrightarrow{n} (i, j)\}$
- ▷ $q(n+1; i, j) = q(n; i-1, j) + q(n; i+1, j) + q(n; i, j-1) + q(n; i, j+1)$
(Caloric functions)
- ▷ $f(i, j) = \frac{1}{4}\{f(i-1, j) + f(i+1, j) + f(i, j-1) + f(i, j+1)\}$
(Preharmonic functions)

Main differences & difficulties

- ▷ A unique solution vs. an unknown ($\leq \infty$) number of solutions
- ▷ Consequence: *guess and prove* techniques do not work

Warning: lattice walk enum. vs. preharmonic functions

Multivariate recurrence relations in both cases



- ▷ $q(n; i, j) = \#_{\mathbb{N}^2} \{(0, 0) \xrightarrow{n} (i, j)\}$
- ▷ $q(n+1; i, j) = q(n; i-1, j) + q(n; i+1, j) + q(n; i, j-1) + q(n; i, j+1)$
(Caloric functions)
- ▷ $f(i, j) = \frac{1}{4} \{f(i-1, j) + f(i+1, j) + f(i, j-1) + f(i, j+1)\}$
(Preharmonic functions)

Main differences & difficulties


- ▷ A unique solution vs. an unknown ($\leq \infty$) number of solutions
- ▷ Consequence: *guess and prove* techniques do not work
- ▷ Generating functions of preharmonic functions satisfy *kernel functional equations*
- ▷ Preharmonic functions \approx homogenized enumeration problem:

$$K(x, y)Q(x, y) = K(x, 0)Q(x, 0) + K(0, y)Q(0, y) - K(0, 0)Q(0, 0) - xy$$

$$K'(x, y)F(x, y) = K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0)$$

Warning: lattice walk enum. vs. preharmonic functions

Multivariate recurrence relations in both cases



- ▷ $q(n; i, j) = \#_{\mathbb{N}^2} \{(0, 0) \xrightarrow{n} (i, j)\}$
- ▷ $q(n+1; i, j) = q(n; i-1, j) + q(n; i+1, j) + q(n; i, j-1) + q(n; i, j+1)$
(Caloric functions)
- ▷ $f(i, j) = \frac{1}{4} \{f(i-1, j) + f(i+1, j) + f(i, j-1) + f(i, j+1)\}$
(Preharmonic functions)

Main differences & difficulties

- ▷ A unique solution vs. an unknown ($\leq \infty$) number of solutions
- ▷ Consequence: *guess and prove* techniques do not work
- ▷ Generating functions of preharmonic functions satisfy *kernel functional equations*
- ▷ Preharmonic functions \approx homogenized enumeration problem:
$$K(x, y)Q(x, y) = K(x, 0)Q(x, 0) + K(0, y)Q(0, y) - K(0, 0)Q(0, 0) - xy$$
$$K'(x, y)F(x, y) = K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0)$$
- ▷ Preharmonic functions \rightsquigarrow counting numbers asymptotics

Introduction & motivations


Applications in probability theory

Applications in combinatorics

Discrete harmonic functions in the quadrant


Absorption probabilities and statistical mechanics

Markov chains: example



Absorption probabilities and statistical mechanics

Markov chains: example




$$f_i = \mathbb{P}_i[\text{hit } 4] \text{ satisfies } \begin{cases} f_1 = 0 \\ f_4 = 1 \\ f_2 = \frac{1}{2}f_1 + \frac{1}{2}f_3 \\ f_3 = \frac{1}{2}f_2 + \frac{1}{2}f_4 \end{cases}$$

$$\text{Solution: } \boxed{f_1 = 0, f_2 = \frac{1}{3}, f_3 = \frac{2}{3}, f_4 = 1}$$

Absorption probabilities and statistical mechanics

Markov chains: example



$$f_i = \mathbb{P}_i[\text{hit } 4] \text{ satisfies } \begin{cases} f_1 = 0 \\ f_4 = 1 \\ f_2 = \frac{1}{2}f_1 + \frac{1}{2}f_3 \\ f_3 = \frac{1}{2}f_2 + \frac{1}{2}f_4 \end{cases}$$


$$\text{Solution: } f_1 = 0, f_2 = \frac{1}{3}, f_3 = \frac{2}{3}, f_4 = 1$$

Markov chains: general theorem

The hitting probabilities are characterized as being the *minimal non-negative solutions* to a system of *linear recurrences*.

Absorption probabilities and statistical mechanics

Markov chains: example




$$f_i = \mathbb{P}_i[\text{hit } 4] \text{ satisfies } \begin{cases} f_1 = 0 \\ f_4 = 1 \\ f_2 = \frac{1}{2}f_1 + \frac{1}{2}f_3 \\ f_3 = \frac{1}{2}f_2 + \frac{1}{2}f_4 \end{cases}$$

$$\text{Solution: } f_1 = 0, f_2 = \frac{1}{3}, f_3 = \frac{2}{3}, f_4 = 1$$

Markov chains: general theorem

The hitting probabilities are characterized as being the *minimal non-negative solutions* to a system of *linear recurrences*.

Ising model




Discrete analyticity and convergence of the Fermionic observable

Smirnov '10


Doob transform

(1/2)

Example: construct a 1D process conditioned to stay in \mathbb{N}




Example: construct a 1D process conditioned to stay in \mathbb{N}




- ▷ Function $f(i) = i$ is positive harmonic and $f(0) = 0$
- ▷ Replace weights $p(i, i \pm 1) = \frac{1}{2}$ by $p^f(i, i \pm 1) = \frac{1}{2} \frac{f(i \pm 1)}{f(i)}$

Example: construct a 1D process conditioned to stay in \mathbb{N}




- ▷ Function $f(i) = i$ is positive harmonic and $f(0) = 0$
- ▷ Replace weights $p(i, i \pm 1) = \frac{1}{2}$ by $p^f(i, i \pm 1) = \frac{1}{2} \frac{f(i \pm 1)}{f(i)}$
- ▷ New weights sum to 1: $f(i-1) + f(i+1) = 2f(i)$




- ▷ Discrete Bessel process

 Biane '90; Mishchenko '05

Example: construct a 1D process conditioned to stay in \mathbb{N}



- ▷ Function $f(i) = i$ is positive harmonic and $f(0) = 0$
- ▷ Replace weights $p(i, i \pm 1) = \frac{1}{2}$ by $p^f(i, i \pm 1) = \frac{1}{2} \frac{f(i \pm 1)}{f(i)}$
- ▷ New weights sum to 1: $f(i-1) + f(i+1) = 2f(i)$



- ▷ Discrete Bessel process 🔗 Biane '90; Mishchenko '05

Construction can be generalized

- ▷ *Random processes* conditioned never to leave *cones* of \mathbb{Z}^d
- ▷ Quantum random walks, eigenvalues of random matrices, non-colliding random walks, etc.

🔗 Dyson '62; Biane '90-'92; Eichelsbacher & König '08

A second way for the conditioning (with the first exit time)

▷ RW $\{S(n)\}_{n \geq 0}$; *first exit time* $\tau = \inf\{n \geq 0 : S(n) = 0\}$

A second way for the conditioning (with the first exit time)

- ▷ RW $\{S(n)\}_{n \geq 0}$; *first exit time* $\tau = \inf\{n \geq 0 : S(n) = 0\}$
- ▷ On the event $\{\tau = \infty\}$, the RW stays in $\{1, 2, 3, \dots\}$

A second way for the conditioning (with the first exit time)

- ▷ RW $\{S(n)\}_{n \geq 0}$; *first exit time* $\tau = \inf\{n \geq 0 : S(n) = 0\}$
- ▷ On the event $\{\tau = \infty\}$, the RW stays in $\{1, 2, 3, \dots\}$
- ▷ Replace $\mathbb{P}[\cdot]$ by $\mathbb{P}[\cdot | \{\tau = \infty\}]$ to obtain a *conditioned RW*


A second way for the conditioning (with the first exit time)

- ▷ RW $\{S(n)\}_{n \geq 0}$; *first exit time* $\tau = \inf\{n \geq 0 : S(n) = 0\}$
- ▷ On the event $\{\tau = \infty\}$, the RW stays in $\{1, 2, 3, \dots\}$
- ▷ Replace $\mathbb{P}[\cdot]$ by $\mathbb{P}[\cdot | \{\tau = \infty\}]$ to obtain a *conditioned RW*
- ▷ Important question: do we have $\mathbb{P}[\cdot | \{\tau = \infty\}] = \mathbb{P}^f$?
 - 📖 Denisov & Wachtel '15 (general cones with zero drift); Courtiel, Melczer, Mishna & R. '16 (Gouyou-Beauchamps model with drift)
- ▷ Difficulty: exit time τ

A second way for the conditioning (with the first exit time)

- ▷ RW $\{S(n)\}_{n \geq 0}$; *first exit time* $\tau = \inf\{n \geq 0 : S(n) = 0\}$
- ▷ On the event $\{\tau = \infty\}$, the RW stays in $\{1, 2, 3, \dots\}$
- ▷ Replace $\mathbb{P}[\cdot]$ by $\mathbb{P}[\cdot | \{\tau = \infty\}]$ to obtain a *conditioned RW*
- ▷ Important question: do we have $\mathbb{P}[\cdot | \{\tau = \infty\}] = \mathbb{P}^f$?
 - 📎 Denisov & Wachtel '15 (general cones with zero drift); Courtiel, Melczer, Mishna & R. '16 (Gouyou-Beauchamps model with drift)
- ▷ Difficulty: exit time τ

Example (1/3) in the quadrant: the simple walk




- ▷ Uniform weights $\frac{1}{4}$
- ▷ $f(i, j) = i \cdot j$
- ▷ *Unique preharmonic function* (up to multiplicative factors)
- ▷ Product form 📎 Picardello & Woess '92

A second way for the conditioning (with the first exit time)

- ▷ RW $\{S(n)\}_{n \geq 0}$; *first exit time* $\tau = \inf\{n \geq 0 : S(n) = 0\}$
- ▷ On the event $\{\tau = \infty\}$, the RW stays in $\{1, 2, 3, \dots\}$
- ▷ Replace $\mathbb{P}[\cdot]$ by $\mathbb{P}[\cdot | \{\tau = \infty\}]$ to obtain a *conditioned RW*
- ▷ Important question: do we have $\mathbb{P}[\cdot | \{\tau = \infty\}] = \mathbb{P}^f$?
 - 📎 Denisov & Wachtel '15 (general cones with zero drift); Courtiel, Melczer, Mishna & R. '16 (Gouyou-Beauchamps model with drift)
- ▷ Difficulty: exit time τ

Example (2/3) in the quadrant: the Tandem walk




- ▷ Uniform weights $\frac{1}{3}$
- ▷ $f(i, j) = i \cdot j \cdot (i + j)$
- ▷ *Unique preharmonic function* (up to multiplicative factors) 📎 Biane '92

A second way for the conditioning (with the first exit time)

- ▷ RW $\{S(n)\}_{n \geq 0}$; *first exit time* $\tau = \inf\{n \geq 0 : S(n) = 0\}$
- ▷ On the event $\{\tau = \infty\}$, the RW stays in $\{1, 2, 3, \dots\}$
- ▷ Replace $\mathbb{P}[\cdot]$ by $\mathbb{P}[\cdot | \{\tau = \infty\}]$ to obtain a *conditioned RW*
- ▷ Important question: do we have $\mathbb{P}[\cdot | \{\tau = \infty\}] = \mathbb{P}^f$?
 - 📎 Denisov & Wachtel '15 (general cones with zero drift); Courtiel, Melczer, Mishna & R. '16 (Gouyou-Beauchamps model with drift)
- ▷ Difficulty: exit time τ

Example (3/3) in the quadrant: the GB walk



- ▷ Uniform weights $\frac{1}{4}$
- ▷ $f(i, j) = i \cdot j \cdot (i + j) \cdot (i + 2j)$
- ▷ *Unique preharmonic function* (up to multiplicative factors) 📎 Biane '92

Martin boundary theory

Rough description

▷ *Martin boundary* $\partial \rightsquigarrow$ set of all harmonic functions

Martin boundary theory

Rough description

- ▷ *Martin boundary* $\partial \rightsquigarrow$ set of all harmonic functions
- ▷ *Minimal Martin boundary* $\partial_m \rightsquigarrow$ integral representation of all harmonic functions: $f = \int_{\partial_m} \{\text{Martin kernel}\} d\mu_f$

 Martin '41;

Hunt '57; Doob '59; Choquet & Deny '60; Ney & Spitzer '66; Picardello & Woess '92

Martin boundary theory

Rough description

- ▷ *Martin boundary* $\partial \rightsquigarrow$ set of all harmonic functions
- ▷ *Minimal Martin boundary* $\partial_m \rightsquigarrow$ integral representation of all harmonic functions: $f = \int_{\partial_m} \{\text{Martin kernel}\} d\mu_f$

 Martin '41;

Hunt '57; Doob '59; Choquet & Deny '60; Ney & Spitzer '66; Picardello & Woess '92

Homogeneous random processes

- ▷ Well understood  Spitzer '64; Ney & Spitzer '66

Martin boundary theory

Rough description

- ▷ *Martin boundary* $\partial \rightsquigarrow$ set of all harmonic functions
- ▷ *Minimal Martin boundary* $\partial_m \rightsquigarrow$ integral representation of all harmonic functions: $f = \int_{\partial_m} \{\text{Martin kernel}\} d\mu_f$



 Martin '41;

Hunt '57; Doob '59; Choquet & Deny '60; Ney & Spitzer '66; Picardello & Woess '92

Homogeneous random processes

- ▷ Well understood  Spitzer '64; Ney & Spitzer '66

Non-homogeneous random processes: difficult problem

- ▷ Walks related to Lie algebras  Biane '90–'92
- ▷ Quadrant walks with **drift** $\sum_{y \in \mathbb{N}} y \cdot p(y)$  Ignatiouk-Robert '10
- ▷ Quadrant walks with **zero drift**

Partial results by  R. '14; Bouaziz, Mustapha & Sifi '15

Martin boundary theory

Rough description

- ▶ *Martin boundary* $\partial \rightsquigarrow$ set of all harmonic functions
- ▶ *Minimal Martin boundary* $\partial_m \rightsquigarrow$ integral representation of all harmonic functions: $f = \int_{\partial_m} \{\text{Martin kernel}\} d\mu_f$



 Martin '41;

Hunt '57; Doob '59; Choquet & Deny '60; Ney & Spitzer '66; Picardello & Woess '92

Homogeneous random processes

- ▶ Well understood  Spitzer '64; Ney & Spitzer '66

Non-homogeneous random processes: difficult problem

- ▶ Walks related to Lie algebras  Biane '90–'92
- ▶ Quadrant walks with **drift** $\sum_{y \in \mathbb{N}} y \cdot p(y)$  Ignatiouk-Robert '10
- ▶ Quadrant walks with **zero drift**

Partial results by  R. '14; Bouaziz, Mustapha & Sifi '15

- ▶ General RW in cones: open problem (**conjecture**: uniqueness \iff **drift** = 0)

Introduction & motivations


Applications in probability theory

Applications in combinatorics

Discrete harmonic functions in the quadrant

Asymptotics of some numbers of walks

Asymptotic statements




▷ *Total number of walks* starting at (k, l) :

$$q(n; k, l; \mathbb{N}^2) = \#_{\mathbb{N}^2} \{ (k, l) \xrightarrow{n} \mathbb{N}^2 \}$$
$$\sim f_1(k, l) \cdot \rho_1^n \cdot n^{\alpha_1}$$

 Not proved yet!

Asymptotics of some numbers of walks

Asymptotic statements



- ▷ *Total number of walks* starting at (k, ℓ) :

$$q(n; k, \ell; \mathbb{N}^2) = \#\mathbb{N}^2\{(k, \ell) \xrightarrow{n} \mathbb{N}^2\}$$
$$\sim f_1(k, \ell) \cdot \rho_1^n \cdot n^{\alpha_1}$$

 Not proved yet!


- ▷ *Excursions* starting at (k, ℓ) :

$$q(n; k, \ell; i, j) = \#\mathbb{N}^2\{(k, \ell) \xrightarrow{n} (i, j)\}$$
$$\sim f_2(k, \ell) \cdot f'_2(i, j) \cdot \rho_2^n \cdot n^{\alpha_2}$$

 Denisov & Wachtel '15

Asymptotics of some numbers of walks

Asymptotic statements



- ▷ *Total number of walks* starting at (k, ℓ) :

$$q(n; k, \ell; \mathbb{N}^2) = \#\mathbb{N}^2\{(k, \ell) \xrightarrow{n} \mathbb{N}^2\} \\ \sim f_1(k, \ell) \cdot \rho_1^n \cdot n^{\alpha_1}$$

Not proved yet!

- ▷ *Excursions* starting at (k, ℓ) :

$$q(n; k, \ell; i, j) = \#\mathbb{N}^2\{(k, \ell) \xrightarrow{n} (i, j)\} \\ \sim f_2(k, \ell) \cdot f'_2(i, j) \cdot \rho_2^n \cdot n^{\alpha_2}$$


Denisov & Wachtel '15

Preharmonicity of the prefactors

- ▷ f_1 is ρ_1 -harmonic & f_2 is ρ_2 -harmonic

Asymptotics of some numbers of walks

Asymptotic statements



- ▷ *Total number of walks* starting at (k, ℓ) :

$$q(n; k, \ell; \mathbb{N}^2) = \#\mathbb{N}^2\{(k, \ell) \xrightarrow{n} \mathbb{N}^2\} \\ \sim f_1(k, \ell) \cdot \rho_1^n \cdot n^{\alpha_1}$$

✎ Not proved yet!

- ▷ *Excursions* starting at (k, ℓ) :

$$q(n; k, \ell; i, j) = \#\mathbb{N}^2\{(k, \ell) \xrightarrow{n} (i, j)\} \\ \sim f_2(k, \ell) \cdot f'_2(i, j) \cdot \rho_2^n \cdot n^{\alpha_2}$$


✎ Denisov & Wachtel '15

Preharmonicity of the prefactors

- ▷ f_1 is ρ_1 -harmonic & f_2 is ρ_2 -harmonic: replace $q(n; k, \ell; \mathbb{N}^2)$ by its asymptotic expansion in the step-by-step construction
- $$q(n+1; k, \ell; \mathbb{N}^2) = \sum_{(i,j) \in \mathcal{S}} q(n; k-i, \ell-j; \mathbb{N}^2)$$

Asymptotics of some numbers of walks

Asymptotic statements



- ▷ *Total number of walks* starting at (k, ℓ) :

$$q(n; k, \ell; \mathbb{N}^2) = \#\mathbb{N}^2\{(k, \ell) \xrightarrow{n} \mathbb{N}^2\} \\ \sim f_1(k, \ell) \cdot \rho_1^n \cdot n^{\alpha_1}$$

 Not proved yet!

- ▷ *Excursions* starting at (k, ℓ) :

$$q(n; k, \ell; i, j) = \#\mathbb{N}^2\{(k, \ell) \xrightarrow{n} (i, j)\} \\ \sim f_2(k, \ell) \cdot f'_2(i, j) \cdot \rho_2^n \cdot n^{\alpha_2}$$


 Denisov & Wachtel '15

Preharmonicity of the prefactors

- ▷ f_1 is ρ_1 -harmonic & f_2 is ρ_2 -harmonic: replace $q(n; k, \ell; \mathbb{N}^2)$ by its asymptotic expansion in the step-by-step construction
- $$q(n+1; k, \ell; \mathbb{N}^2) = \sum_{(i,j) \in \mathcal{S}} q(n; k-i, \ell-j; \mathbb{N}^2)$$
- ▷ f'_2 is ρ_2 -harmonic for the *reversed step set* $\mathcal{S}' = -\mathcal{S}$

Asymptotics of some numbers of walks

Asymptotic statements



- ▷ *Total number of walks* starting at (k, l) :

$$q(n; k, l; \mathbb{N}^2) = \#_{\mathbb{N}^2} \{ (k, l) \xrightarrow{n} \mathbb{N}^2 \} \\ \sim f_1(k, l) \cdot \rho_1^n \cdot n^{\alpha_1}$$

 Not proved yet!

- ▷ *Excursions* starting at (k, l) :

$$q(n; k, l; i, j) = \#_{\mathbb{N}^2} \{ (k, l) \xrightarrow{n} (i, j) \} \\ \sim f_2(k, l) \cdot f'_2(i, j) \cdot \rho_2^n \cdot n^{\alpha_2}$$


 Denisov & Wachtel '15

Preharmonicity of the prefactors

- ▷ f_1 is ρ_1 -harmonic & f_2 is ρ_2 -harmonic: replace $q(n; k, l; \mathbb{N}^2)$ by its asymptotic expansion in the step-by-step construction $q(n+1; k, l; \mathbb{N}^2) = \sum_{(i,j) \in \mathcal{S}} q(n; k-i, l-j; \mathbb{N}^2)$
- ▷ f'_2 is ρ_2 -harmonic for the *reversed step set* $\mathcal{S}' = -\mathcal{S}$
- ▷ Drift zero: unique harmonic function $\implies f_1, f_2$ and f'_2

Random generation


Aim: generate efficiently a long walk (e.g., confined to a region)



A walk of length 18000



Random generation

Aim: generate efficiently a long walk (e.g., confined to a region)




A walk of length 18000

Different methods

- ▷ *Recursive method* (step-by-step construction)
- ▷ *Bijections* (if existing) For Kreweras see  Bernardi '07
- ▷ *Rejection algorithms*
 Bacher & Sportiello '16; Lumbroso, Mishna & Ponty '16




Random generation

Aim: generate efficiently a long walk (e.g., confined to a region)





A walk of length 18000

Different methods

- ▷ *Recursive method* (step-by-step construction)
- ▷ *Bijections* (if existing) For Kreweras see  Bernardi '07
- ▷ *Rejection algorithms*
 Bacher & Sportiello '16; Lumbroso, Mishna & Ponty '16
- ▷ Preharmonic functions and *Doob transform*  Fusy '16
(Difficulty: after Doob transform, non-uniform walks)



Potential theoretic tools

Counting numbers are caloric functions

- ▶ Asymptotics of numbers of quadrant walks (also with inhomogeneities)  D'Arco, Lacivita & Mustapha '16
- ▶ Asymptotics in three quarter of plane  Mustapha '16

Potential theoretic tools

Counting numbers are caloric functions



- ▷ Asymptotics of numbers of quadrant walks (also with inhomogeneities)  D'Arco, Lacivita & Mustapha '16
- ▷ Asymptotics in three quarter of plane  Mustapha '16

Zero drift case: classical inequalities Varopoulos '99–'09

General principle: there is a canonical function (*the réduite of the cone* f_c : $\Delta[f_c] = 0$) containing “all” the information

Potential theoretic tools

Counting numbers are caloric functions

- ▷ Asymptotics of numbers of quadrant walks (also with inhomogeneities)  D'Arco, Lacivita & Mustapha '16
- ▷ Asymptotics in three quarter of plane  Mustapha '16



Zero drift case: classical inequalities Varopoulos '99–'09

General principle: there is a canonical function (*the réduite of the cone* f_c : $\Delta[f_c] = 0$) containing “all” the information:

- ▷ $q(n; k, \ell; \mathbb{N}^2) \approx f(k, \ell) \cdot \rho^n \cdot n^\alpha$ as $n \rightarrow \infty$

Potential theoretic tools

Counting numbers are caloric functions

- ▷ Asymptotics of numbers of quadrant walks (also with inhomogeneities)  D'Arco, Lacivita & Mustapha '16
- ▷ Asymptotics in three quarter of plane  Mustapha '16



Zero drift case: classical inequalities Varopoulos '99–'09

General principle: there is a canonical function (*the réduite of the cone* f_c : $\Delta[f_c] = 0$) containing “all” the information:

- ▷ $q(n; k, \ell; \mathbb{N}^2) \approx f(k, \ell) \cdot \rho^n \cdot n^\alpha$ as $n \rightarrow \infty$
- ▷ $\alpha =$ homogeneity degree of f_c

Potential theoretic tools

Counting numbers are caloric functions

- ▷ Asymptotics of numbers of quadrant walks (also with inhomogeneities)  D'Arco, Lacivita & Mustapha '16
- ▷ Asymptotics in three quarter of plane  Mustapha '16



Zero drift case: classical inequalities Varopoulos '99–'09

General principle: there is a canonical function (*the réduite of the cone* f_c : $\Delta[f_c] = 0$) containing “all” the information:

- ▷ $q(n; k, \ell; \mathbb{N}^2) \approx f(k, \ell) \cdot \rho^n \cdot n^\alpha$ as $n \rightarrow \infty$
- ▷ $\alpha =$ homogeneity degree of f_c
- ▷ $f \sim f_c$ asymptotically

Potential theoretic tools

Counting numbers are caloric functions


- ▷ Asymptotics of numbers of quadrant walks (also with inhomogeneities)  D'Arco, Lacivita & Mustapha '16
- ▷ Asymptotics in three quarter of plane  Mustapha '16

Zero drift case: classical inequalities Varopoulos '99–'09

General principle: there is a canonical function (*the réduite of the cone* f_c : $\Delta[f_c] = 0$) containing “all” the information:

- ▷ $q(n; k, \ell; \mathbb{N}^2) \approx f(k, \ell) \cdot \rho^n \cdot n^\alpha$ as $n \rightarrow \infty$
- ▷ $\alpha =$ homogeneity degree of f_c
- ▷ $f \sim f_c$ asymptotically

Non-zero drift case: Cramér's transform & ongoing work

- ▷ Works if drift with ≤ 0 coordinates
- ▷ Ongoing work in the remaining cases  Garbit, Mustapha & R.

Introduction & motivations


Applications in probability theory

Applications in combinatorics

Discrete harmonic functions in the quadrant

Functional equation & Tutte's invariants

A functional equation reminiscent of the enumeration




- ▷ $F(x, y) = \sum_{i, j \geq 1} f(i, j) x^{i-1} y^{j-1}$
- ▷ $K'(x, y) = xy \{ \sum_{-1 \leq k, \ell \leq 1} p(k, \ell) x^{-k} y^{-\ell} - 1 \}$
- ▷ *Kernel functional equation:*

$$K'(x, y)F(x, y) = K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0)$$

Functional equation & Tutte's invariants

A functional equation reminiscent of the enumeration



- ▷ $F(x, y) = \sum_{i, j \geq 1} f(i, j) x^{i-1} y^{j-1}$
- ▷ $K'(x, y) = xy \{ \sum_{-1 \leq k, \ell \leq 1} p(k, \ell) x^{-k} y^{-\ell} - 1 \}$
- ▷ *Kernel functional equation:*


$$\begin{aligned} K'(x, y)F(x, y) = \\ K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0) \end{aligned}$$

Definition of Tutte's invariants

- ▷ Introduced to count q -colored triangulations & planar maps
📖 Tutte '73; Bernardi & Bousquet-Mélou '11
- ▷ Define X_0 & X_1 by $K'(X_0, y) = K'(X_1, y) = 0$
- ▷ Tutte's invariant: function $I \in \mathbb{Q}[[x]]$ such that $I(X_0) = I(X_1)$

Functional equation & Tutte's invariants

A functional equation reminiscent of the enumeration



- ▷ $F(x, y) = \sum_{i, j \geq 1} f(i, j) x^{i-1} y^{j-1}$
- ▷ $K'(x, y) = xy \left\{ \sum_{-1 \leq k, \ell \leq 1} p(k, \ell) x^{-k} y^{-\ell} - 1 \right\}$
- ▷ *Kernel functional equation:*

$$K'(x, y)F(x, y) = K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0)$$

Definition of Tutte's invariants


- ▷ Introduced to count q -colored triangulations & planar maps
 📖 Tutte '73; Bernardi & Bousquet-Mélou '11
- ▷ Define X_0 & X_1 by $K'(X_0, y) = K'(X_1, y) = 0$
- ▷ Tutte's invariant: function $I \in \mathbb{Q}[[x]]$ such that $I(X_0) = I(X_1)$

The sections $K'(x, 0)F(x, 0)$ & $K'(0, y)F(0, y)$ are invariants

- ▷ Evaluate the functional equation at X_0 & X_1
- ▷ Make the difference of the two identities

Functional equation & Tutte's invariants

A functional equation reminiscent of the enumeration



- ▷ $F(x, y) = \sum_{i, j \geq 1} f(i, j) x^{i-1} y^{j-1}$
- ▷ $K'(x, y) = xy \left\{ \sum_{-1 \leq k, \ell \leq 1} p(k, \ell) x^{-k} y^{-\ell} - 1 \right\}$
- ▷ *Kernel functional equation:*

$K'(x, y)F(x, y) = K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0)$

Definition of Tutte's invariants

- ▷ Introduced to count q -colored triangulations & planar maps
📖 Tutte '73; Bernardi & Bousquet-Mélou '11
- ▷ Define X_0 & X_1 by $K'(X_0, y) = K'(X_1, y) = 0$
- ▷ Tutte's invariant: function $I \in \mathbb{Q}[[x]]$ such that $I(X_0) = I(X_1)$

The sections $K'(x, 0)F(x, 0)$ & $K'(0, y)F(0, y)$ are invariants

- ▷ Evaluate the functional equation at X_0 & X_1
- ▷ Make the difference of the two identities

Does this characterize the sections?

Example: the SRW

A product-form generating function

$$f(i, j) = i \cdot j \implies F(x, y) = \sum_{i, j \geq 1} i \cdot j \cdot x^{i-1} y^{j-1} = \frac{1}{(1-x)^2(1-y)^2}$$

$$\text{Kernel: } K'(x, y) = xy \left\{ \frac{x}{4} + \frac{1}{4x} + \frac{y}{4} + \frac{1}{4y} - 1 \right\} = \frac{y(x-1)^2}{4} + \frac{x(y-1)^2}{4}$$

Example: the SRW

A product-form generating function

$$f(i, j) = i \cdot j \implies F(x, y) = \sum_{i, j \geq 1} i \cdot j \cdot x^{i-1} y^{j-1} = \frac{1}{(1-x)^2(1-y)^2}$$

$$\text{Kernel: } K'(x, y) = xy \left\{ \frac{x}{4} + \frac{1}{4x} + \frac{y}{4} + \frac{1}{4y} - 1 \right\} = \frac{y(x-1)^2}{4} + \frac{x(y-1)^2}{4}$$

Verification of the functional equation

$$K'(x, y)F(x, y) = K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0)$$

Example: the SRW

A product-form generating function

$$f(i, j) = i \cdot j \implies F(x, y) = \sum_{i, j \geq 1} i \cdot j \cdot x^{i-1} y^{j-1} = \frac{1}{(1-x)^2(1-y)^2}$$

$$\text{Kernel: } K'(x, y) = xy \left\{ \frac{x}{4} + \frac{1}{4x} + \frac{y}{4} + \frac{1}{4y} - 1 \right\} = \frac{y(x-1)^2}{4} + \frac{x(y-1)^2}{4}$$

Verification of the functional equation

$$\begin{aligned} K'(x, y)F(x, y) &= K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0) \\ &= \frac{x}{4} \times \frac{1}{(1-x)^2} + \frac{y}{4} \times \frac{1}{(1-y)^2} - 0 \times 1 \end{aligned}$$

Example: the SRW

A product-form generating function

$$f(i, j) = i \cdot j \implies F(x, y) = \sum_{i, j \geq 1} i \cdot j \cdot x^{i-1} y^{j-1} = \frac{1}{(1-x)^2(1-y)^2}$$

$$\text{Kernel: } K'(x, y) = xy \left\{ \frac{x}{4} + \frac{1}{4x} + \frac{y}{4} + \frac{1}{4y} - 1 \right\} = \frac{y(x-1)^2}{4} + \frac{x(y-1)^2}{4}$$

Verification of the functional equation

$$\begin{aligned} K'(x, y)F(x, y) &= K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0) \\ &= \frac{x}{4} \times \frac{1}{(1-x)^2} + \frac{y}{4} \times \frac{1}{(1-y)^2} - 0 \times 1 \end{aligned}$$

Tutte's invariants

$$\triangleright I(X_0) = I(X_1) \xrightarrow{X_0 X_1 = 1} I(x) = I\left(\frac{1}{x}\right) \implies I \text{ function of } x + \frac{1}{x}$$

$$\triangleright \boxed{K'(x, 0)F(x, 0) = \frac{x}{4} \frac{1}{(1-x)^2} = \frac{1}{4} \frac{1}{x + \frac{1}{x} - 2}} \text{ is an invariant}$$

Example: the SRW

A product-form generating function

$$f(i, j) = i \cdot j \implies F(x, y) = \sum_{i, j \geq 1} i \cdot j \cdot x^{i-1} y^{j-1} = \frac{1}{(1-x)^2(1-y)^2}$$

$$\text{Kernel: } K'(x, y) = xy \left\{ \frac{x}{4} + \frac{1}{4x} + \frac{y}{4} + \frac{1}{4y} - 1 \right\} = \frac{y(x-1)^2}{4} + \frac{x(y-1)^2}{4}$$

Verification of the functional equation

$$\begin{aligned} K'(x, y)F(x, y) &= K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0) \\ &= \frac{x}{4} \times \frac{1}{(1-x)^2} + \frac{y}{4} \times \frac{1}{(1-y)^2} - 0 \times 1 \end{aligned}$$

Tutte's invariants

- ▷ $I(X_0) = I(X_1) \xrightarrow{X_0 X_1 = 1} I(x) = I\left(\frac{1}{x}\right) \implies I$ function of $x + \frac{1}{x}$
- ▷ $K'(x, 0)F(x, 0) = \frac{x}{4} \frac{1}{(1-x)^2} = \frac{1}{4} \frac{1}{x + \frac{1}{x} - 2}$ is an invariant

Why *this* function of $x + \frac{1}{x}$?


- ▷ Of order 1 in $x + \frac{1}{x} \rightsquigarrow$ *Minimality* (conformal mappings)
- ▷ $F(1, 0) = \infty \rightsquigarrow$ *Liouville's theorem*

Tutte's invariants & conformal mappings

A general theorem

$K'(x, 0)F(x, 0) = w(x)$, *characterized by*

- ▷ Conformal mapping of a certain domain
- ▷ $w(x) = w(\bar{x})$
- ▷ $w(1) = \infty$
- ▷ Same for $K'(0, y)F(0, y)$




Tutte's invariants & conformal mappings

A general theorem

$K'(x, 0)F(x, 0) = w(x)$, *characterized by*


- ▶ Conformal mapping of a certain domain
- ▶ $w(x) = w(\bar{x})$
- ▶ $w(1) = \infty$
- ▶ Same for $K'(0, y)F(0, y)$



Going back to the SRW

$K'(x, 0)F(x, 0) = \frac{x}{4(1-x)^2}$, *characterized by*

- ▶ Conformal mapping of the unit disc
- ▶ $w(e^{i\theta}) = w(e^{-i\theta})$
- ▶ $w(1) = \infty$
- ▶ Same for $K'(0, y)F(0, y) = \frac{y}{4(1-y)^2}$




Tutte's invariants & conformal mappings

A general theorem

$K'(x, 0)F(x, 0) = w(x)$, *characterized by*


- ▷ Conformal mapping of a certain domain
- ▷ $w(x) = w(\bar{x})$
- ▷ $w(1) = \infty$
- ▷ Same for $K'(0, y)F(0, y)$



Going back to the SRW

$K'(x, 0)F(x, 0) = \frac{x}{4(1-x)^2}$, *characterized by*

- ▷ Conformal mapping of the unit disc
- ▷ $w(e^{i\theta}) = w(e^{-i\theta})$
- ▷ $w(1) = \infty$
- ▷ Same for $K'(0, y)F(0, y) = \frac{y}{4(1-y)^2}$



Question

How deep is this *connection conformal maps/harmonic functions*?

