

MPRI 2-38-1: Algorithms and combinatorics for geometric graphs

Lecture 1

Preliminaries on planar graphs

september 18, 2024

Luca Castelli Aleardi




Part 0

Introduction and historical background

(a short digression on planar graphs and their applications)


Origins of Graph Theory (back to Euler)

Solutio problematis ad geometriam situs pertinentis
(1735, presented to the St. Petersburg Academy)



(Königsberg, Prussia)

(from Wikipedia)



Eulerian path: it visits every edge exactly once

Theorem (Euler 1735, Hierholzer 1873)


A graph G contains an *Eulerian walk (path)* if and only if G is connected and the number of vertices of odd degree is 0 or 2.

Theorem


A connected graph contains an *Eulerian circuit* if and only if there are no vertices of odd degree.

Planar graphs

Design of integrated circuits (VLSI)




french roads network




Planar graphs in computational geometry and geometric modeling

(Delaunay triangulations, Voronoi diagrams, 3D meshes, ...)




Voronoi diagram
Delaunay triangulation

3D Geometric modeling
Pixar characters (image by De Goes et al.)




GIS Technology



Terrain modelling

triangulations were already
used in 18th century: approxi-
mation of the meridian
(Delambre et Méchain, 1792)



3D paper sculpture (*DT Workshop*)




Major results (on planar graphs) in graph theory

Kuratowski theorem (1930)




(cfr Wagner's theorem, 1937)

- G contains neither K_5 nor $K_{3,3}$ as minors




Thm (Steinitz, 1916)

3-connected planar graphs are skeletons of convex polyhedra




Thm (Koebe-Andreev-Thurston)

Every planar graph with n vertices is isomorphic to the intersection graph of n disks in the plane.



Thm (Schnyder '90)




(dimension of partial orders)

- G planar iff $\dim(G) \leq 3$

Efficient algorithms on planar graphs

Graph drawing

Planarity testing



	Tutte barycentric layout	Schnyder layout	FFP layout
fish model ($n = 241$)			
random triang. ($n = 100$)			

Minimum spanning tree




Planar Separators

Minimum cut

Part I

What is a planar graph?

(some terminology: embedded graphs, topological and combinatorial maps)



Graphs

A graph $G = (V, E)$ is a pair of:



- a set of *vertices* $V = (v_1, \dots, v_n)$
- a collection of $E = (e_1, \dots, e_m)$ elements of the cartesian product $V \times V = \{(u, v) \mid u \in V, v \in V\}$ (called *edges*).

circuit: a closed walk without repeated vertices

$$A_G[i, j] = \begin{cases} 1 & v_i \text{ adjacent } v_j \\ 0 & \text{otherwise} \end{cases}$$

$$A_G = \begin{bmatrix} 0 & 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 1 & 1 \\ 1 & 1 & 0 & 1 & 1 \\ 1 & 1 & 1 & 0 & 1 \\ 0 & 1 & 1 & 1 & 0 \end{bmatrix}$$


$$D_G = \begin{matrix} & e_1 & e_2 & e_3 & e_4 & e_5 & e_6 & e_7 & e_8 & e_9 \\ \begin{bmatrix} 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ \dots & \dots & & & & & & & \\ \dots & & & & & & & & \end{bmatrix} & v_1 \\ & v_2 \end{matrix}$$



Planar drawings: some basic notion of topology

topological space: a set X with a collection of *open sets* (subsets of X) satisfying:



- X itself and the empty set are open
- the union of open sets is open
- any finite intersection of open sets is open



$f : X \rightarrow Y$ is *continuous*: the inverse image of an open set of Y is open

$f : X \rightarrow Y$ is *homeomorphism*: f, f^{-1} are bijective and continuous

a *path* is a continuous map $p : [0, 1] \rightarrow X$
(the path is *simple* if p is 1-to-1)



Remark:
we consider topological spaces which are *Haussdorff* (any two distinct points have disjoint neighborhoods)

Planar drawings of planar graphs

an *embedding* of G into R^2 is a 1-to-1 continuous map satisfying:


- (i) graph vertices are represented as points ;
- (ii) edges are represented as paths (curves);
- (iii) the images of vertices are distinct points
- (iv) the images of edges simple (no self-intersections at the interior)
- (v) the interior of the images of edges are disjoint (no crossings)
- (vi) edges cannot pass through a vertex (except at its extremities)

faces of a graph embedding: connected component of the image of the vertices/edges of G

cellular embedding: the faces are homeomorphic to open disks

planar graph: a graph admitting an embedding in the plane

plane graph: a planar graph + a cellular embedding




two cellular embeddings defining the same planar graph

Planar drawings of planar graphs


an *embedding* of G into R^2 is a 1-to-1 continuous map satisfying:

- (i) graph vertices are represented as points ;
- (ii) edges are represented as paths (curves);
- (iii) the images of vertices are distinct points
- (iv) the images of edges simple (no self-intersections at the interior)
- (v) the interior of the images of edges are disjoint (no crossings)
- (vi) edges cannot pass through a vertex (except at its extremities)



plane graph: a planar graph + a cellular embedding


(*topological*) *map*: cellular embedding up to homeomorphism (equivalence class)



The Jordan curve theorem

Theorem

Any simple closed curve C in the plane partitions $\mathbb{R}^2$ into two disjoint arcwise-connected open sets.



($Ext(C)$ and $Int(C)$ are closed sets)


$$Ext(C) \cap Int(C) = C$$

Remark:

Any arc joining a point p in the (open) interior to a point q in the (open) exterior must meet C at least once.

Jordan curve Theorem (reformulation)

Let G a graph embedded on $\mathbb{S}^2$. Then G disconnects $\mathbb{S}^2$ if and only if it contains a circuit





The Jordan curve theorem: application

Theorem

Any simple closed curve C in the plane partitions $\mathbb{R}^2$ into two disjoint arcwise-connected open sets.

Theorem

The graph K_5 is not planar



Proof (topological)

(by contradiction) Let G be a planar embedding of K_5

K_5 is complete $\longrightarrow$ it contains $C := \{v_1, v_2, v_3, v_1\}$ (simple cycle)

G planar $\longrightarrow f(C)$ simple closed curve (separating the plane)


w.l.o.g. assume $v_4 \in \text{Int}(C)$

(G planar, no edge crossings)

$(v_4, v_1) \in \text{Int}(C)$

$(v_4, v_2) \in \text{Int}(C)$

$(v_4, v_3) \in \text{Int}(C)$





The Jordan curve theorem: application

Theorem

Any simple closed curve C in the plane partitions $\mathbb{R}^2$ into two disjoint arcwise-connected open sets.

Theorem

The graph K_5 is not planar




Proof (topological)

Consider 3 cycles

$$C_1 := \{v_2, v_3, v_4, v_2\}$$

$$C_2 := \{v_3, v_1, v_4, v_3\} \longrightarrow v_i \in \text{ext}(C_i) \ (i \in \{1, 2, 3\})$$

$$C_3 := \{v_1, v_2, v_4, v_1\}$$



K_5 is complete $\longrightarrow$ it contains $(v_5, v_1), (v_5, v_2), (v_5, v_3) \longrightarrow v_5 \in \text{ext}(C_i) \ (i \in \{1, 2, 3\})$
(Jordan curve theorem)

$v_5 \in \text{ext}(C) \xrightarrow{\text{(Jordan curve theorem)}} (v_5, v_4) \text{ meets } C \text{ somewhere}$



(edge crossing, contradicting the planarity of G)



Planar graphs and graphs embeddable on the sphere are the same

Theorem

A graph G is embeddable on the sphere $\mathbb{S}^2$ if and only if it is embeddable on the plane






Planar graphs and graphs embeddable on the sphere are the same

Theorem

A graph G is embeddable on the sphere $\mathbb{S}^2$ if and only if it is embeddable on the plane

Proof



Stereographic projection $\Pi : \mathbb{S}^2 \setminus N \longrightarrow \mathbb{R}^2$
(homeomorphism: Π and its inverse are bijective and continuous)

$$\Pi^{-1}(x, y) \begin{pmatrix} 2x/\chi \\ 2y/\chi \\ 1 - 2/\chi \end{pmatrix}$$

$$\chi := x^2 + y^2 + 1$$

Remark

To get a planar embedding of a graph G , just take a point N in the interior of a face of G on $\mathbb{S}^2$, and project on $\mathbb{R}^2$

$$\Pi(x, y, z) \begin{pmatrix} \frac{x}{1-z} \\ \frac{y}{1-z} \end{pmatrix}$$

Combinatorial maps: representations and data structures



Cellularly embedded planar graphs as *combinatorial maps*

Let G a cellular graph embedding

The *combinatorial map* associated to G is the set of closed walks, obtained walking around the boundary of each face (in our example in cw direction)

2 permutations on the set H of the $2m$ darts

- (i) α involution without fixed point;
- (ii) ϕ gives the cyclic ordering of the darts (edges) around each face




$$\phi = (1, 2, 3, 4)(17, 23, 18, 22)(5, 10, 8, 12) \dots$$

$$\alpha = (2, 18)(3, 5)(4, 7)(12, 13)(9, 15) \dots$$

2 permutations on the set H of the $2m$ darts

- (i) α involution without fixed point;
- (ii) σ gives the cyclic ordering of the darts (edges) around each vertex



$$\sigma = (1, 20, 18)(4, 5, 13)(3, 12, 7) \dots$$

$$\alpha = (2, 18)(3, 5)(4, 7)(12, 13)(9, 15) \dots$$

(*) $\alpha\sigma\phi = Id$; **The two representations are dual to each other**

(*) the group generated by σ , α et ϕ transitively on H .

$$3 \xrightarrow{\phi} 4 \xrightarrow{\sigma} 5 \xrightarrow{\alpha} 3$$



Cellularly embedded planar graphs as *combinatorial maps*

Let G a cellular graph embedding

The *combinatorial map* associated to G is the set of closed walks, obtained walking around the boundary of each face (in our example in cw direction)





2 permutations on the set H of the $2m$ darts

- (i) α involution without fixed point;
- (ii) ϕ gives the cyclic ordering of the darts (edges) around each face



2 permutations on the set H of the $2m$ darts

- (i) α involution without fixed point;
- (ii) σ gives the cyclic ordering of the darts (edges) around each vertex




Graph: adjacency lists representation

easy to implement

quite compact

not efficient for traversal



for each face (of degree d), store:

- d references to adjacent vertices

for each vertex, store:

- 1 reference to its coordinates

```
class Point{
    double x;
    double y;
}
```

geometric information

Memory cost

$$\sum_i \deg(v_i) = 2 \times e$$

Size (number of references)

```
class Vertex{
    Point p;
    List<Vertex> neigh
}
```

combinatorial information

Queries/Operations

List all vertices

Test adjacency between u and v

Find the 3 neighboring faces of f


List the neighbors of vertex v

vertex locations

v_0	6 5 1 2	(x_0, y_0, z_0)	(x_1, y_1, z_1)
v_1	0		
v_2	0		
v_3	4 5		
	5 6 3 6		
	0 4 3		
	4 0		
	8		
	7		

Half-edge data structure: polygonal (orientable) meshes

2 half-edges per edge



```
class Halfedge{
    Halfedge prev, next, opposite;
    Vertex v;
    Face f;
}
class Vertex{
    Halfedge e;
    Point p;
}
class Face{
    Halfedge e;
}
```

combinatorial information

```
class Point{
    double x;
    double y;
}
```

geometric information

Size (number of references)


$$f + 5 \times h + n \approx 2n + 5 \times (2e) + n$$

```
public int degree() {
    Halfedge<X> e, p;
    if(this.halfedge==null) return 0;

    e=halfedge; p=halfedge.next;
    int cont=1;
    while(p!=e) {
        cont++;
        p=p.next;
    }
    return cont;
}
```

Flag representation


4 flags per edge



```
class Flag{
  Flag ei, fi, vi;
  Vertex u;
}
```

```
class Vertex{
  Flag f;
  Point p;
}
```

combinatorial information



$$0 \leq f \leq 4 * e - 1$$

$$0 \leq v \leq n - 1$$

navigation around vertices

```
vertexDegree(Flag f) {
  int j=0;
  Flag g=f;
  do {
    ++j;
    g=g.ei().fi();
  } while (g!=f);
  return j;
}
```


navigation around faces

```
faceDegree(Flag f) {
  int j=0;
  Flag g=f;
  do {
    ++j;
    g=g.ei().vi();
  } while (g!=f);
  return j;
}
```

Duality

Definition

Given a cellular graph embedding G on the sphere, its *dual graph* G^* is a graph embedding for which: we put a (dual) vertex f^* in the interior of a face $f \in G$; and create a dual edge e^* crossing an edge $e \in G$




Remarks:

- The dual of a plane graph is connected (**exercise**)
- A dual graph embedding is also cellular
- The combinatorial map of the dual graph is uniquely defined
- $(G^*)^* \cong G$


Duality

Definition

Given a cellular graph embedding G on the sphere, its *dual graph* G^* is a graph embedding for which: we put a (dual) vertex f^* in the interior of a face $f \in G$; and create a dual edge e^* crossing an edge $e \in G$



(genus 0) triangle mesh



planar triangulation
(simple connected plane graph, with all faces of degree 3)

Remark:

- A simple connected plane graph is a planar triangulation if and only if its dual is a cubic graph

Duality

Exercise:


Given a plane graph G with m edges show that:

$$\sum_{f \in F} \text{degree}(f) = 2m$$

$$\sum_{v \in V} \text{degree}(v) = 2m$$


Duality: edge contractions and deletions

edge contraction and vertex split



Remark:



Edge contractions and edge deletions preserve some properties



Property


Let G be a connected cellularly embedded graph, and e a non cut edge. Then

$$(G \setminus e)^* \cong G^*/e^*$$



Remark:

What happens if e belongs to a separating triangle?



Duality



Lemma

Let us consider a graph embedding $G = (V, E)$ and its dual $G^* = (F^*, E^*)$, and a subset of edges $E' \subset E$. Then we have



- (V, E') is acyclic if and only if $(F^*, (E \setminus E')^*)$ is connected

Corollary:

(V, E') is a spanning tree if and only if $(F^*, (E \setminus E')^*)$ is a spanning tree.



(V, E') is acyclic



$(F^*, (E \setminus E')^*)$ is connected

Remove the (dual) blue edges which are crossing the (red) edges in E'

Duality


Lemma

Let us consider a graph embedding $G = (V, E)$ and its dual $G^*(F^*, E^*)$, and a subset of edges $E' \subset E$. Then we have


- (V, E') is acyclic if and only if $(F^*, (E \setminus E')^*)$ is connected.

Proof

(V, E') is acyclic $\Leftrightarrow (\mathbb{S}^2 \setminus E')$ is connected



$(F^*, (E \setminus E')^*)$




Claim: $(\mathbb{S}^2 \setminus E')$ is connected $\Leftrightarrow (F^*, (E \setminus E')^*)$ is connected

Consider points $x \in f$ and $x' \in f'$ (in the embedding G)

There is a path $P_{xx'}$ $\begin{cases} \text{avoiding the edges in } E' \text{ and} \\ P_{xx'} \text{ is lying only in } f \text{ and } f' \end{cases}$

$\Leftrightarrow f^*$ and f'^* are adjacent by some edge not in E'

$\Leftrightarrow f^*$ and f'^* are adjacent in $(F^*, (E \setminus E')^*)$



Part II


Euler formula and its consequences

Euler-Poincaré characteristic: topological invariant

$$\chi := n - e + f$$

One of the (11) world's most beautiful equations


(according to livescience.com)



planar map


$$n - e + f = 2$$

Euler's relation




Euler-Poincaré characteristic: topological invariant


$$\chi := n - e + f$$



planar map




(convex) polyhedron




$$n - e + f = 2$$

Euler's relation

$$\chi = 0$$



$$\chi = -4$$




$$n = 1660$$

$$e = 4992$$

$$f = 3328$$

$$g = 3$$

$$n - e + f = 2 - 2g$$



$$n = 364$$

$$e = 675$$

$$f = 302$$

$$b = 11$$

$$g = 0$$

$$n - e + f = 2 - b$$

Euler's relation: first proof (induction on the faces)

Theorem (Euler's relation)

Given a connected plane graph G we have:

$$v(G) - e(G) + f(G) = 2$$

Let us first prove a preliminary result

Lemma

If G is a tree then we have: $e(G) = v(G) - 1$

proof: (induction on the nodes)


$$v(G \setminus x) = v(G) - 1$$

$$e(G \setminus x) = e(G) - 1$$

base case of the induction

$$v(G) = 1$$

$$e(G) = 0$$



Claim

Any tree contains at least one leaf
(exercise)

Euler's relation: first proof (induction on the faces)

Theorem (Euler's relation)

Given a connected plane graph G we have:

$$v(G) - e(G) + f(G) = 2$$

Let us first prove a preliminary result

Lemma

If G is a tree then we have: $e(G) = v(G) - 1$

proof: (induction on the nodes)


$$v(G \setminus x) = v(G) - 1$$

$$e(G \setminus x) = e(G) - 1$$

base case of the induction

$$v(G) = 1$$

$$e(G) = 0$$



Claim

Any tree contains at least one leaf

Claim 2

If a graph G all vertices have degree at least 2, then G contains a cycle.

(exercise)

Euler's relation: first proof (induction on the faces)

Theorem (Euler's relation)

Given a connected plane graph G we have:

$$v(G) - e(G) + f(G) = 2$$

Let us first prove a preliminary result

Lemma

If G is a tree then we have: $e(G) = v(G) - 1$

proof: (induction on the nodes)


$$v(G \setminus x) = v(G) - 1$$

$$e(G \setminus x) = e(G) - 1$$

base case of the induction

$$v(G) = 1$$

$$e(G) = 0$$



(remove one leaf)


Claim

Any tree contains at least one leaf

Claim 2

If a graph G all vertices have degree at least 2, then G contains a cycle.

(solution) assume G is simple
(otherwise the statement is trivial)



let $P := v_0, v_1 \dots v_{k-1} v_k$ (path of maximal length in G)

$degree(v_k) \geq 2$

- $u \notin P$ $P \cup \{u\}$ is longer
- $u \in P$ $u = v_i$ ($i \leq k-2$)
it defines a cycle

Euler's relation: first proof (induction on the faces)


Theorem (Euler's relation)

Given a connected plane graph G we have:

$$v(G) - e(G) + f(G) = 2$$

proof: (induction on the faces)

(base case) $f(G) = 1$



G is a tree (use previous Lemma)

$$e(G) = v(G) - 1$$

(general case) $f(G) \geq 2$


There is a non cut edge $e \longrightarrow G \setminus e$ is connected $\xrightarrow{\text{(induction hypothesis)}} v(G \setminus e) - e(G \setminus e) + f(G \setminus e) = 2$
 $f(G \setminus e) = f(G) - 1$

$$f(G \setminus e) = f(G) - 1$$


$$e(G \setminus e) = e(G) - 1$$

$$v(G \setminus e) = v(G)$$

$$v(G) - e(G) + f(G) = 2$$



remove an arbitrary
non cut edge e



Euler's relation: by induction (variation)

Theorem (Euler's relation)

Given a connected plane graph G we have:

$$v(G) - e(G) + f(G) = 2$$

$$\chi := n - e + f$$

proof: (induction on the faces)

base case: $f = 1$



$$\chi(t) = 3 - 3 + 2 = 2$$

$$\chi(M) = \chi(M^t)$$

exercise: prove the following invariant
invariant: the boundary (exterior) is a simple cycle
perform the removal according to a shelling order



$$e''' = e'' - 2$$

$$f''' = f'' - 1$$


$$n''' = n'' - 1$$

$$e' = e - 1 \quad f' = f - 1$$


$$e'' = e' - 1 \quad f'' = f' - 1$$




remove a boundary edge



remove a boundary edge





remove a triangle



Euler's relation: second proof (via the dual)

$$n - e + f = 2$$








Pepakura software for unfolding polyhedral surfaces

Euler's relation for polyhedral surfaces

Overview of the proof

$$n - e + f = 2$$

planar graph G



f faces

patron


take any spanning tree T
 $n - 1$ edges

$f - 1$ edges f vertices

$$e = (n - 1) + (f - 1)$$


dual spanning tree T^* avoiding the edges of T

take the dual graph T^* avoiding the edges of T : this is a spanning tree of G^*




Exercise 1: give a proof of Euler formula

(using spherical geometry)




Geodesic: arc of great circle (between p and q)



Girard Theorem

$$\alpha + \beta + \gamma = \pi + \frac{\text{area}(A,B,C)}{r^2}$$

Hints:




Spherical drawing of an octahedron

- $\alpha + \beta + \gamma + \delta = 2\pi$

- $\sum_{i \leq f} \text{area}(P_i) = 4\pi r^2$

(since faces are non overlapping)

edges are drawn as non crossing geodesic arcs



Theorem (Girard)

Given a spherical polygon P_i of size k_i , the sum of internal angles α_{ij} satisfy

$$\sum_{j \leq k} \alpha_{ij} = (k_i - 2)\pi + \frac{\text{area}(P_i)}{r^2}$$

Exercise 2 (soccer ball theorem)

Given a plane graph, where every face is a pentagon or a hexagon, and such that every vertex has degree 3, show that there must be exactly twelve pentagonal faces.



Euler's relation: consequences

Corollary: linear dependence between edges, vertices and faces

$$f \leq 2n - 4$$

$$e \leq 3n - 6$$

proof (double counting argument)

$$f = f_1 + f_2 + f_3 + \dots$$

$$n = n_1 + n_2 + n_3 + \dots$$

all faces have degree at least 3 ($\mathcal{G}$ simple simple), then we get

$$f = f_3 + f_4 + \dots$$

every edge appears twice

$$2e = 3 \cdot f_3 + 4 \cdot f_4 + \dots$$

then we get

$$2e - 3f \geq 0$$

Euler's relation: consequences

Corollary: linear dependence between edges, vertices and faces

$$f \leq 2n - 4$$

$$e \leq 3n - 6$$

given $2e - 3f \geq 0$

by applying Euler formula, we obtain


$$3n - 6 = 3(e - f + 2) - 6 = 3e - 3f$$

$$3n - 6 = e + (2e - 3f) \geq e$$

$$3n - 6 \geq e$$

Euler's relation: consequences

assume $\mathcal{G}$ is a *simple* planar graph: no multiple edges, no loops



furthermore, assume all **faces have degree at least 3**, then we get

$$f = f_3 + f_4 + \dots$$

every edge appears twice

$$2e = 3 \cdot f_3 + 4 \cdot f_4 + \dots$$


then we get

$$2e - 3f \geq 0$$

by applying Euler formula, we obtain

$$3n - 6 = 3(e - f + 2) = 3e - 3f \geq 0$$

$$e \leq 3n - 6$$



furthermore, assume there are **no cycles of length 3**, then we get

$$f = f_4 + f_5 + \dots$$

every edge appears twice

$$2e = 4 \cdot f_4 + 5 \cdot f_5 + \dots$$

then we get

$$2e - 4f \geq 0$$


use again Euler formula

$$3n - 6 = 3(e - f + 2) = 3e - 3f \geq 0$$

$$e \leq 2n - 4$$

Euler's relation for polyhedral surfaces

can we construct a regular
(genus 0) mesh, where every
vertex has degree 6?



Euler's relation for polyhedral surfaces

we just showed $2e - 3f \geq 0$

proof (double counting argument)

Assume all the vertices have degrees ≥ 6 :

the total number of vertices is: $n = n_6 + n_7 + n_8 + \dots$

using a double counting of edges: $2e = 6 \cdot n_6 + 7 \cdot n_7 + 8 \cdot n_8 + \dots$




$$2e - 6 \cdot n \geq 0$$

$$\left. \begin{array}{l} 2e - 6 \cdot n \geq 0 \\ 2e - 3f \geq 0 \end{array} \right\} \longrightarrow 6(e - n - f) = (2e - 6n) + 2(2e - 3f) \geq 0$$

$$e - n - f \geq 0 \longrightarrow e \geq n + f$$

contradicting Euler formula: $e = n + f - 2$



Euler's relation and Kuratowski theorem (easy direction)

theorem (Kuratowski 1930)

G is planar iff it contains no subdivision of K_5 nor $K_{3,3}$



theorem (Wagner, 1937)

G is planar iff it does not contain K_5 nor $K_{3,3}$ as minors

Lemma

The graphs K_5 and $K_{3,3}$ are not planar

Exercise: give a combinatorial proof




$K_{3,3}$ bipartite:

Euler's relation and Kuratowski theorem (easy direction)

Lemma


The graphs K_5 and $K_{3,3}$ are not planar

Proof: (combinatorial)



$$e \leq 3n - 6 = 9$$

but we have $e(K_5) = \binom{5}{2} = 10$



$K_{3,3}$ bipartite:

no cycle of length 3: faces have degree ≥ 4

$$4f(G) \leq \sum_{f \in F} \deg(f) = 2e(G) = 18$$

so the number of faces is $f(G) \leq 4$

$$2 = v(G) - e(G) + f(G) \leq 6 - 9 + 4 = 1$$

Euler's relation and Kuratowski theorem (easy direction)


theorem (Kuratowski 1930)

G is planar iff it contains no subdivision of K_5 nor $K_{3,3}$

theorem (Wagner, 1937)


G is planar iff it does not contain K_5 nor $K_{3,3}$ as minors

A graph G' is a *minor* of a graph G it can be obtained from G with a sequence of vertex/edge deletions and edge contractions



K_5 is a minor of the Petersen graph

A graph G' is a *subdivision* of a graph G if it can be obtained from G with a sequence of edge subdivisions



Subdivisions of K_5 and $K_{3,3}$

Remark

Minors of planar graphs are planar

Remark

A graph G is planar if and only if every subdivision of G is planar

Part III



3-connectedness and planar graphs

3-connectedness

Defini


G is 3-connected if

$\left\{ \begin{array}{l} \text{is connected and} \\ \text{the removal of one or two vertices} \\ \text{does not disconnect } G \end{array} \right.$




at least 3 vertices are required to disconnect the graph

cut-vertex




cut-pair



Menger Theorem

if G is 3-connected then for every pair of vertices u and v there exist 3 vertex disjoint paths (intersecting only at u and v)




(see Lecture 5, for a simple proof in the triangulated planar case)



3-connected planar graphs: Whitney theorem

Thm (Whitney, 1933)

3-connected planar graphs admit an unique embedding (up to homeomorphism and inversion of the sphere $\mathbb{S}^2$).






Remark: why 3-connectedness is important?

3-connected planar graphs: Whitney theorem

Thm (Whitney, 1933)

3-connected planar graphs admit an unique embedding (up to homeomorphism and inversion of the sphere $\mathbb{S}^2$).



$f_i :=$ number of faces of degree i


$$f_3 = 4$$

$$f_4 = 2$$


$$f_5 = 1$$

$$f_6 = 0$$

$$f_7 = 1$$



$$|F| = 9$$



$$f_3 = 4$$

$$f_4 = 2$$

$$f_5 = 0$$

$$f_6 = 2$$

$$f_7 = 0$$

two different (non equivalent) embeddings of the same graph

Bridges: some terminology

G a connected graph

C a cycle

Bridges := subgraphs induced by the edges of $E(G) \setminus E(C)$

Remarks

bridges can only intersect at the vertices of C

trivial bridges do not have inner vertices: loops, chords

from any two vertices of a bridge there exists one path internally disjoint from C

if G is non-separable then there are two vertices of attachment

k -bridge is a bridge with k vertices of attachment

equivalent bridges: same point of attachment (B_1 and B_2 , which are 3-bridges)


B_2 and B_4 are said to *avoid each other*

B_3 and B_4 are said to be *overlapping*

Two bridges B and B' are *skew* if there exist 4 vertices of attachment u, v (on B) and u', v' on B' which are listed consecutively on C : u, u', v, v'

B_3 and B_4 are skew

B_1 and B_2 are not skew




Bridges of cycles: properties

Lemma 1


Given a cycle in a graph G the overlapping bridges are either skew or else equivalent 3-bridges.

proof:



Case 1 either B or B' is a chord (2-bridge)
they must be skew (as B_3 and B_4)

Case 2 both B and B' have at least 3 points of attachment
 B and B' are not equivalent
they must be skew (as B_1 and B_2)



Case 3a B and B' are equivalent
both B and B' are 3-bridges (as B_4 and B_5)

Case 3b B and B' are equivalent
both B and B' are k -bridges ($k \geq 4$)
they must be skew

Bridges (in planar graphs): properties

Lemma 2

Given a cycle C in a plane graph G the inner (outer) bridges avoid each other.



(they are not overlapping)

proof:

Case 1 B and B' are skew $\longrightarrow$ there are u, u', v, v' consecutive on C

take the two disjoint paths P and P' (included in B and B')

The graph $H = P \cup C \cup P'$ is planar (subgraph of G)



Define a graph K adding a vertex on the outer face (and connecting it to vertices on C)

The graph K should be planar by construction

But K is also a subdivision of K_5 (non planar)

(contradiction)



Case 2 B and B' are equivalent 3-bridges

exercise

3-connected planar graphs: Whitney theorem

Thm (Tutte, 1963)

A cycle in a 3-connected planar graph is a facial cycle (bounding a face) if and only if it is non-separating.



Def:

a cycle is *non-separating* if it has no chords and at most one non trivial bridge

3-connected planar graphs: Whitney theorem

Thm (Tutte, 1963)

A cycle in a 3-connected planar graph is a facial cycle (bounding a face) if and only if it is non-separating.

proof:


Assume C is not a facial cycle




There is at least one inner and one outer bridge (not loops)



C cannot be a non-separating cycle




both bridges are non trivial



bridge is a chord


Let C a facial cycle (assume it is bounding the outer face)

All bridges are in the interior of C and avoid each other (previous Lemma)



the chord defines a 2-cut $\{x, y\}$

contradicting 3-connectedness



there are at least two bridges (avoiding each other: their segments are internally disjoint)

the pair $\{x, y\}$ is a vertex cut of the inner vertices of the bridges

contradicting 3-connectedness


Part IV

Exercices

Exercise 1 (Euler formula via spherical geometry)

Give an alternative geometric proof of Euler Formula, using Girard theorem

$$n - e + f = 2$$




Girard Theorem

$$\alpha + \beta + \gamma = \pi + \frac{\text{area}(A,B,C)}{r^2}$$

Exercise 2 (soccer ball theorem)


Given a plane graph, where every face is a pentagon or a hexagon, and such that every vertex has degree 3, show that there must be exactly twelve pentagonal faces.




Using spherical geometry

An alternative geometric proof of Euler Formula

$$n - e + f = 2$$




Geodesic: arc of great circle (between p and q)




Girard Theorem

$$\alpha + \beta + \gamma = \pi + \frac{\text{area}(A,B,C)}{r^2}$$




Spherical drawing of an octahedron

$$\alpha + \beta + \gamma + \delta = 2\pi$$




$$\sum_{i \leq f} \text{area}(P_i) = 4\pi r^2$$

(since faces are non overlapping)




central projection




edges are drawn as non crossing geodesic arcs

Using spherical geometry

An alternative geometric proof of Euler Formula




Geodesic: arc of great circle (between p and q)




Girard Theorem

$$\alpha + \beta + \gamma = \pi + \frac{\text{area}(A,B,C)}{r^2}$$



Spherical drawing of an octahedron

$$\alpha + \beta + \gamma + \delta = 2\pi$$



- $$\sum_{j \leq k_i} \alpha_{ij} = (k_i - 2)\pi + \frac{\text{area}(P_i)}{r^2}$$
 - $$\sum_{r=1}^{\deg(v)} \alpha_r^v = 2\pi \quad \text{(sum of angles around each vertex } v)$$
 - $$\sum_{i \leq f} \text{area}(P_i) = 4\pi r^2$$
- compute the sum of angles in 2 ways
- $$\sum_{v \leq n} \sum_{r=1}^{\deg(v)} \alpha_r^v = \sum_{i \leq f} \sum_j \alpha_{ij} \rightarrow 2e\pi - 2f\pi + 4\pi = n2\pi$$
- take the unit sphere ($r = 1$)

Exercise 2 (soccer ball theorem)



Given a plane graph, where every face is a pentagon or a hexagon, and such that every vertex has degree 3, show that there must be exactly twelve pentagonal faces.

f_i := number of faces of degree i

Use a double counting argument (edges around vertices): $3v = 2e$

Count edges around faces): $5f_5 + 6f_6 = 2e = 3v$

Use Euler relation: $(5f_5 + 6f_6) - e + v = 2$

from previous relation (using $3v = 2e$): $-v + 2f_5 + 2f_6 = 4$

combining with: $5f_5 + 6f_6 = 3v$ we get $f_5 = 12$

MPRI 2-38-1: Algorithms and combinatorics for geometric graphs

Lecture 1, part II

Graph Drawing: Tutte barycentric method

september 18, 2024

Luca Castelli Aleardi



Graph drawing: introduction and applications

Graph drawing and data visualization

Global transportation system



Social networks







Roads, railways, ...



Parameterization problem
(known in Geometry Processing)

Compute a crossing-free drawing of planar graphs



Bennis et al., 1991
Maillot et al., 1993

Graph drawing: motivation


$$A_G = \begin{bmatrix} 0 & 1 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 & 0 \end{bmatrix}$$

Challenge: what kind of graph does A_G represent?

Graph drawing: motivation

$$A_G = \begin{bmatrix} 0 & 1 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 & 0 \end{bmatrix}$$

Challenge: what kind of graph does A_G represent?




adjacency matrix

$$A_G[i, j] = \begin{cases} 1 & \text{if } v_i \text{ is adjacent to } v_j \\ 0 & \text{otherwise} \end{cases}$$

Graph drawing: motivation

$$A_G = \begin{bmatrix} 0 & 1 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 & 0 \end{bmatrix}$$



Challenge: what kind of graph does A_G represent?



Graph drawing: motivation

$$A_G = \begin{bmatrix} 0 & 1 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 & 0 \end{bmatrix}$$




Challenge: what kind of graph does A_G represent?



Graph drawing: motivation

$$A_G = \begin{bmatrix} 0 & 1 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 & 0 \end{bmatrix}$$





Challenge: what kind of graph does A_G represent?



Graph drawing: motivation

$$A_G = \begin{bmatrix} 0 & 1 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 & 0 \end{bmatrix}$$






Challenge: what kind of graph does A_G represent?



Graph drawing: motivation

$$A_G = \begin{bmatrix} 0 & 1 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 & 0 \end{bmatrix}$$







Challenge: what kind of graph does A_G represent?



Graph drawing: motivation

$$A_G = \begin{bmatrix} 0 & 1 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 & 0 \end{bmatrix}$$


Challenge: what kind of graph does A_G represent?



Major results in Graph Drawing (for planar graphs)


Fáry theorem (1947) (exercise)

- Every (simple) planar graph admits a straight line planar embedding (no edge crossings)




Thm (Koebe-Andreev-Thurston) (not covered)

Every planar graph with n vertices is isomorphic to the intersection graph of n disks in the plane.



Thm (Steinitz, 1916)

3-connected planar graphs are the 1-skeletons of convex polyhedra




(Lecture 1)

Thm (Tutte barycentric method, 1963)

Every 3-connected planar graph G admits a barycentric representation ρ in $\mathbb{R}^2$.




$$\rho(v_i) = \sum_{j \in N(i)} w_{ij} \rho(v_j) \quad (\sum_j w_{ij} = 1 \text{ and } w_{ij} > 0)$$



Graph drawing paradigms

(Lecture 5)


Thm (Schnyder 1990)



Thm (De Fraysseix, Pack Pollack 1989)

FPP algorithm

(Lecture 2)




shift algorithm via Canonical orderings

linear time algorithms

$O(n) \times O(n)$ grid drawings

not trivial to implement

extremely fast: they can process millions of vertices per second



Spring embedder (Eades, 1984)
(Fruchterman and Reingold, 1991)



force-directed paradigm

easy to implement

pretty slow: $O(n^2)$ or $O(n \log n)$ time per iteration

$$F_a(v) = c_1 \cdot \sum_{(u,v) \in E} \log(\text{dist}(u, v)/c_2)$$

$$F_r(v) = c_3 \cdot \sum_{u \in V} \frac{1}{\sqrt{\text{dist}(u, v)}}$$



images from Kaufman Wagner (Springer, 2001)

[Tutte'63]

Tutte barycentric embedding

minimize the spring energy


$$E(p) := \sum_{(i,j) \in E} |\mathbf{x}(v_i) - \mathbf{x}(v_j)|^2 = \sum_{(i,j) \in E} (x_i - x_j)^2 + (y_i - y_j)^2$$

solve large sparse linear systems

$$\mathbf{x}(v_i) = \sum_{j \in N(i)} \frac{1}{\deg(v_i)} \mathbf{x}(v_j)$$

easy to implement

not very fast: they can process $\approx 10^4$ vertices per second




Straight-line planar drawings of planar graphs

Problem definition (Planarity testing, Embedding a planar graph)

Input: a planar graph


Output: the planar map (cellulaly embedded graph)



Problem definition (drawing in the plane)

Input: a planar graph (or planar map)

Output: a straight-line planar drawing (crossing-free)



Input of the problem: planar map

(a, b, c) (d, e, g) (i, g, b)



(a, c, d) (e, b, g) (i, b, a)

(d, c, e) (a, f, h)

(c, b, e) (a, h, i)

(a, d, f) (i, h, f)

(f, d, g) (i, f, g)





Computing a planar embedding

3-connectedness

Defini


G is 3-connected if

$\left\{ \begin{array}{l} \text{is connected and} \\ \text{the removal of one or two vertices} \\ \text{does not disconnect } G \end{array} \right.$




at least 3 vertices are required to disconnect the graph

cut-vertex




cut-pair



Menger Theorem

if G is 3-connected then for every pair of vertices u and v there exist 3 vertex disjoint paths (intersecting only at u and v)


(see Lecture on Schnyder woods, for a simple proof in the triangulated planar case)



3-connected planar graphs: Whitney theorem

Thm (Whitney, 1933)

3-connected planar graphs admit an unique embedding (up to homeomorphism and inversion of the sphere $\mathbb{S}^2$).






Remark: why 3-connectedness is important?

3-connected planar graphs: Whitney theorem

Thm (Whitney, 1933)

3-connected planar graphs admit an unique embedding (up to homeomorphism and inversion of the sphere $\mathbb{S}^2$).



$f_i :=$ number of faces of degree i


$$f_3 = 4$$

$$f_4 = 2$$


$$f_5 = 1$$

$$f_6 = 0$$

$$f_7 = 1$$



$$|F| = 9$$



$$f_3 = 4$$

$$f_4 = 2$$

$$f_5 = 0$$

$$f_6 = 2$$


$$f_7 = 0$$

two different (non equivalent) embeddings of the same graph

Bridges

G a connected graph

C a cycle



Bridges:= subgraphs induced by the edges of $E(G) \setminus E(C)$

Remarks

bridges can only intersect at the vertices of C

trivial bridges do not have inner vertices: loops, chords

from any two vertices of a bridge there exists one path internally disjoint from C

if G is non-separable then there are two vertices of attachment

k -bridge is a bridge with k vertices of attachment

equivalent bridges: same point of attachment (B_1 and B_2 , which are 3-bridges)


B_2 and B_4 are said to *avoid each other*

B_3 and B_4 are said to be *overlapping*

Two bridges B and B' are *skew* if there exist 4 vertices of attachment u, v (on B) and u', v' on B' which are listed consecutively on C : u, u', v, v'

B_3 and B_4 are skew

B_1 and B_2 are not skew




Block decomposition

Definition

A *block* is a maximal sub-graph (with respect to inclusion) that has no cut vertex

cut-vertex




Remark

A graph G is planar if and only if all its blocks are planar.

Lemma

Given a graph G all its blocks can be computed in linear time.

proof: Compute a DFS tree from an arbitrary vertex r



case 1:

$v = r$


For each vertex v compute:

$\text{depth}(v)$

$\text{lowpoint}(v)$

(process vertices in post-order)

$\text{lowpoint}(v) :=$ smallest depth of the extremity of a back (red) edge (w, z) (where w is a descendant of v)



case 2:

$v \neq r$



Bridges

Lemma


Given a 2-connected graph G we can in linear time either compute a circuit of G having at least two bridges, or certify that G is planar.

proof: Compute an arbitrary cycle C

Assume there is a single bridge B_1 (otherwise we are already done)



compute a path $P :=$ in B_1 from v_1 and v_2



assuming $P \neq B_1$

otherwise $G = C \cup P$ is planar

Bridges

Lemma

Let C a circuit of G . The graph G is planar if and only if:



The conflict graph of the bridges of C is bipartite (bridges are either outside or inside)

For every bridge B (with respect to C), the graph $H = B \cup C$ is planar

proof:

One direction: assume G is planar

Two bridges B and B' drawn both inside (or outside) cannot be overlapping
(no edge in the conflict graph between them)



The original graph would be non planar

Bridges

Lemma

Let C a circuit of G . The graph G is planar if and only if:



The conflict graph of the bridges of C is bipartite

For every bridge B (with respect to C), the graph $H = B \cup C$ is planar


proof:

Other direction (we want to embed the graph, without crossings)

Solution: since (inner) bridges are without conflict, we can add all (inner) bridges iteratively one by one



B_3 is not overlapping with B_1 and B_2



Embedding algorithm

Embed(G, C)

Compute the bridges of G with respect to C

Compute the conflict graph of $B_1, B_2, B_3, \dots$

if the conflict graph is not bipartite, return non-planar

For each bridge B of G (not a path):

let $G' := C \cup B$

let $C' := \text{extract}(G', C)$ (apply previous Lemma)

embed(G', C') (recursive call)

if G' is non-planar, return non-planar

return planar

Embedding algorithm $O(n^3)$

Embed(G, C)

Compute the bridges of G with respect to C $O(n)$

Compute the conflict graph of $B_1, B_2, B_3, \dots$ $O(n^2)$

if the conflict graph is not bipartite, return non-planar $O(n)$

For each bridge B of G (not a path):

let $G' := C \cup B$

let $C' := \text{extract}(G', C)$ (apply previous Lemma) $O(n')$

embed(G', C') (recursive call) $O(n)$ recursive calls

if G' is non-planar, return non-planar

return planar


Triangulating a planar graph

Lemma

Let G be a simple plane graph (cellularly embedded). Then it is possible to triangulate G in linear time obtaining a simple triangulation T (super graph of G).

proof:

Any idea?



Triangulating a planar graph



Lemma

Let G be a simple plane graph (cellularly embedded). Then it is possible to triangulate G in linear time obtaining a simple triangulation T (super graph of G).

proof:

Solution: triangulate faces

Problem: eliminate loops and multiple edges






Triangulating a planar graph

Lemma

Let G be a simple plane graph (cellularly embedded). Then it is possible to triangulate G in linear time obtaining a simple triangulation T (super graph of G).

proof:

Idea: eliminate loops and multiple edges (via edge flipping)






Triangulating a planar graph

Lemma

Let G be a simple plane graph (cellularly embedded). Then it is possible to triangulate G in linear time obtaining a simple triangulation T (super graph of G).

proof:

Idea: eliminate loops and multiple edges (via edge flipping)






Triangulating a planar graph

Lemma

Let G be a simple plane graph (cellularly embedded). Then it is possible to triangulate G in linear time obtaining a simple triangulation T (super graph of G).

proof:

Idea: eliminate loops and multiple edges (via edge flipping)




Tutte's planar embedding

Preliminaries: barycentric coordinates

$$q = \sum_i^n \alpha_i v_i \text{ (avec } \sum_i \alpha_i = 1)$$



coefficients $(\alpha_1, \dots, \alpha_n)$ are called *barycentric coordinates* of q
(relative to $v_1, \dots, v_n$)

Geometric interpretation
of barycentric coordinates



$$q = \alpha_0 v_0 + \alpha_1 v_1 + \alpha_2 v_2$$

$$q = \frac{\text{area}(v, v_1, v_2)v_0 + \text{area}(v_0, v, v_2)v_1 + \text{area}(v_0, v_1, v)v_2}{\text{area}(v_0, v_1, v_2)}$$




Tutte's theorem



Thm (Tutte barycentric method, 1963)

Every 3-connected planar graph G admits a convex representation ρ in R^2 .



Tutte's theorem




Thm (Tutte barycentric method, 1963)

Every 3-connected planar graph G admits a convex representation ρ in R^2 .

$$\rho : (V_G) \longrightarrow R^2$$

ρ is convex

the images of the faces of G are convex polygons



Tutte's theorem



Thm (Tutte barycentric method, 1963)

Every 3-connected planar graph G admits a convex representation ρ in R^2 .




$$\rho : (V_G) \longrightarrow R^2$$

ρ is barycentric the images of interior vertices are barycenters of their neighbors

$$\rho(v_i) = \sum_{j \in N(i)} w_{ij} \rho(v_j)$$

where w_{ij} satisfy $\sum_j w_{ij} = 1$, and $w_{ij} > 0$

according to Tutte: $w_{ij} = \frac{1}{\deg(v_i)}$





Tutte's theorem: main steps

- chose a cycle F (the outer face of G) in the right way
a cycle such that $G \setminus F$ is connected
(deletion of vertices and edges)

Tutte's theorem: main steps

- chose a cycle F (the outer face of G) in the right way
a cycle such that $G \setminus F$ is connected
(deletion of vertices and edges)
- choose a convex polygon P of size $k = |F|$
such that $\rho(F) = P$






Tutte's theorem: main steps

- chose a cycle F (the outer face of G) in the right way
a cycle such that $G \setminus F$ is connected
(deletion of vertices and edges)
- choose a convex polygon P of size $k = |F|$
such that $\rho(F) = P$
- solve equations for images of inner vertices $\rho(v_i)$:

$$\rho(v_i) = \sum_{j \in N(i)} w_{ij} \rho(v_j)$$

$$\rho(v_i) - \sum_{j \in N(i)} w_{ij} \rho(v_j) = 0$$


according to Tutte: $w_{ij} = \frac{1}{\deg(v_i)}$



Tutte's theorem: main steps



- chose a cycle F (the outer face of G) in the right way
a cycle such that $G \setminus F$ is connected
(deletion of vertices and edges)
- choose a convex polygon P of size $k = |F|$
such that $\rho(F) = P$
- solve two linear systems:

$$\left\{ \begin{array}{l} (I - W) \cdot \mathbf{x} = \mathbf{b}_x \\ (I - W) \cdot \mathbf{y} = \mathbf{b}_y \end{array} \right. \longleftrightarrow \left\{ \begin{array}{l} \rho_x(v_i) - \sum_{j \in N(i)} w_{ij} \rho_x(v_j) = 0 \\ \rho_y(v_i) - \sum_{j \in N(i)} w_{ij} \rho_y(v_j) = 0 \end{array} \right.$$



Tutte's theorem: main steps

- chose a cycle F (the outer face of G) in the right way
a cycle such that $G \setminus F$ is connected
(deletion of vertices and edges)
- choose a convex polygon P of size $k = |F|$
such that $\rho(F) = P$
- solve a linear system:



Validity of Tutte's theorem: main results


- show that the linear system admit a (unique) solution:

$$\begin{cases} (I - W) \cdot \mathbf{x} = \mathbf{b}_x \\ (I - W) \cdot \mathbf{y} = \mathbf{b}_y \end{cases} \quad \text{matrix } (I - W) \text{ is invertible}$$

- a barycentric drawing is planar: no edge crossing
- a 3-connected planar graph G has a non-separating cycle



Claim (existence of no-separating cycles)

In a 3-connected planar graph peripheral cycles are exactly the faces (of the embedding)




Validity of Tutte's theorem: main results

why 3-connectness and peripheral cycles are important:





Advantages of Tutte's drawing

- the drawing is guaranteed to be planar (no edge crossing)
- no need of the map structure
graph structure + a peripheral cycle
- very easy to implement: no need of sophisticated data structure or preprocessing



linear systems to solves

- nice drawings
(detection of symmetries)





Drawbacks of Tutte's drawing

- requires to solve linear systems of equations (of size n)

$$\begin{cases} (I - W) \cdot \mathbf{x} = \mathbf{b}_x \\ (I - W) \cdot \mathbf{y} = \mathbf{b}_y \end{cases} \quad \begin{array}{l} \text{complexity } O(n^3) \\ \text{or } O(n^{3/2}) \text{ with methods more involved} \end{array}$$

- exponential size of the resulting vertex coordinates (with respect to n)

- drawings are not always "nice"



Tutte's spring embedder: iterative version

- choose an outer face F , and a convex polygon P
- put exterior vertices $v \in F$ on the polygon
- repeat (until convergence)



for each inner vertex $v \in V_i$ compute

$$x_v = \frac{1}{\deg(v)} \sum_{(u,v) \in E} x_u$$

$$y_v = \frac{1}{\deg(v)} \sum_{(u,v) \in E} y_u$$

V_i inner vertices

(u, v) edge connecting v and u




Tutte's spring embedder: several interpretations

- choose an outer face F , and a convex polygon P
- put exterior vertices $v \in F$ on the polygon
- repeat (until convergence)

for each inner vertex $v \in V_i$ compute

$$x_v = \frac{1}{\deg(v)} \sum_{(u,v) \in E} x_u$$

$$y_v = \frac{1}{\deg(v)} \sum_{(u,v) \in E} y_u$$



Force directed method, with total force:

$$\mathbf{F}(v) = F_a(v) + F_r(v) = \sum_{(u,v) \in E} (\mathbf{p}_u - \mathbf{p}_v)$$

Resolution of linear systems

$$\rho(v_i) = \frac{1}{d_i} \sum_{j \in N(i)} \rho(v_j)$$

$$\begin{cases} (I - W) \cdot \mathbf{x} = \mathbf{b}_x \\ (I - W) \cdot \mathbf{y} = \mathbf{b}_y \end{cases}$$

Energy minimization

$$E(\rho) := \sum_{(i,j) \in E} |\rho(v_i) - \rho(v_j)|^2 = \sum_{(i,j) \in E} (x_i - x_j)^2 + (y_i - y_j)^2$$

find ρ minimizing

$$\begin{cases} E(\rho) \\ \text{subject to } \rho(v_k) = p_k = (x_k, y_k) \text{ (for exterior vertices } v_k) \end{cases}$$

Related drawing paradigm: force-directed algorithms

Spring electrical model (Fruchterman and Reingold, 1991)

(not covered)

```

area := W * L; {W and L are the width and length of the frame}
G := (V, E); {the vertices are assigned random initial positions}
k := sqrt(area / |V|);
function fa(x) := begin return x2 / k end;
function fr(x) := begin return k2 / x end;
for i := 1 to iterations do begin
    {calculate repulsive forces}
    for v in V do begin
        {each vertex has two vectors: .pos and .disp}
        v.disp := 0;
        for u in V do
            if (u ≠ v) then begin
                {δ is the difference vector between the positions of the two vertices}
                δ := v.pos - u.pos;
                v.disp := v.disp + (δ / |δ|) * fr(|δ|)
            end
        end
        {calculate attractive forces}
        for e in E do begin
            {each edge is an ordered pair of vertices .v and .u}
            δ := e.v.pos - e.u.pos;
            e.v.disp := e.v.disp - (δ / |δ|) * fa(|δ|);
            e.u.disp := e.u.disp + (δ / |δ|) * fa(|δ|)
        end
    end
    outside frame}
    for v in V do begin
        v.pos := v.pos + (v.disp / |v.disp|)
        v.pos.x := min(W / 2, max(-W / 2, v.pos.x));
        v.pos.y := min(L / 2, max(-L / 2, v.pos.y))
    end
end
end
    
```

$$F_a(u) = \sum_{(u,v) \in E} \frac{\|\mathbf{x}(u) - \mathbf{x}(v)\|}{K} (\mathbf{x}(v) - \mathbf{x}(u))$$

$$F_r(u) = \sum_{v \in V, v \neq u} \frac{-CK^2(\mathbf{x}(v) - \mathbf{x}(u))}{\|\mathbf{x}(u) - \mathbf{x}(v)\|^2}$$









Figure 64. Three-dimensional layouts of an icosahedron variant



ash85


Related drawing paradigm: spectral drawing

(not covered)


$$L_G[i, j] = \begin{cases} \deg(v_i) & \text{si } i = j \\ -A_G[i, j] & \text{otherwise} \end{cases}$$

$$E(\rho) := \sum_{(ij) \in E} \|\rho(v_i) - \rho(v_j)\|^2$$

$$\begin{cases} \min_{\underline{x}} E(\underline{x}) := \underline{x}^T L_G \underline{x} \\ \text{constraint: } \underline{x}^T \cdot \underline{x} = 1 \\ x_M = \sum_i x_i = 0 \quad \underline{x}^T \cdot \mathbf{1}_n = 0 \end{cases} \longrightarrow (x_1, \dots, x_d) = \left(\frac{v_2[i]}{\sqrt{\lambda_2}}, \frac{v_3[i]}{\sqrt{\lambda_3}}, \dots, \frac{v_{d+1}[i]}{\sqrt{\lambda_{d+1}}} \right)$$




(4elt graph, force-directed layout)




(b) The 4elt graph [10] $|V| = 15,606$, $|E| = 45,878$.

(4elt graph, spectral layout)
(image from Koren, 2005)



(degree-normalized (Koren))

$$\min_{\underline{x}} E(\underline{x}) := \frac{\underline{x}^T L_G \underline{x}}{\underline{x}^T \Delta \underline{x}}$$



(images from TD, INF562)

Tutte's theorem: the proof

First: existence and uniqueness of barycentric representations

Second: the barycentric representation defines a planar drawing (no edge crossing)

Third: characterization of non-separating cycles

(Some notions of) Spectral graph theory

Lemma (Laplacian and the number of spanning trees)


Let Q_G be the laplacian of a graph G , with n vertices.

Then the number of distinct spannig trees of G is:

(proof not covered)

$$(i \leq n) \quad \tau(G) = \det(Q_G[i])$$

$Q_G[i]$:= laplacian of Q_G , after removing the i -th row and column



$$Q_G[1] = \begin{bmatrix} \cancel{5} & \cancel{-3} & \cancel{-2} \\ -3 & 4 & -1 \\ -2 & -1 & 3 \end{bmatrix}$$

$$Q_G[2] = \begin{bmatrix} 5 & \cancel{-3} & \cancel{-2} \\ \cancel{-3} & \cancel{4} & \cancel{-1} \\ -2 & -1 & 3 \end{bmatrix}$$

$$Q_G[3] = \begin{bmatrix} 5 & -3 & \cancel{-2} \\ -3 & 4 & -1 \\ \cancel{-2} & \cancel{-1} & \cancel{3} \end{bmatrix}$$

$$\det(Q_G[1]) = \begin{bmatrix} 4 & -1 \\ -1 & 3 \end{bmatrix} = 11$$

$$\det(Q_G[2]) = \begin{bmatrix} 5 & -2 \\ -2 & 3 \end{bmatrix} = 11$$

$$\det(Q_G[3]) = \begin{bmatrix} 5 & -3 \\ -3 & 4 \end{bmatrix} = 11$$

First: existence and uniqueness of barycentric representations

Theorem (Tutte)

Let G be a 3-connected planar graph with n vertices, and F a peripheral cycle (such that $G \setminus F$ is connected). Let P be a convex polygon, such that $\rho(F) = P$. Then the barycentric representation ρ exists (and is unique)

Goal: show that the two systems above admit a solution (unique)

Let us denote $\rho(v_i) := (x_i, y_i) = \mathbf{x}_i$ the coordinates of vertex v_i

$$\begin{cases} (I - W) \cdot \mathbf{x} = \mathbf{b}_x \\ (I - W) \cdot \mathbf{y} = \mathbf{b}_y \end{cases} \longleftrightarrow \begin{aligned} \rho(v_i) &= \sum_{j=1}^n w_{ij} \rho(v_j) & i = 1, \dots, (n-k) \\ \rho(v_i) &= \sum_{j \in N(i)} w_{ij} \rho(v_j) \end{aligned}$$

$\mathbf{x} = [x_1, x_2, \dots, x_{n-k}]$
 $\mathbf{y} = [y_1, y_2, \dots, y_{n-k}]$
(coordinates of inner vertices)

(one equation for each inner vertex)

Theorem (Tutte)

Let G be a 3-connected planar graph with n vertices, and F a peripheral cycle (such that $G \setminus F$ is connected). Let P be a convex polygon, such that $\rho(F) = P$. Then the barycentric representation ρ exists (and is unique)

$$\begin{cases} (I - W) \cdot \mathbf{x} = \mathbf{b}_x \\ (I - W) \cdot \mathbf{y} = \mathbf{b}_y \end{cases} \iff \rho(v_i) = \sum_{j=1}^n w_{ij} \rho(v_j) \quad \begin{matrix} i = 1, \dots, (n - k) \\ \text{(one equation for} \\ \text{each inner vertex)} \end{matrix}$$

($I - W$ is not symmetric)

$$\mathbf{x} = [x_1, x_2, \dots, x_{n-k}]$$

$$\mathbf{y} = [y_1, y_2, \dots, y_{n-k}]$$

(coordinates of inner vertices)


Example (to help intuition)

$$\begin{bmatrix} 1 & -\frac{1}{4} \\ -\frac{1}{3} & 1 \end{bmatrix} \begin{bmatrix} x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} b_{4x} \\ b_{5x} \end{bmatrix} \quad \begin{cases} \rho(v_4) = \frac{1}{4}\rho(v_1) + \frac{1}{4}\rho(v_2) + \frac{1}{4}\rho(v_3) + \frac{1}{4}\rho(v_5) \\ \rho(v_5) = \frac{1}{3}\rho(v_2) + \frac{1}{3}\rho(v_3) + \frac{1}{3}\rho(v_4) \end{cases}$$

$$\begin{bmatrix} 1 & -\frac{1}{4} \\ -\frac{1}{3} & 1 \end{bmatrix} \begin{bmatrix} y_4 \\ y_5 \end{bmatrix} = \begin{bmatrix} b_{4y} \\ b_{5y} \end{bmatrix} \quad \begin{cases} \rho(v_4) - \frac{1}{4}\rho(v_5) = \frac{1}{4}\rho(v_1) + \frac{1}{4}\rho(v_2) + \frac{1}{4}\rho(v_3) \\ -\frac{1}{3}\rho(v_4) + \rho(v_5) = \frac{1}{3}\rho(v_2) + \frac{1}{3}\rho(v_3) \end{cases}$$

$$N(v_4) = \{v_1, v_2, v_3, v_5\}$$

$$N(v_5) = \{v_2, v_3, v_4\}$$



Theorem (Tutte)

Let G be a 3-connected planar graph with n vertices, and F a peripheral cycle (such that $G \setminus F$ is connected). Let P be a convex polygon, such that $\rho(F) = P$. Then the barycentric representation ρ exists (and is unique)

($I - W$ is not symmetric)

$$\begin{cases} (I - W) \cdot \mathbf{x} = \mathbf{b}_x \\ (I - W) \cdot \mathbf{y} = \mathbf{b}_y \end{cases}$$

$$\mathbf{x} = [x_1, x_2, \dots, x_{n-k}]$$

$$\mathbf{y} = [y_1, y_2, \dots, y_{n-k}]$$

(coordinates of inner vertices)

(M is symmetric)

$$\begin{cases} M \cdot \underline{x} = \underline{a_x} \\ M \cdot \underline{y} = \underline{a_y} \end{cases}$$

$$i = 1, \dots, (n - k)$$

$$\rho(v_i) = \sum_{j \in N(i)} w_{ij} \rho(v_j) \iff \deg(v_i) \rho(v_i) - \sum_{j \in N(i)} \rho(v_j) = 0$$

$$\begin{bmatrix} 3 & 1 & 1 & 1 & 0 \\ -1 & 4 & -1 & -1 & -1 \\ -1 & -1 & 4 & -1 & -1 \\ -1 & -1 & -1 & 4 & -1 \\ 0 & -1 & -1 & -1 & 3 \end{bmatrix}$$

(k constraints, for the k outer vertices)

$$M = \begin{bmatrix} 4 & -1 \\ -1 & 3 \end{bmatrix}$$



$$N(v_4) = \{v_1, v_2, v_3, v_5\}$$

$$N(v_5) = \{v_2, v_3, v_4\}$$

$$\begin{cases} \rho(v_4) - \frac{1}{4}\rho(v_5) = \frac{1}{4}\rho(v_1) + \frac{1}{4}\rho(v_2) + \frac{1}{4}\rho(v_3) \\ -\frac{1}{3}\rho(v_4) + \rho(v_5) = \frac{1}{3}\rho(v_2) + \frac{1}{3}\rho(v_3) \end{cases}$$

$$\begin{cases} 4\rho(v_4) - \rho(v_5) = \rho(v_1) + \rho(v_2) + \rho(v_3) \\ -\rho(v_4) + 3\rho(v_5) = \rho(v_2) + \rho(v_3) \end{cases}$$

($n - k$) equations for inner vertices



Existence and uniqueness of barycentric representations

$$\deg(v_i)\rho(v_i) - \sum_{j \in N(i)} \rho(v_j) = 0 \qquad \begin{cases} M \cdot \underline{x} = \underline{a_x} \\ M \cdot \underline{y} = \underline{a_y} \end{cases}$$

$$\tau(G/F) > 0 \longrightarrow \det(M) = \tau(Q_{G/F}) > 0$$

(G after the contraction of F is still connected)


$G \setminus F$ has at least one spanning tree

$$F := \{v_1, v_2, v_3\}$$


(k constraints, for the k outer vertices)

$$Q_G = \begin{bmatrix} -3 & -1 & -1 & -1 & 0 \\ -1 & 4 & -1 & -1 & -1 \\ -1 & -1 & 4 & -1 & -1 \\ -1 & -1 & -1 & 4 & -1 \\ 0 & -1 & -1 & -1 & 3 \end{bmatrix}$$

($n - k$) equations for inner vertices



$$G/F = G/\{v_1, v_2, v_3\}$$



$$\begin{cases} \rho(v_4) - \frac{1}{4}\rho(v_5) = \frac{1}{4}\rho(v_1) + \frac{1}{4}\rho(v_2) + \frac{1}{4}\rho(v_3) \\ -\frac{1}{3}\rho(v_4) + \rho(v_5) = \frac{1}{3}\rho(v_2) + \frac{1}{3}\rho(v_3) \end{cases}$$

$$\begin{cases} 4\rho(v_4) - \rho(v_5) = \rho(v_1) + \rho(v_2) + \rho(v_3) \\ -\rho(v_4) + 3\rho(v_5) = \rho(v_2) + \rho(v_3) \end{cases}$$

$$Q_{G/F} = \begin{bmatrix} -5 & 3 & 2 \\ -3 & 4 & -1 \\ -2 & -1 & 3 \end{bmatrix}$$

$$M = \begin{bmatrix} 4 & -1 \\ -1 & 3 \end{bmatrix}$$

Existence and uniqueness of barycentric representations

$$\deg(v_i)\rho(v_i) - \sum_{j \in N(i)} \rho(v_j) = 0 \quad \left\{ \begin{array}{l} M \cdot \underline{x} = \underline{a_x} \\ M \cdot \underline{y} = \underline{a_y} \end{array} \right.$$

the outer face F is not separating

$$\tau(G \setminus F) > 0$$

since $G \setminus F$ is connected

$$\tau(G/F) > 0 \longrightarrow \det(M) = \tau(Q_{G/F}) > 0$$


(G after the contraction of F is still connected)

$G \setminus F$ has at least one spanning tree


M admits inverse $\square$

(k constraints, for the k outer vertices)

$$Q_G = \begin{bmatrix} 3 & -1 & -1 & -1 & 0 \\ -1 & 4 & -1 & -1 & -1 \\ -1 & -1 & 4 & -1 & -1 \\ -1 & -1 & -1 & 4 & -1 \\ 0 & -1 & -1 & -1 & 3 \end{bmatrix}$$



$$G/F = G/\{v_1, v_2, v_3\}$$



$$Q_G[1, 2, 3] := M = \begin{bmatrix} 4 & -1 \\ -1 & 3 \end{bmatrix}$$

$$Q_{G/F} = \begin{bmatrix} 5 & 3 & 2 \\ -3 & 4 & -1 \\ -2 & -1 & 3 \end{bmatrix}$$

$$\left. \begin{array}{l} \rho(v_4) - \frac{1}{4}\rho(v_5) = \frac{1}{4}\rho(v_1) + \frac{1}{4}\rho(v_2) + \frac{1}{4}\rho(v_3) \\ -\frac{1}{3}\rho(v_4) + \rho(v_5) = \frac{1}{3}\rho(v_2) + \frac{1}{3}\rho(v_3) \end{array} \right\}$$

($n - k$) equations for inner vertices

First: existence and uniqueness of barycentric representations

Lemma *The barycentric representation ρ exists (and is unique)*

Second proof: (via energy minimization)

Let us denote $\rho(v_i) := (x_i, y_i) = \mathbf{x}_i$

spring energy for edge (v_i, v_j)

$$E(v_i, v_j) := D_{ij} \|\rho(v_i) - \rho(v_j)\|^2$$

Consider the spring energy of the whole system (of all inner edges):

$$E(\rho) := \sum_{e=(i,j) \in E} D_{ij} \|\rho(v_i) - \rho(v_j)\|^2 = \sum_{(i,j) \in E} D_{ij} [(x_i - x_j)^2 + (y_i - y_j)^2]$$

Rewrite the sum:



$$E(\rho) := \frac{1}{2} \sum_{v_i \in V} \sum_{j \in N_i} D_{ij} \|\mathbf{x}_i - \mathbf{x}_j\|^2$$

to find the coordinates $\mathbf{x}_i$ minimizing the energy, compute the gradient of E :

$$\frac{\partial E}{\partial \mathbf{x}_i} = 0 \quad \sum_{j \in N_i} D_{ij} (\mathbf{x}_i - \mathbf{x}_j) = \sum_{j \in N_i} D_{ij} \mathbf{x}_i - \sum_{j \in N_i} D_{ij} \mathbf{x}_j = 0 \quad \mathbf{x}_i = \sum_{j \in N_i} \left[\frac{D_{ij}}{\sum_{j \in N_i} D_{ij}} \right] \mathbf{x}_j = \sum_{j \in N_i} w_{ij} \mathbf{x}_j$$

Remark: the solution is **not degenerate**, because of boundary constraints
(to be proved... later)

physical analogy



(spring energy)

$$E(x) = \frac{1}{2} k x^2$$

$$F(x) = kx$$

$$w_{ij} := \left[\frac{D_{ij}}{\sum_{j \in N_i} D_{ij}} \right]$$



barycentric coordinates

Second: the barycentric representation defines a planar drawing

Theorem (Tutte)

Let G be a 3-connected planar graph with n vertices, and F a non-separating cycle (such that $G \setminus F$ is connected). Let P be a convex polygon, such that $\rho(F) = P$. Then the barycentric representation defines a planar drawing (no edge crossing)

Proof: (we follow the presentation given by Jeff Erickson)




Second: the barycentric representation defines a planar drawing

Lemma (outer face)

In any Tutte embedding the image of every inner vertex v is a point lying in the interior of the outer face (the convex polygon)


Proof:

Remark: (every inner vertex v is a barycentric combination of its neighbors u_i)





$v :=$ farthest point


All the neighbors of v should lie on the same line



G is connected, then (by induction) there is a path $P = \{z_1, z_2, \dots\}$




v is in the interior of C
then we are done



All the neighbors of v should lie on the same boundary between x and y , and also all the inner vertices reachable from v

there is a cut-pair x, y
contradicting the 3-conn.






3-connectedness is important

Second: the barycentric representation defines a planar drawing

Lemma (both sides)

Given an inner vertex v and a line l passing through its image $\rho(v)$ either all neighbors of v lie on l , or there are neighbors on both sides of l .

Proof:



impossible (v must be in the interior or the convex hull)

Lemma (convexity)


Every face in the Tutte embedding is a convex polygon.

Proof:

By contradiction, assume f is not convex


There must be a reflex angle at v

All neighbors of v must lie in the concave region between (u, v) and (w, v)



all neighbors must lie in the half-plane H^+

contradicting previous Lemma



Second: the barycentric representation defines a planar drawing

Lemma (half-plane)

Let H^+ be an half-plane containig at least one vertex of G . Then the sub-graph of G induced by all the vertices lying in H^+ is connected.


Proof:

$t :=$ vertex with larget y -coord
(remark: t must lie on the convex hull)

Let u be an arbitrary vertex in H^+

claim: there is a path from u to t
(with non-decreasing y -coordinates)

assume $u^y < t^y$ (otherwise the claim is trivial)



G is connected, then there is $v \in U$ with neighbors in both H^+ and H^-
(because previous Lemma)

apply induction to the vertex w neighbor of v : since $w^y > v^y$ we can
find a path from v to the boundary

Second: the barycentric representation defines a planar drawing

Lemma (non-degeneracy)

No vertex is collinear with all its neighbors.


Proof: (by contradiction u has all its neighbors on l)

The induced graphs $G(V^+)$ and $G(V^-)$ are connected (previous lemma)

U := set of vertices reachable from u and whose neighbors all lie on l


W := set of vertices lying on l having at least one neighbor not in U

G is 3-connected, so $|W| \geq 3$



contract all edges in $G(V^+)$ and $G(V^-)$
(edge contraction preserve planarity)

($K_{3,3}$ contradicts the planarity of G)




The Jordan curve theorem

Theorem

Any simple closed curve C in the plane partitions $\mathbb{R}^2$ into two disjoint arcwise-connected open sets.

($Ext(C)$ and $Int(C)$ are closed sets)

$$Ext(C) \cap Int(C) = C$$




Remark:

Any arc joining a point p in the (open) interior to a point q in the (open) exterior must meet C at least once.

Jordan curve Theorem (reformulation)

Let G a graph embedded on $\mathbb{S}^2$. Then G disconnects $\mathbb{S}^2$ if and only if it contains a circuit



Second: the barycentric representation defines a planar drawing



Lemma (Geelen)

Let us consider an edge $e = (u, v)$ incident to two faces f and f' , whose remaining vertices are in two sets S and S' . Consider an arbitrary path P from one vertex in S to one vertex in S' . Then every path from u to v either consists of the edge (u, v) or contains a vertex of the path P .

Proof:

$P' :=$ a curve crossing (u, v) , lying inside $f \cup f'$

Consider an arbitrary planar embedding of G



The closed curve $C = P \cup P'$ separates u from v

Then every path from u to v must cross C (by Jordan curve thm)

Second: the barycentric representation defines a planar drawing

Lemma (Split Faces)

Let us consider an edge $e = (u, v)$ incident to two faces f and f' , whose remaining vertices are in two sets S and S' . Consider a line l passing through u and v . Then the vertices in S and S' lie on opposite sides with respect to l (and there is no vertex on l).

Proof: (by contradiction: assume there is $s, s' \in H^-$)

$\exists s_1, s'$ are strictly below l (degeneracy Lemma)

The graph included in H^- is connected (half-space Lemma): then there exists a path P from s to s'


$\exists u', v' \in H^+$ above l (degeneracy Lemma)

there exists a path Q' from u' to v' (above l)

consider the path $Q := Q' \cup (u, u') \cup (v, v')$ (above l)

Apply Geelen's Lemma: path Q should cross P (impossible)

contradiction: the path Q does not cross P , and avoids the arc (u, v)



Second: the barycentric representation defines a planar drawing

Final Lemma (no edge crossings)

The Tutte embedding of G is crossing-free (faces are non-overlapping).

Proof:

Claim: a generic point cannot lie into two different faces f_1 and f_2

Strategy: draw a (generic) line from p to infinity (crossing edges only at their interior)

