

Proposed direct test of a certain type of noncontextuality in quantum mechanics

Karl Svozil*

*Institut für Theoretische Physik, Vienna University of Technology,
Wiedner Hauptstraße 8-10/136, A-1040 Vienna, Austria*

Abstract

The noncontextuality of quantum mechanics can be directly tested by a novel type of experiment measuring two entangled particles with more than two outcomes per particle. The two associated contexts are “interlinked” by common observables.

PACS numbers: 03.65.Ta,03.65.Ud

Keywords: quantum contextuality

Quantum value indefiniteness [1] refers to the impossibility of a consistent coexistence of certain complementary, operationally incompatible quantum observables. It is inferred from three sources: (i) from quantum violations of constraints on classical probability distributions termed ‘*conditions of possible experience*’ by Boole [2], also known as the Boole-Bell type inequalities [3], (ii) from the Kochen-Specker theorem [4–6], as well as (iii) from the Greenberger-Horne-Zeilinger [7, 8] theorem. Formally, these results are related to the “scarcity” or even total absence of two-valued states identifiable as (classical) truth assignments on the entire range of quantum observables. In what follows, quantum contextuality [9–13] will be identified with the assertion that the result of a measurement depends on what other observables are comeasured alongside of it. It is one conceivable (but not necessary [14]) quasi-classical interpretation of quantum value indefiniteness, thereby counterfactually maintaining the “physical existence” of the full domain of possible physical observables.

There exist other notions of contextuality based upon violations of some bounds on, or conditions imposed by, classical probabilities. In their extreme form, these amount to all-or-nothing type contradictions between noncontextual hidden variables and quantum mechanics. The corresponding experimental tests indicate the occurrence of this type of quantum contextuality [15–23]. These findings utilize subsequent measurements of quantum observables contributing to a contradiction with their classical counterparts, but they have no direct bearing on the novel type of experiments proposed here which aim at testing another, more direct form of quantum contextuality.

A quantum mechanical context [13] is a “maximal collection of comeasurable observables” within the nondistributive structure of quantum propositions. It can be formalized by a single “maximal” self-adjoint operator, such that every collection of mutually compatible comeasurable operators (such as projections corresponding to yes–no propositions) are functions thereof [24, § 84].

Different contexts can be *interlinked* at one or more common observable(s) whose Hilbert space representation is identical and independent of the contexts they belong to. The context independence of the representation of observables by operators (e.g., projectors) in Hilbert space suggests that quantum contextuality, if it exists, manifests itself in random and uncontrollable single-particle outcomes. A necessary condition for the interlinking of two or more contexts by link observable(s) is the requirement that the dimensionality of the Hilbert space must exceed two, since for lower dimensional Hilbert spaces the maximal operators “decay” into separate, isolated “trivial” Boolean sublogics without any common observable. This is also the reason for similar dimensional condi-

tions on the theorems by Gleason, as well as by Kochen and Specker.

In what follows we propose an experiment capable of directly testing the contextuality hypothesis; alas via counterfactual elements of physical reality. Indeed, counterfactual reasoning might be considered less desirable than direct measurements, as it involves an additional logical inference step rather than a straight empirical finding.

In the proposed experiment, two different contexts or, equivalently, two noncommuting maximal observables, are simultaneously measured on a pair of spin one particles in a singlet state [11, 25, 26]. The contexts are fine-tuned to allow a common *single observable interlinking* them. Although the proposal possesses some conceptual similarities to Einstein-Podolsky-Rosen type experiments, the quantum states as well as the structure of the observables are different.

We shall first consider the contexts originally proposed by Kochen and Specker [4, pp. 71-73], referring to the change in the energy of the lowest orbital state of orthohelium resulting from the application of a small electric field with rhombic symmetry. The terms *Kochen-Specker contexts* and (maximal) *Kochen-Specker operators* will be used synonymously. More explicitly, the maximal Kochen-Specker operators associated with this link configuration can be constructed from the spin one observables (e.g., Refs. [27, 28]) in arbitrary directions measured in spherical coordinates

$$J(\theta, \phi) = \begin{pmatrix} \cos \theta & \frac{e^{-i\phi} \sin \theta}{\sqrt{2}} & 0 \\ \frac{e^{i\phi} \sin \theta}{\sqrt{2}} & 0 & \frac{e^{-i\phi} \sin \theta}{\sqrt{2}} \\ 0 & \frac{e^{i\phi} \sin \theta}{\sqrt{2}} & -\cos \theta \end{pmatrix}, \quad (1)$$

where $0 \leq \theta \leq \pi$ stands for the polar angle in the x - z -plane taken from the z -axis, and $0 \leq \phi < 2\pi$ is the azimuthal angle in the x - y -plane taken from the x -axis. The orthonormalized eigenvectors associated with the eigenvalues $+1, 0, -1$ of $J(\theta, \phi)$ in Eq. (1) are

$$\begin{aligned} x_{+1} &= e^{i\delta_{+1}} \left(e^{-i\phi} \cos^2 \frac{\theta}{2}, \frac{1}{\sqrt{2}} \sin \theta, e^{i\phi} \sin^2 \frac{\theta}{2} \right), \\ x_0 &= e^{i\delta_0} \left(-\frac{1}{\sqrt{2}} e^{-i\phi} \sin \theta, \cos \theta, \frac{1}{\sqrt{2}} e^{i\phi} \sin \theta \right), \\ x_{-1} &= e^{i\delta_{-1}} \left(e^{-i\phi} \sin^2 \frac{\theta}{2}, -\frac{1}{\sqrt{2}} \sin \theta, e^{i\phi} \cos^2 \frac{\theta}{2} \right), \end{aligned} \quad (2)$$

where $\delta_{\pm 1}$, and δ_0 stand for arbitrary phases.

For real $\alpha \neq \beta \neq \gamma \neq \alpha$, the maximal Kochen and Specker operators [4] are defined by

$$\begin{aligned} C_{KS}(\alpha, \beta, \gamma) &= \frac{1}{2} [(\alpha + \beta - \gamma)J^2(\frac{\pi}{2}, 0) + (\alpha - \beta + \gamma)J^2(\frac{\pi}{2}, \frac{\pi}{2}) + (\beta + \gamma - \alpha)J^2(0, 0)], \\ C'_{KS}(\alpha, \beta, \gamma) &= \frac{1}{2} [(\alpha + \beta - \gamma)J^2(\frac{\pi}{2}, \frac{\pi}{4}) + (\alpha - \beta + \gamma)J^2(\frac{\pi}{2}, \frac{3\pi}{4}) + (\beta + \gamma - \alpha)J^2(0, 0)]. \end{aligned} \quad (3)$$




FIG. 1: Diagrammatical representation of two interlinked Kochen-Specker contexts: Greechie (orthogonality) diagram representing two tripods with a common leg: points stand for individual basis vectors, and entire contexts — in this case the one-dimensional linear subspaces spanned by the vectors of the orthogonal tripods — are drawn as smooth curves.

Their common spectrum of eigenvalues is α , β and γ , corresponding to the eigenvectors $(0, 1, 0)$, $(1, 0, 1)$, $(-1, 0, 1)$ of C_{KS} , and $(0, 1, 0)$, $(-i, 0, 1)$, $(i, 0, 1)$ of C'_{KS} , respectively. The resulting orthogonality structure of propositions is depicted in Fig. 1.

In order to be able to use the type of counterfactual inference employed by an Einstein-Podolsky-Rosen setup, a multipartite quantum state has to be chosen which satisfies the *uniqueness property* [29] with respect to the two Kochen-Specker contexts such that knowledge of a measurement outcome of one particle entails the certainty that, if this observable were measured on the other particle(s) as well, the outcome of the measurement would be a unique function of the outcome of the measurement actually performed. Consider the two spin-one particle singlet state $|\varphi_s\rangle = (1/\sqrt{3})(-|00\rangle + |-+\rangle + |+-\rangle)$ and identify with the spin states the directions in Hilbert space according to Eqs. (2); i.e., with $|+\rangle = (1, 0, 0)$, $|0\rangle = (0, 1, 0)$, and $|-\rangle = (0, 0, 1)$; hence in the Kronecker product representation, $|\varphi_s\rangle = (1/\sqrt{3})(0, 0, 1, 0, -1, 0, 1, 0, 0)$. This singlet state is form invariant under spatial rotations (but not under all unitary transformations [28]) and satisfies the uniqueness property (see below), just as the ordinary Bell singlet state of two spin one-half quanta (we cannot use these because they are limited to 2×2 dimensions, with merely two dimensions per quantum). Hence, it is possible to employ a similar counterfactual argument and establish two elements of physical reality according to the Einstein-Podolsky-Rosen criterion for the two interlinked Kochen-Specker contexts C_{KS} as well as C'_{KS} .




FIG. 2: Greechie diagram of two contexts in four-dimensional Hilbert space interconnected by two link observables.

When combined with the singlet state $|\varphi_s\rangle$, two “collinear” Kochen-Specker contexts yield

$$\begin{aligned} \text{Tr} \{ |\varphi_s\rangle \langle \varphi_s| \cdot [C_{KS}(\alpha, \beta, \gamma) \otimes C_{KS}(\delta, \varepsilon, \zeta)] \} &= \\ = \text{Tr} \{ |\varphi_s\rangle \langle \varphi_s| \cdot [C'_{KS}(\alpha, \beta, \gamma) \otimes C'_{KS}(\delta, \varepsilon, \zeta)] \} &= \frac{1}{3} [\alpha\delta + \beta\varepsilon + \gamma\zeta]. \end{aligned} \quad (4)$$

As a consequence, in this configuration the uniqueness property manifests itself by the unique joint occurrence of the outcomes associated with $\alpha \leftrightarrow \delta$ (corresponding to the proposition associated with the link observable between C_{KS} and C'_{KS}), as well as $\beta \leftrightarrow \varepsilon$ and $\gamma \leftrightarrow \zeta$. Thus, by counterfactual inference, if the contexts measured on both sides are identical, whenever α , β or γ is registered on one side, δ , ε or ζ is measured on the other side, respectively, and *vice versa*.

We are now in the position to formulate a testable criterion for (non)contextuality: Contextuality predicts that there exist outcomes associated with α on one context C_{KS} which are accompanied by the outcomes ε or ζ for the other context C'_{KS} ; likewise δ should be accompanied by β and γ . The quantum mechanical expectation values can be obtained from

$$\text{Tr} \{ |\varphi_s\rangle \langle \varphi_s| \cdot [C_{KS}(\alpha, \beta, \gamma) \otimes C'_{KS}(\delta, \varepsilon, \zeta)] \} = \frac{1}{6} [2\alpha\delta + (\beta + \gamma)(\varepsilon + \zeta)]. \quad (5)$$

As a consequence, the outcomes α - ε , α - ζ , as well as β - δ and γ - δ indicating contextuality do not occur. This is in contradiction with the contextuality hypothesis.

Another context configuration in four-dimensional Hilbert space drawn in Fig. 2 consists of two

contexts which are interconnected by *two* common link observables. The two context operators

$$C(\alpha, \beta, \gamma, \delta) = \text{diag}(\alpha, \beta, \gamma, \delta), \quad C'(\alpha, \beta, \gamma, \delta) = \text{diag}\left(\frac{\alpha+\beta}{2}, \frac{\alpha-\beta}{2}, \gamma, \delta\right) \quad (6)$$

have identical eigenvalue spectra containing mutually different real eigenvalues α , β , γ and δ .

Consider the singlet state of two spin-3/2 observables $|\psi_s\rangle = \frac{1}{2}(|\frac{3}{2}, -\frac{3}{2}\rangle - |-\frac{3}{2}, \frac{3}{2}\rangle - |\frac{1}{2}, -\frac{1}{2}\rangle + |-\frac{1}{2}, \frac{1}{2}\rangle)$ satisfying the uniqueness property for all spatial directions. The four different spin states can be identified with the cartesian basis of fourdimensional Hilbert space $|\frac{3}{2}\rangle = (1, 0, 0, 0)$, $|\frac{1}{2}\rangle = (0, 1, 0, 0)$, $|-\frac{1}{2}\rangle = (0, 0, 1, 0)$, and $|-\frac{3}{2}\rangle = (0, 0, 0, 1)$, respectively. When combined with the singlet state $|\psi_s\rangle$, two ‘‘collinear’’ contexts yield

$$\begin{aligned} \text{Tr}\{|\psi_s\rangle\langle\psi_s| \cdot [C(\alpha, \beta, \gamma, \delta) \otimes C(\varepsilon, \zeta, \eta, \nu)]\} &= \frac{1}{4}[\alpha\nu + \beta\eta + \gamma\zeta + \delta\varepsilon], \\ \text{Tr}\{|\psi_s\rangle\langle\psi_s| \cdot [C'(\alpha, \beta, \gamma, \delta) \otimes C'(\varepsilon, \zeta, \eta, \nu)]\} &= \frac{1}{8}[2(\alpha\nu + \beta\eta) + (\gamma + \delta)(\varepsilon + \zeta)]. \end{aligned} \quad (7)$$

As a consequence, in this configuration the uniqueness property manifests itself by the unique joint occurrence of the outcomes associated with $\alpha \leftrightarrow \nu$ and $\beta \leftrightarrow \eta$ (corresponding to the proposition associated with the two link observables between C and C'), as well as $\gamma \leftrightarrow \zeta$ and $\delta \leftrightarrow \varepsilon$. Thus, by counterfactual inference, if the contexts measured on both sides are identical, whenever α or β , and γ or δ is registered on one side, ν or η , and ζ or ε is measured on the other side, respectively, and *vice versa*.

Compared to the previous Kochen-Specker contexts, this configuration has the additional advantage that — in the absence of any criterion for outcome preference — Jayne’s principle [30] suggests that contextuality predicts totally uncorrelated outcomes associated with a maximal unbiased of the two common link observables, resulting in the equal occurrence of the joint outcomes $\gamma\text{--}\eta$, $\gamma\text{--}\nu$, $\delta\text{--}\eta$, and $\delta\text{--}\nu$. The quantum mechanical predictions are based on the expectation values

$$\text{Tr}\{|\psi_s\rangle\langle\psi_s| \cdot [C(\alpha, \beta, \gamma, \delta) \otimes C'(\varepsilon, \zeta, \eta, \nu)]\} = \frac{1}{8}[(\gamma + \delta)(\varepsilon + \zeta) + 2(\beta\eta + \alpha\nu)]. \quad (8)$$

As a consequence, there are no outcomes $\gamma\text{--}\eta$, $\gamma\text{--}\nu$, $\delta\text{--}\eta$, and $\delta\text{--}\nu$, which is in contradiction to the contextuality postulate.

One of the conceivable criticisms against the presented arguments is that the configurations considered, although containing complementary contexts, still allow even a full, separable set of two-valued states, and therefore need no contextual interpretation. However, it is exactly these Kochen-Specker type contexts which enter the Kochen-Specker argument. Hence, they should not

be interpreted as separate, isolated sublogics, but as parts of a continuum of sublogics, containing the finite structure devised by Kochen and Specker and others.

One could also point out that it might suffice to *prepare* the particle in some link state “along” one context, and then *measure* its state “along” a different context “containing” the same link observable. This could for instance in the three-dimensional configuration be realized by two successive three-port beam splitters arranged serially. In such a configuration, if the outcomes of the two beam splitters do not coincide at the link observable, then noncontextuality is disproved; likewise, if there is a perfect correlation between the link state prepared and the link observable measured, then contextuality could be disproved. This configuration might be criticized by proponents of contextuality as being too restrictive, since there is a preselection, effectively fixing the preparation state corresponding to the link observable.

Third, one could reprehend that the entangled particles cannot be thought of as isolated and that the singlet state enforces noncontextuality by the way it is constructed. This criticism could be counterpointed by noting that it is exactly this kind of configurations which yield violations of Boole-Bell type *conditions of physical experience*.

The situation can be summarized as follows. The direct measurement of more than one context on a single particle is blocked by quantum complementarity. For the counterfactual “workaround” to measure two noncommuting interlinked contexts on pairs of spin-one and spin three-half particles in singlet states, quantum mechanics predicts noncontextual behavior. Because of the lack of a uniqueness property, counterfactual inference of configurations with more than two particles are impossible .

* Electronic address: `svozil@tuwien.ac.at`; URL: <http://tph.tuwien.ac.at/~svozil>

- [1] A. Peres, *American Journal of Physics* **46**, 745 (1978), <http://dx.doi.org/10.1119/1.11393>.
- [2] G. Boole, *Philosophical Transactions of the Royal Society of London* **152**, 225 (1862).
- [3] I. Pitowsky, *The British Journal for the Philosophy of Science* **45**, 95 (1994), <http://dx.doi.org/10.1093/bjps/45.1.95>.
- [4] S. Kochen and E. P. Specker, *Journal of Mathematics and Mechanics* (now *Indiana University Math-*

- ematics Journal) **17**(1), 59 (1967), reprinted in [31, pp. 235–263], <http://dx.doi.org/10.1512/iumj.1968.17.17004>.
- [5] K. Svozil and J. Tkadlec, *Journal of Mathematical Physics* **37**(11), 5380 (November 1996), <http://dx.doi.org/10.1063/1.531710>.
- [6] A. Cabello, J. M. Estebarez, and G. García-Alcaine, *Physics Letters A* **212**(4), 183 (1996), [http://dx.doi.org/10.1016/0375-9601\(96\)00134-X](http://dx.doi.org/10.1016/0375-9601(96)00134-X).
- [7] D. M. Greenberger, M. A. Horne, and A. Zeilinger, in M. Kafatos, ed., *Bell's Theorem, Quantum Theory, and Conceptions of the Universe* (Kluwer Academic Publishers, Dordrecht, 1989), pp. 73–76.
- [8] N. D. Mermin, *Reviews of Modern Physics* **65**, 803 (1993), <http://dx.doi.org/10.1103/RevModPhys.65.803>.
- [9] N. Bohr, in P. A. Schilpp, ed., *Albert Einstein: Philosopher-Scientist* (The Library of Living Philosophers, Evanston, Ill., 1949), pp. 200–241, <http://www.emr.hibu.no/lars/eng/schilpp/Default.html>.
- [10] J. S. Bell, *Reviews of Modern Physics* **38**, 447 (1966), reprinted in [32, pp. 1-13], <http://dx.doi.org/10.1103/RevModPhys.38.447>.
- [11] P. Heywood and M. L. G. Redhead, *Foundations of Physics* **13**(5), 481 (1983), <http://dx.doi.org/10.1007/BF00729511>.
- [12] M. Redhead, *Incompleteness, Nonlocality, and Realism: A Prolegomenon to the Philosophy of Quantum Mechanics* (Clarendon Press, Oxford, 1990).
- [13] K. Svozil, in K. Engesser, D. M. Gabbay, and D. Lehmann, eds., *Handbook of Quantum Logic and Quantum Structures* (Elsevier, Amsterdam, 2008), pp. 551–586, arXiv:quant-ph/0609209, <http://arxiv.org/abs/quant-ph/0609209>.
- [14] K. Svozil, *Information Sciences* **179**, 535 (2009), <http://dx.doi.org/10.1016/j.ins.2008.06.012>.
- [15] A. Cabello and G. García-Alcaine, *Physical Review Letters* **80**(9), 1797 (1998), quant-ph/9709047, <http://dx.doi.org/10.1103/PhysRevLett.80.1797>.
- [16] M. Michler, H. Weinfurter, and M. Zukowski, *Physical Review Letters* **84**(24), 5457 (Jun 2000), quant-ph/0009061, <http://dx.doi.org/10.1103/PhysRevLett.84.5457>.
- [17] Y. Hasegawa, R. Loidl, G. Badurek, M. Baron, and H. Rauch, *Physical Review Letters* **97**(23), 230401, 230401 (4 pages) (2006), <http://dx.doi.org/10.1103/PhysRevLett.97.230401>.
- [18] A. Cabello, *Physical Review Letters* **101**(21), 210401, 210401 (4 pages) (2008), <http://dx.doi.org/10.1103/PhysRevLett.101.210401>.

org/10.1103/PhysRevLett.101.210401.

- [19] B. R. L. Cour, *Physical Review A (Atomic, Molecular, and Optical Physics)* **79**(1), 012102, 012102 (13 pages) (2009), <http://link.aps.org/abstract/PRA/v79/e012102>.
- [20] C. Simon, M. Żukowski, H. Weinfurter, and A. Zeilinger, *Physical Review Letters* **85**(9), 1783 (Aug 2000), quant-ph/0009074, <http://dx.doi.org/10.1103/PhysRevLett.85.1783>.
- [21] Y.-F. Huang, C.-F. Li, Y.-S. Zhang, J.-W. Pan, and G.-C. Guo, *Physical Review Letters* **90**(25), 250401 (Jun 2003), quant-ph/0209038, <http://dx.doi.org/10.1103/PhysRevLett.90.250401>.
- [22] G. Kirchmair, F. Zähringer, R. Gerritsma, M. Kleinmann, O. Ghne, A. Cabello, R. Blatt, and C. F. Roos, *Nature* **460**, 494 (2009), arXiv:0904.1655, <http://dx.doi.org/10.1038/nature08172>.
- [23] H. Bartosik, J. Klepp, C. Schmitzer, S. Sponar, A. Cabello, H. Rauch, and Y. Hasegawa, *Experimental test of quantum contextuality in neutron interferometry* (2009), eprint arXiv:0904.4576, arXiv:0904.4576.
- [24] P. R. Halmos, *Finite-dimensional vector spaces* (Springer, New York, Heidelberg, Berlin, 1974).
- [25] A. Stairs, *Philosophy of Science* **50**, 578 (1983), <http://www.jstor.org/stable/187557>.
- [26] H. R. Brown and G. Svetlichny, *Foundations of Physics* **20**, 1379 (1990), <http://dx.doi.org/10.1007/BF01883492>.
- [27] L. I. Schiff, *Quantum Mechanics* (McGraw-Hill, New York, 1955).
- [28] M. E. Rose, *Elementary Theory of Angular Momentum* (John Wiley & Sons, New York, 1957).
- [29] K. Svozil, *New Journal of Physics* **8**, 39 (2006), quant-ph/0401113, <http://dx.doi.org/10.1088/1367-2630/8/3/039>.
- [30] E. T. Jaynes, *Probability Theory: The Logic Of Science* (Cambridge University Press, Cambridge, 2003), <http://bayes.wustl.edu/etj/prob/book.pdf>.
- [31] E. Specker, *Selecta* (Birkhäuser Verlag, Basel, 1990).
- [32] J. S. Bell, *Speakable and Unspeakable in Quantum Mechanics* (Cambridge University Press, Cambridge, 1987).