


Bijections autour des bois de Schnyder

Éric Fusy (LIX, École Polytechnique)

travaux avec Olivier Bernardi, Dominique Poulalhon et Gilles Schaeffer

Schnyder structures on simple triangulations


Let T be a **simple triangulation** (topological, **up to isotopy**)



Schnyder structures on simple triangulations


1) Schnyder labellings

[Schnyder'89]




T can be endowed with a **labelling** of the corners by $\{1, 2, 3\}$ such that


inner faces



inner vertices




outer vertices



Schnyder structures on simple triangulations


1) Schnyder labellings

[Schnyder'89]




T can be endowed with a **labelling** of the corners by $\{\color{red}{\times}, \color{blue}{\times}, \color{green}{\times}\}$ such that


inner faces



inner vertices



outer vertices




Schnyder structures on simple triangulations


2) Schnyder woods

[Schnyder'89]


T can be endowed with a **tricoloration** + **orientation** of the inner edges such that



inner vertices



outer vertices




Schnyder structures on simple triangulations


2) Schnyder woods

[Schnyder'89]


T can be endowed with a **tricoloration** + **orientation** of the inner edges such that



inner vertices



outer vertices




Schnyder woods \leftrightarrow Schnyder labellings

Schnyder structures on simple triangulations


2) Schnyder woods

[Schnyder'89]


T can be endowed with a **tricoloration** + **orientation** of the inner edges such that



inner vertices



outer vertices




Schnyder structures on simple triangulations


2) Schnyder woods

[Schnyder'89]


T can be endowed with a **tricoloration** + **orientation** of the inner edges such that






inner vertices



outer vertices



yields a **spanning tree** in each color




Schnyder structures on simple triangulations


2) Schnyder woods

[Schnyder'89]


T can be endowed with a **tricoloration** + **orientation** of the inner edges such that



inner vertices



outer vertices




Schnyder structures on simple triangulations


3) 3-orientations

[Schnyder'89]

T can be endowed with an **orientation** of its inner edges such that






inner vertices



outdeg=3

outer vertices




outdeg=0


Schnyder structures on simple triangulations

The **3 incarnations** of Schnyder structures:


Schnyder labelling



Schnyder wood




3-orientation




Schnyder structures on simple triangulations

The **3 incarnations** of Schnyder structures:






Schnyder labelling



Schnyder wood




3-orientation



Applications of Schnyder woods

[Schnyder'89,90]

Associate **3 coordinates** to each vertex of T
(mapping from V to \mathbb{R}^3)



Applications of Schnyder woods


[Schnyder'89,90]

Associate **3 coordinates** to each vertex of T
(mapping from V to \mathbb{R}^3)

2 faces in blue area

3 faces in green area


$A \rightarrow (4, 2, 3)$



Applications of Schnyder woods

[Schnyder'89,90]

Associate **3 coordinates** to each vertex of T
(mapping from V to \mathbb{R}^3)




a_1	\rightarrow	$(9, 0, 0)$
a_2	\rightarrow	$(0, 9, 0)$
a_3	\rightarrow	$(0, 0, 9)$
A	\rightarrow	$(4, 2, 3)$
B	\rightarrow	$(5, 3, 1)$
C	\rightarrow	$(1, 4, 4)$
D	\rightarrow	$(2, 1, 6)$

all in $x + y + z = 9$

Applications of Schnyder woods

[Schnyder '89,90]


Associate **3 coordinates** to each vertex of T
(mapping from V to \mathbf{R}^3)



a_1	\rightarrow	$(9, 0, 0)$
a_2	\rightarrow	$(0, 9, 0)$
a_3	\rightarrow	$(0, 0, 9)$
A	\rightarrow	$(4, 2, 3)$
B	\rightarrow	$(5, 3, 1)$
C	\rightarrow	$(1, 4, 4)$
D	\rightarrow	$(2, 1, 6)$

all in $x + y + z = 9$

Straight-line drawing algo



Planarity criterion

$G = (V, E)$ is planar iff

$$\exists \Phi : V \cup E \rightarrow \mathbf{R}^3$$


such that

$$\forall p \neq q \in (V \cup E)^2$$

$$\Phi(p) \leq_{\mathbf{R}^3} \Phi(q)$$




$$p \in V, q \in E \text{ and } p \in q$$



Bijection for Schnyder woods


[Bonichon'02]
revisited in the dual setting



Bijection for Schnyder woods

[Bonichon'02]

Take the (3-regular) **dual** of the triangulation




Bijection for Schnyder woods

[Bonichon'02]

In black the **dual tree** of the red tree


In orange the **dual of the red edges**



Bijection for Schnyder woods

[Bonichon'02]


move corner-labels toward black vertices



Bijection for Schnyder woods

[Bonichon'02]


Erase the triangulation, keep the dual



Bijection for Schnyder woods

[Bonichon'02]

Cut the orange edges at their middle




Bijection for Schnyder woods

[Bonichon'02]

Cut the orange edges at their middle

⇒ **binary tree** such that there is a
parenthesis **matching of the leaves**





Bijection for Schnyder woods

[Bonichon'02]

binary tree such that there is a parenthesis **matching of the leaves**

rectilinear representation





Bijection for Schnyder woods

[Bonichon'02]

binary tree such that there is a parenthesis **matching of the leaves**

rectilinear representation





Bijection for Schnyder woods

[Bonichon'02]

binary tree such that there is a parenthesis **matching of the leaves**

rectilinear representation
(encoded by **two words**)





Bijection for Schnyder woods

[Bonichon'02]

binary tree such that there is a parenthesis **matching of the leaves**

rectilinear representation
(encoded by **two words**)





Bijection for Schnyder woods

[Bonichon'02]

binary tree such that there is a parenthesis **matching of the leaves**

rectilinear representation
(encoded by **two words**)




Bijection for Schnyder woods


[Bonichon'02]

binary tree such that there is a parenthesis **matching of the leaves**

rectilinear representation
(encoded by **two words**)



\Rightarrow






Bijection for Schnyder woods

[Bonichon'02]

binary tree such that there is a parenthesis **matching of the leaves**

rectilinear representation
(encoded by **two words**)




non-crossing pair of Dyck paths

Bijection for Schnyder woods [Bonichon'02]


[Bonichon'02]

binary tree such that there is a parenthesis **matching of the leaves**


rectilinear representation (encoded by **two words**)



⇒



⇐



non-crossing pair of Dyck paths

Total number s_n of Schnyder woods over triangulations with $n + 3$ vertices is


$$s_n = \text{Cat}_n \text{Cat}_{n+2} - \text{Cat}_{n+1} \text{Cat}_{n+1}$$

$$= \frac{6(2n)!(2n+2)!}{n!(n+1)!(n+2)!(n+3)!}$$

Lattice property for Schnyder woods

[Ossona de Mendez'94], [Brehm'00]


Theorem: Let T be a simple triangulation. Then the set of Schnyder structures of T is a **distributive lattice**





Lattice property for Schnyder woods

[Ossona de Mendez'94], [Brehm'00]

Theorem: Let T be a simple triangulation. Then the set of Schnyder structures of T is a **distributive lattice**




on Schnyder woods





Lattice property for Schnyder woods

[Ossona de Mendez'94], [Brehm'00]

Theorem: Let T be a simple triangulation. Then the set of Schnyder structures of T is a **distributive lattice**




on 3-orientations




Lattice property for Schnyder woods

[Ossona de Mendez'94], [Brehm'00]





Theorem: Let T be a simple triangulation. Then the set of Schnyder structures of T is a **distributive lattice**







on 3-orientations



min



max




The min is the **unique 3-orientation** of T with **no clockwise circuit**

Orientations and mobiles

Let \mathcal{O} be the set of **orientations** on planar maps such that:


- there is **no clockwise circuit**
- Each inner vertex can **access** the outer (unoriented simple) cycle
- the outer cycle is a **sink**




Orientations and mobiles

Let \mathcal{O} be the set of **orientations** on planar maps such that:

- there is **no clockwise circuit**
- Each inner vertex can **access** the outer (unoriented simple) cycle
- the outer cycle is a **sink**





Let \mathcal{M} be the set of **mobiles**, i.e., bipartite plane trees with **arrows** (called buds) at **black vertices**



“Master bijection”

[Bernardi, F'10]



Theorem: The above construction Φ is a **bijection** between \mathcal{O} and \mathcal{M} .
Moreover,



degrees of inner faces	\longleftrightarrow	degrees of black vertices
outdegrees of inner vertices	\longleftrightarrow	degrees of white vertices

Specialization to simple triangulations

- From the lattice property (**taking the min**) we have



family of simple triangulations \leftrightarrow subfamily \mathcal{F} of \mathcal{O} where:

- faces have degree 3
- inner vertices have outdegree 3






Specialization to simple triangulations

- From the lattice property (**taking the min**) we have
family of simple triangulations \leftrightarrow subfamily \mathcal{F} of \mathcal{O} where:
 - faces have degree 3
 - inner vertices have outdegree 3






- From the **master bijection specialized to \mathcal{F}** , we have
 $\mathcal{F} \leftrightarrow$ subfamily of mobiles where all vertices have **degree 3**






Counting formula

The **bijection** when there is a **marked inner face**:




Counting formula


The **bijection** when there is a **marked inner face**:




Each of the **3 parts** (when non empty) is **of the form**



\Rightarrow




\Rightarrow






quaternary tree

Counting formula

The **bijection** when there is a **marked inner face**:



Each of the **3 parts** (when non empty) is **of the form**




quaternary tree

Let $t_n = \#\{(\text{rooted}) \text{ triang. with } n + 3 \text{ vertices}\}$, $F(x) = \sum_n t_n x^{2n+1}$




Then $F'(x) = (1 + u)^3$ where $u = u(x)$ is specified by $u = \underbrace{x^2(1 + u)^4}_{\text{quat. trees}}$

Counting formula

The **bijection** when there is a **marked inner face**:



Each of the **3 parts** (when non empty) is **of the form**



quaternary tree

Let $t_n = \#\{(\text{rooted}) \text{ triang. with } n + 3 \text{ vertices}\}$, $F(x) = \sum_n t_n x^{2n+1}$

Then $F'(x) = (1 + u)^3$ where $u = u(x)$ is specified by $u = \underbrace{x^2(1 + u)^4}_{\text{quat. trees}}$


\Rightarrow
(Lagrange)

$$t_n = \frac{2(4n + 1)!}{(n + 1)!(3n + 2)!}$$

[Tutte'62]


Colored formulation of the bijection

Take the **Schnyder labelling** corresponding to the **minimal 3-orientation**




Colored formulation of the bijection

Take the **Schnyder labelling** corresponding to the **minimal 3-orientation**






Colored formulation of the bijection

Replace each




by






Colored formulation of the bijection


Replace each



by





Local rules:






Colored formulation of the bijection

- **Apply**




to each **inner white vertex**

- **Erase** the 3 **outer vertices** and their incident half-edges




Colored formulation of the bijection

• **Apply**




to each **inner white vertex**

• **Erase** the 3 **outer vertices** and their incident half-edges



Same bijection as before, because




Summary and extensions

- We have **two formulations of a bijection** for (simple) triangulations


Ⓐ

oriented




Ⓑ

colored



Summary and extensions

- We have **two formulations of a bijection** for (simple) triangulations



Let $t_n = \# \{(\text{rooted}) \text{ triang. with } n + 3 \text{ vertices}\}$, $F(x) = \sum_n t_n x^{2n+1}$



- Yields the **counting formulas** (one for GF, one for coefficients):

① $F'(x) = (1 + u)^3$ where $u = x^2(1 + u)^4$

② $t_n = \frac{2(4n+1)!}{(n+1)!(3n+2)!}$

Summary and extensions

- We have **two formulations of a bijection** for (simple) triangulations



Let $t_n = \#\{(\text{rooted}) \text{ triang. with } n + 3 \text{ vertices}\}$, $F(x) = \sum_n t_n x^{2n+1}$


- Yields the **counting formulas** (one for GF, one for coefficients):

(1) $F'(x) = (1 + u)^3$ where $u = x^2(1 + u)^4$

(2) $t_n = \frac{2(4n+1)!}{(n+1)!(3n+2)!}$

- We now give **two extensions**:


3-connected maps



Bijection extends **(B)**

Counting: (bivariate) extends **(2)**

d -angulations of girth d




Bijection extends **(A)**

Counting: (bivariate) extends **(1)**


Extension to 3-connected maps

3-connectivity

3-connected graph = needs **delete** at least 3 vertices to disconnect it




not 3-connected




3-connected

3-connectivity

3-connected graph = needs **delete** at least 3 vertices to disconnect it




not 3-connected




3-connected

3-connectivity

3-connected graph = needs **delete** at least 3 vertices to disconnect it





not 3-connected




3-connected

Rk: a triangulation is **3-connected** iff it is **simple**




3-connectivity

3-connected graph = needs **delete** at least 3 vertices to disconnect it





not 3-connected




3-connected

Rk: a triangulation is **3-connected** iff it is **simple**




Quasi 3-connected maps

A planar map G with 3 marked outer vertices $\{R, B, G\}$ is called **quasi 3-connected** if $G + \text{triangle formed by } \{R, B, G\}$ is 3-connected



Quasi 3-connected maps


A planar map G with 3 marked outer vertices $\{R, B, G\}$ is called **quasi 3-connected** if $G + \text{triangle formed by } \{R, B, G\}$ is 3-connected



not 3-connected


Quasi 3-connected maps

A planar map G with 3 marked outer vertices $\{R, B, G\}$ is called **quasi 3-connected** if $G + \text{triangle formed by } \{R, B, G\}$ is 3-connected



Quasi 3-connected maps


A planar map G with 3 marked outer vertices $\{R, B, G\}$ is called **quasi 3-connected** if $G + \text{triangle formed by } \{R, B, G\}$ is 3-connected



Let $\mathcal{Q}_{i,j}$ = set quasi 3-conn. maps with $i + 3$ vertices and j inner faces

Quasi 3-connected maps

A planar map G with 3 marked outer vertices $\{R, B, G\}$ is called **quasi 3-connected** if $G + \text{triangle formed by } \{R, B, G\}$ is 3-connected






Let $\mathcal{Q}_{i,j}$ = set quasi 3-conn. maps with $i + 3$ vertices and j inner faces

Rk: Extremal case $j = 2i + 1$ gives **triangulations** with $i + 3$ vertices

Duality for quasi 3-connected maps

The family of quasi 3-connected maps is **stable by duality**




$$Q_{i,j}^* = Q_{j,i}$$

Duality seen with the corner-map


Corner-map: obtained by **replacing each face by a star** (3 outer faces)

G



Duality seen with the corner-map


Corner-map: obtained by **replacing each face by a star** (3 outer faces)



Duality seen with the corner-map

Corner-map: obtained by **replacing each face by a star** (3 outer faces)

C




C is a dissection of an hexagon by **quadrangular faces**

Rk: quasi **3-connectivity** of $G \Leftrightarrow$ each **4-cycle** of C delimits a **face**

Duality seen with the corner-map

Corner-map: obtained by **replacing each face by a star** (3 outer faces)




G and G^* have the **same corner-map**


3-connected Schnyder labellings

Let G be a quasi 3-connected map. [Miller'02], [Felsner'04]


G can be endowed with a **labelling** of the **corners** by $\{\times, \times, \times\}$ such that




inner faces




inner vertices



outer vertices




outer face(s)




3-connected Schnyder labellings

Let G be a quasi 3-connected map. [Miller'02], [Felsner'04]


G can be endowed with a **labelling** of the **corners** by $\{\times, \times, \times\}$ such that




inner faces




inner vertices



outer vertices




outer face(s)




Rk: also incarnations as **Schnyder woods**, **3-orientations** (omitted)


Duality for 3-connected Schnyder labellings




Duality for 3-connected Schnyder labellings



Duality for 3-connected Schnyder labellings





Duality for 3-connected Schnyder labellings




Duality for 3-connected Schnyder labellings

duality is well seen on **corner map** C



Local rule:





same rule at black & white vertices

Lattice property in the 3-connected case

[Felsner'04] formulated on the associated **corner map** C




Theorem: Let G be a quasi 3-connected map. Then the set of Schnyder labellings of G is a **distributive lattice**





Lattice property in the 3-connected case

[Felsner'04] formulated on the associated **corner map** C

Theorem: Let G be a quasi 3-connected map. Then the set of Schnyder labellings of G is a **distributive lattice**






Rk: **extends** flip for triangulations





Lattice property in the 3-connected case

[Felsner'04] formulated on the associated **corner map** C

Theorem: Let G be a quasi 3-connected map. Then the set of Schnyder labellings of G is a **distributive lattice**





Rk: **extends** flip for triangulations




Bijection for quasi 3-connected maps

[F, Poulalhon, Schaeffer'08]




$\text{min}_{\text{lattice}}$

\approx




Bijection for quasi 3-connected maps

[F, Poulalhon, Schaeffer'08]




Bijection for quasi 3-connected maps

[F, Poulalhon, Schaeffer'08]




Bijection for quasi 3-connected maps

[F, Poulalhon, Schaeffer'08]






$i + 3$ vertices
 j inner faces





$\text{min}_{\text{lattice}}$

\approx




i black vert.
 j white vert.





unrooted **binary tree**
+ vertex bicoloration
+ edge tricoloration

Counting formula




G with a marked inner face





3 rooted binary trees

Counting formula



G with a marked inner face




3 rooted binary trees

Let $q_{i,j} = \#\{\text{quasi 3-conn. maps with } i + 3 \text{ vertices and } j \text{ inner faces}\}$



Let $F(x_{\circ}, x_{\bullet}) = \sum_{i,j} q_{i,j} x_{\circ}^i x_{\bullet}^j$

$$\frac{\partial}{\partial x_{\bullet}} F(x_{\circ}, x_{\bullet}) = (1 + U)^3, \text{ where } \begin{cases} U &= x_{\circ} \cdot (1 + V)^2, \\ V &= x_{\bullet} \cdot (1 + U)^2 \end{cases}$$

Counting formula



G with a marked inner face



3 rooted binary trees

Let $q_{i,j} = \#\{\text{quasi 3-conn. maps with } i + 3 \text{ vertices and } j \text{ inner faces}\}$

Let $F(x_{\circ}, x_{\bullet}) = \sum_{i,j} q_{i,j} x_{\circ}^i x_{\bullet}^j$

$$\frac{\partial}{\partial x_{\bullet}} F(x_{\circ}, x_{\bullet}) = (1 + U)^3, \text{ where } \begin{cases} U &= x_{\circ} \cdot (1 + V)^2, \\ V &= x_{\bullet} \cdot (1 + U)^2 \end{cases}$$


\Rightarrow Lagrange $q_{i,j} = \frac{3}{(2i+1)(2j+1)} \binom{2i+1}{j} \binom{2j+1}{i}$ [Mullin & Schellenberg'68]

recover triangulations counting formula in the (extremal) case $j = 2i + 1$

Extension to d -angulations of girth d

The girth parameter

The **girth** of a graph is the length of a shortest cycle within the graph



Girth = 3


Rk: Simple \Leftrightarrow girth ≥ 3

If **girth** = d then all faces have **degree at least d**

(in particular a triangulation is simple iff it has girth 3)

d -angulations of girth d


For $d \geq 3$ we consider d -angulations (all faces have degree d) of **girth d**



a pentagulation of girth 5

d -angulations of girth d

For $d \geq 3$ we consider d -angulations (all faces have degree d) of **girth d**



a pentagulation of girth 5


Rk: By the Euler relation,

$$\frac{\#(\text{inner edges})}{\#(\text{inner vertices})} = \frac{d}{d-2}$$

$d/(d-2)$ -orientations for d -angulations of girth d

[Bernardi-F'10]: Let G be a d -angulation of girth d . Then $(d-2)G$ admits an orientation where each inner vertex has **outdegree** d

Such an orientation is called a $d/(d-2)$ -orientation





$d/(d-2)$ -orientations for d -angulations of girth d

[Bernardi-F'10]: Let G be a d -angulation of girth d . Then $(d-2)G$ admits an orientation where each inner vertex has **outdegree** d

Such an orientation is called a $d/(d-2)$ -**orientation**

\Leftrightarrow **assignment** of (outgoing) **flows** to half-edges





total flow at inner edge = $d - 2$
total flow at inner vertex = d

$d/(d-2)$ -orientations for d -angulations of girth d

[Bernardi-F'10]: Let G be a d -angulation of girth d . Then $(d-2)G$ admits an orientation where each inner vertex has **outdegree** d

Such an orientation is called a $d/(d-2)$ -**orientation**

\Leftrightarrow **assignment** of (outgoing) **flows** to half-edges







Rk: also formulations as Schnyder labellings/woods

total flow at inner edge = $d - 2$
total flow at inner vertex = d



Lattice property for $d/(d-2)$ -orientations

The set of $d/(d-2)$ -orientations of a fixed d -angulation of girth d is a **distributive lattice**




Lattice property for $d/(d-2)$ -orientations


The set of $d/(d-2)$ -orientations of a fixed d -angulation of girth d is a **distributive lattice**





unique one with no "clockwise circuit"



- flow:
- 0 - - - - -
 - 1 —————
 - 2 = = = = =
 - 3 **—————**




Master bijection in the flow-formulation



degrees of inner faces	↔	degrees of black vertices
total flows at inner vertices	↔	total weights at white vertices
total flows at inner edges	↔	total weights at edges

Specialization to d -angulations of girth d



Bijection d -angulations of girth $d \leftrightarrow$ weighted mobiles such that

- each black vertex has degree d
- each white vertex has total weight d
- each edge has total weight $d - 2$ (weight > 0 at \circ , weight $= 0$ at \bullet)



[Albenque, Poulalhon'11]: other bijection (with blossoming trees)

Generating function expression

For $i \in [0..d]$, $\mathcal{L}_i :=$ family of such mobiles with a **root-leg of weight i**
 Let $L_i(x)$ be the GF of \mathcal{L}_i where x marks black nodes

Examples:

$d = 5$



For $d \geq 3$, $F_d(x) :=$ GF of (rooted) d -angulations of girth d by inner faces

- **Bijection** when an **inner face** is **marked**

$$\Rightarrow F'(x) = (1 + L_{d-2})^d$$

- **Root-decomposition** of mobiles in $\mathcal{L}_i \Rightarrow (L_0, L_1, \dots, L_d)$ are given by

$$\begin{cases} L_0 &= x \cdot (1 + L_{d-2})^{d-1}, \\ L_d &= 1, \\ L_i &= \sum_{j>0} L_{d-2-j} L_{i+j} \quad \text{for } i = 1..d-1 \end{cases}$$