

## EXERCICES, MPRI 2-38-1

return by Tuesday October 2nd, 2018

### 1. INTERVAL GRAPHS

We consider a finite set  $V$  and define  $\binom{V}{2} := \{\{u, v\} \mid u \neq v \in V\}$ . Consider a set  $\mathcal{I} := \{I_v \mid v \in V\}$  where  $I_v := [x_v, y_v]$  is an interval of  $\mathbb{R}$ . The *interval graph* of  $\mathcal{I}$  is the graph  $G_{\mathcal{I}}$  with vertex set  $V$  and edge set  $\{\{u, v\} \in \binom{V}{2} \mid I_u \cap I_v \neq \emptyset\}$ .

**Q 1.** What is the interval graph of  $\{[1, 4], [2, 6], [3, 8], [5, 9], [7, 10]\}$ ? Give a set of intervals with interval graph  $G = (V, E)$  where  $V = \{a, b, c, d, e\}$  and  $E = \{ab, ac, bc, cd, ce, de\}$ .

**Q 2.** Consider an interval graph  $G_{\mathcal{I}} = (V, E)$ . Show that:

- all induced cycles in  $G_{\mathcal{I}}$  are triangles,
- there is a partial order  $\prec$  on  $V$  whose comparability graph is the complement of  $G_{\mathcal{I}}$ , *i.e.* such that  $\{u, v\}$  is an edge in  $G_{\mathcal{I}}$  if and only if  $u$  and  $v$  are incomparable in  $\prec$ .

In fact, this is a characterization of interval graphs, but we skip the proof here.

### 2. BOXICITY

Consider a set  $\mathcal{B} := \{B_v \mid v \in V\}$  where  $B_v := [x_v^1, y_v^1] \times \dots \times [x_v^d, y_v^d]$  is a *box* in  $\mathbb{R}^d$  for some  $d \geq 1$ . The *box graph* of  $\mathcal{B}$  is the graph  $G_{\mathcal{B}}$  with vertex set  $V$  and edge set  $\{\{u, v\} \in \binom{V}{2} \mid B_u \cap B_v \neq \emptyset\}$ . See Figure 1 for an example. Given a graph on  $V$ , the *boxicity* of  $G$  is the smallest possible dimension  $d$  such that there exists a set  $\mathcal{B} = \{B_v \mid v \in V\}$  of boxes whose box graph  $G_{\mathcal{B}}$  is  $G$ .




FIGURE 1. A set  $\mathcal{B}$  of rectangles (2-dimensional boxes) and the corresponding box graph  $G_{\mathcal{B}}$ .

**Q 3.** What is the boxicity of a complete graph?

**Q 4.** Show that a cycle of length at least 4 has boxicity 2.

**Q 5.** Consider the intersection  $G \cap H = (V, E \cap F)$  of two graphs  $G = (V, E)$  and  $H = (V, F)$ . Show that the boxicity of  $G \cap H$  is at most the sum of the boxicities of  $G$  and  $H$ .

**Q 6.** What is the boxicity of an interval graph? Show that the boxicity of  $G = (V, E)$  is the minimal number  $d$  of interval graphs  $G_{\mathcal{I}^1} = (V, E^1), \dots, G_{\mathcal{I}^d} = (V, E^d)$  such that  $E = E^1 \cap \dots \cap E^d$ .

**Q 7.** Consider a graph  $G = (V, E)$  and an induced subgraph  $H = (U, E \cap \binom{U}{2})$  for some  $U \subseteq V$ . Show that the boxicity of  $H$  is at most the boxicity of  $G$ .

### 3. GENERAL UPPER BOUND

We now show an upper bound on the boxicity of any graph  $G$ .

**Q 8.** According to Q3, we can consider a graph  $G = (V, E)$  that is not complete. Let  $u, v$  be two non-adjacent vertices of  $G$  and let  $H = G \setminus \{u, v\}$  be the graph  $G$  where  $u$  and  $v$  were deleted. Assume that  $H$  is the intersection of  $d$  interval graphs  $G_{\mathcal{I}^1}, \dots, G_{\mathcal{I}^d}$ . Define  $d + 1$  sets of intervals  $\mathcal{I}^1, \dots, \mathcal{I}^{d+1}$  as follows:

- For all  $i \in [d]$ , let  $\mathcal{I}^i := \mathcal{J}^i \cup \{J_u, J_v\}$  where  $J_u$  and  $J_v$  are intervals that are large enough to intersect all intervals in  $\mathcal{J}^1 \cup \dots \cup \mathcal{J}^d$ .
- Let  $\mathcal{I}^{d+1} := \{I_w \mid w \in V\}$  where

$$I_w := \begin{cases} \{-1\} & \text{if } w = u \\ \{1\} & \text{if } w = v \\ \{0\} & \text{if } (u, w) \notin E \text{ and } (v, w) \notin E \\ [-1, 0] & \text{if } (u, w) \in E \text{ but } (v, w) \notin E \\ [0, 1] & \text{if } (u, w) \notin E \text{ but } (v, w) \in E \\ [-1, 1] & \text{if } (u, w) \in E \text{ and } (v, w) \in E. \end{cases}$$

Show that  $G$  is the intersection of the interval graphs  $G_{\mathcal{I}^1}, \dots, G_{\mathcal{I}^{d+1}}$ .

**Q 9.** Deduce from the previous question that a graph on  $n$  vertices has boxicity at most  $n/2$ .

**Q 10.** Consider the graph  $U_p$  on  $p = 2q$  vertices obtained by deleting a perfect matching  $M$  from the complete graph  $K_{2q}$ . Show that the boxicity of  $U_p$  is at least  $q = p/2$ . (Hint: Assume that  $U_p$  is the intersection of  $d$  interval graphs  $G_{\mathcal{I}^1}, \dots, G_{\mathcal{I}^d}$ . Show that each edge of the matching  $M$  is missing in at least one of the interval graphs  $G_{\mathcal{I}^k}$  and that two edges of the matching  $M$  cannot be missing in the same interval graph  $G_{\mathcal{I}^k}$ .)

#### 4. SCHNYDER WOODS AND BOXICITY

Thomassen proved that planar graphs have boxicity at most 3. The goal of this section is to prove this result using Schnyder woods.

**Q 11.** Show that any planar graph is an induced subgraph of a triangulation. Deduce from Q 7 that the special case of triangulations suffices to prove Thomassen's result.

Consider now a triangulation  $T = (V, E)$  endowed with a Schnyder wood  $(T^1, T^2, T^3)$ . In other words,  $T^1, T^2, T^3$  are three spanning trees of  $T$ , which partition the edges of  $T$  (except the edges of the outer face which are all contained in two of these trees), and which fulfill Schnyder's local conditions around each vertex. Note that in contrast to the general case seen in the course, only the edges of the external face are bioriented since  $T$  is a triangulation. Consider a vertex  $v \in V$ . We denote by  $R^i(v)$  the region of  $T$  bounded by the paths from  $v$  to the root of the trees  $T^{i-1}$  and  $T^{i+1}$ , and we let  $r^i(v) = |R^i(v)|$ . We define  $x_v^i := r^i(v)$  and  $y_v^i := r^i(v^i)$ , where  $v^i$  is the parent of  $v$  in the tree  $T^i$ . Note that when  $v$  is the root of  $T^i$ , the vertex  $v^i$  is not defined, but we let  $y_v^i := r^i(v) + 1$ . Consider the box  $B_v := [x_v^1, y_v^1] \times [x_v^2, y_v^2] \times [x_v^3, y_v^3]$ . Finally, let  $\mathcal{B} := \{B_v \mid v \in V\}$ . We have represented an example in Figure 2.




FIGURE 2. A triangulation with a Schnyder wood (left), the corresponding orthogonal surface (middle), and the corresponding box representation (right). The three external boxes have been reduced to let the other ones apparent.




FIGURE 3. The octahedral triangulation (left), and some framework to draw box representations corresponding to two Schnyder woods on  $O$  (middle and right).

**Q 12.** Consider the triangulation  $O$  of Figure 3 (left) (note that it is the graph of an octahedron). Compute all possible Schnyder woods on  $O$  (Hint: starting from one Schnyder wood, all the other are obtained by returning oriented cycles surrounding a face of  $O$ .) For each Schnyder wood of  $O$ , compute the boxes  $B_v$  for all vertices  $v$  of  $O$ . Finally, draw these boxes as in Figure 2. You can use the framework of Figure 3 (middle and right) to help your drawing (don't forget to insert this in your exam).

**Q 13.** Consider two adjacent vertices  $u, v \in V$ , and assume that  $u$  is a child of  $v$  in  $T^i$ . Show that

$$x_v^{i-1} \leq x_u^{i-1} \leq y_v^{i-1} \quad y_u^i = x_v^i \quad x_v^{i+1} \leq x_u^{i+1} \leq y_v^{i+1}$$

and conclude that  $\{u, v\} \in G_{\mathcal{B}}$ .

**Q 14.** Consider now two non-adjacent vertices  $u, v \in V$ , let  $i$  be such that  $u \in R_i(v)$  and let  $u^i$  be the parent of  $u$  in  $T^i$ . Show that  $y_{u^i}^i < x_v^i$  when

- $v$  lies on the path from  $u$  to the root of  $T^i$ , or
- $u$  does not lie on the paths from  $v$  to the root of the trees  $T^{i-1}$  and  $T^{i+1}$ .

Deduce that if  $u$  and  $v$  are non-adjacent vertices in  $T$ , then  $B_u \cap B_v = \emptyset$  so that  $\{u, v\} \notin G_{\mathcal{B}}$ .

**Q 15.** Conclude from the two previous questions that  $T = (V, E)$  is the box graph  $G_{\mathcal{B}}$  for the set of boxes  $\mathcal{B} := \{B_v \mid v \in V\}$ .

**Q 16.** We now consider the planar graph of Figure 4 (left) with two marked vertices  $u, v$ . Computing  $x^3(u)$  and  $y^3(v)$ , show that the recipe given for triangulations does not directly work for arbitrary 3-connected planar graphs. This issue is illustrated in Figure 4 where you can see that the boxes corresponding to  $u$  and  $v$  are disjoint, while  $u$  and  $v$  should be adjacent.




FIGURE 4. A planar graph with a Schnyder wood (left), the corresponding orthogonal surface (middle), and the corresponding incorrect box representation (right). The three external boxes have been reduced to let the other ones apparent.