

CELEBRATING LODAY'S ASSOCIAHEDRON

VINCENT PILAUD, FRANCISCO SANTOS, AND GÜNTER M. ZIEGLER

ABSTRACT. We survey Jean-Louis Loday's vertex description of the associahedron, and its far reaching influence in combinatorics, discrete geometry and algebra. We present in particular four topics where it plays a central role: lattice congruences of the weak order and their quotientopes, cluster algebras and their generalized associahedra, nested complexes and their nestohedra, and operads and the associahedron diagonal.

CONTENTS

Introduction	2
1. Permutahedra, associahedra, cubes	3
1.1. Lattices	3
1.2. Polytopes	4
1.3. Hopf algebras	6
1.4. Three surprising geometric properties	7
2. Lattice congruences and quotientopes	8
2.1. Sylvester congruence	8
2.2. Cambrian and permutoree congruences	8
2.3. All congruences	10
2.4. Beyond the braid arrangement	11
3. Cluster algebras, brick polytopes, and quiver representation theory	13
3.1. Cluster algebras and generalized associahedra	13
3.2. Subword complexes and brick polytopes	14
3.3. Quiver representation theory and gentle associahedra	18
4. Graph associahedra and nestohedra	19
4.1. Graph associahedra	19
4.2. Hypergraph associahedra and nestohedra	20
5. Operads and diagonals	21
5.1. Loop spaces and operads	21
5.2. Cellular diagonal of the associahedron	22
6. Further generalizations	23
Acknowledgements	24
References	24

VP was supported by the French ANR grant CHARMS (ANR-19-CE40-0017), and by the French–Austrian project PAGCAP (ANR-21-CE48-0020 & FWF I 5788).

FS was supported by grant PID2019-106188GB-I00 funded by MCIN/AEI/10.13039/501100011033 and by project CLAPPo (21.SI03.64658) of Universidad de Cantabria and Banco Santander.

GMZ is a member of The Berlin Mathematics Research Center MATH+, funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy (EXC-2046/1, project ID: 390685689).

INTRODUCTION

The associahedron is a polytope whose face lattice encodes Catalan families: its vertices correspond to parenthesizations of a non-associative product, triangulations of a convex polygon, or binary trees; its edges correspond to applications of the associativity rule, diagonal flips, or edge rotations; and in general its faces correspond to partial parenthesizations, diagonal dissections, or Schröder trees. It was defined combinatorially in early works of D. Tamari [Tam51] and J. Stasheff [Sta63] with motivation from associativity and loop spaces. The associahedron now appears as a fundamental structure throughout mathematics, in particular for moduli spaces and topology [Sta63, Sta97, Kel01], operads and rewriting theory [Sta04, LV12, Val14, Str12, MTTV21], cluster algebras [FZ03a, CFZ02, Rea06, HLT11, HPS18], quiver representation theory [BKT14, BMCLD⁺18, PPPP19, AHI⁺22], combinatorial Hopf algebras [LR98, CP17, PP18, Pil18], diagonal harmonics [BPR12, PRV17], physics of scattering amplitudes [AHBHY18], etc.

While some 3-dimensional associahedra were drawn in D. Tamari's PhD thesis [Tam51] and constructed by J. Milnor for the PhD defense of J. Stasheff, the first systematic polytopal realizations were constructed by M. Haiman [Hai84] and C. Lee [Lee89]. Since then, three families of realizations were largely developed: the secondary polytope realizations [GKZ08, BFS90], the \mathbf{g} -vectors realizations [RSS03, Lod04, HL07, Pos09, HLT11, PS12, PS15a, HPS18, LP18] and the \mathbf{d} -vector realizations [CFZ02, CSZ15, MP17]. See [CSZ15] for a discussion of some of these realizations, of their connections, and of their respective advantages.

On the occasion of the 75th anniversary of *Archiv der Mathematik* and the 20th anniversary of J.-L. Loday's paper [Lod04], we review in this paper the far-reaching influence of Loday's associahedron. It has to be mentioned that this same realization was already described in [SS93] by simple facet inequalities, one for each interval of $[n]$, but it became extremely useful and popular only when J.-L. Loday provided his elementary description of the vertices, one for each binary tree. Namely, each binary tree T corresponds to a vertex with coordinates $\ell(T, i) \cdot r(T, i)$, where $\ell(T, i)$ and $r(T, i)$ respectively denote the numbers of leaves in the left and right subtrees of the i th node of T (in infix labeling). The same realization of associahedron was later described in [Pos09] as the Minkowski sum of all faces of the standard simplex corresponding to intervals of $[n]$. See Section 1.

This realization has several advantages. Some were already underlined by J.-L. Loday in [Lod04]: “it admits simple vertex and facet descriptions, respects the symmetry, and fits with the classical realization of the permutohedron”. But we believe that the real reason that makes this realization ubiquitous in the literature is that its normal fan transparently encodes each binary tree by a very natural cone: the cone with one inequality $x_i \leq x_j$ for each edge $i \rightarrow j$ in the tree. This implies in particular that the natural surjective map from permutations to binary trees [Ton97] translates geometrically to fans and polytopes, and that the oriented graph of this associahedron is the Hasse diagram of the Tamari lattice [Tam51] (Section 1).

Loday's construction has served as a prototype for several constructions generalizing the associahedron. In this survey, we present four specific topics in which it was instrumental:

- quotientopes (Section 2),
- cluster algebras, subword complexes, and quiver representation theory (Section 3),
- graph associahedra and nestohedra (Section 4),
- operads and diagonals (Section 5).

Additional material on some of these topics can be found in other more detailed surveys: see in particular [CSZ15, CZ12] for realization spaces, [Rea12, Rea16a, Rea16b] for lattice quotients of the weak order, [FWZ] for cluster algebras, [FR07, Hoh12] for generalized associahedra, [Tho12, Tho21] for associahedra and Tamari lattices in representation theory, [Sta04, LV12, Val14, Gir18] for operad theory, and the book [MHPS12] for many more connections of the Tamari lattice.

To summarize, the simplicity of Loday's description of the associahedron has fundamentally contributed to break the psychological barrier of realizing this “mythical polytope” [Hai84]. It resulted in several constructions of polytopes with a combinatorial flavor similar to that of the associahedron, some of which are directly constructed from this associahedron (by Cartesian products and Minkowski sums, as sections or projections, or as polyhedral decompositions). We conclude this survey with a (incomplete and partial) list of these constructions (Section 6), and invite the reader to discover many more descendants of Loday's associahedron.

We hope that this survey can serve as an invitation to the “multiple facets of the associahedron” [Lod05]. Besides bibliographic pointers to the original literature, we have reproduced many pictures that should already give an idea of the topics covered in this survey. Much of the material of this survey is derived from the Habilitation thesis of V. Pilaud [Pil20], who should be considered the main contributor.

1. PERMUTAHEDRA, ASSOCIAHEDRA, CUBES

J.-L. Loday’s paper [Lod04] is a tale of three magical families (permutations, binary trees, and binary sequences) and their fantastic adventures in the lands of lattices, polytopes, and Hopf algebras. This section proposes a brief recollection of these structures, summarized by the table:

	permutations	binary trees	binary sequences
Lattices	Weak order	Tamari lattice [Tam51]	Boolean lattice
Polytopes	Permutahedron $\text{Perm}(n)$	Loday’s associahedron $\text{Asso}(n)$ [Lod04]	Parallelepiped $\text{Para}(n)$ generated by $e_{i+1} - e_i$
Hopf algebras	Malvenuto–Reutenauer Hopf algebra [MR95]	Loday–Ronco Hopf algebra [LR98]	Recoil Hopf algebra [GKL ⁺ 95]

1.1. Lattices. We start with three lattices illustrated in Figure 1:




FIGURE 1. Weak order on \mathfrak{S}_4 (left), Tamari lattice on \mathfrak{T}_4 (middle), boolean lattice on \mathfrak{B}_4 (right).

Weak order. We denote by \mathfrak{S}_n the set of permutations of $[n] := \{1, \dots, n\}$. The (right) weak order on \mathfrak{S}_n is the partial order \leq_w defined by inclusion of inversion sets (an inversion of $\sigma \in \mathfrak{S}_n$ is a pair of values $i, j \in [n]$ such that i is smaller than j but i appears after j in σ). Its cover relations are given by transpositions of two consecutive letters. Its minimal and maximal elements are the permutations $1 \dots n$ and $n \dots 1$, respectively. It was shown in [GR63, Bjö84] that the weak order is a lattice (*i.e.* minimal upper bounds and maximal lower bounds are unique).

Tamari lattice. We denote by \mathfrak{T}_n be the set of rooted binary trees with n internal nodes (equivalently, with $n+1$ leaves). The Tamari lattice on \mathfrak{T}_n is the lattice \leq_T whose cover relations are given by right rotations on binary trees. Its minimal and maximal elements are the left and right combs, respectively. We consider the n internal nodes of a binary tree labeled by $[n]$ in infix order: first the left subtree is labeled, then the root, then the right subtree, recursively in subtrees. Observe that this makes node $i+1$ be a descendant or an ancestor of i .

Boolean lattice. We denote by \mathfrak{B}_n the set of binary sequences of $n-1$ signs + or -. The boolean lattice on \mathfrak{B}_n is the lattice \leq_B defined by $\chi \leq_B \zeta$ if and only if $\chi_i \leq_B \zeta_i$ for all $i \in [n-1]$ (for $- \leq +$). Its cover relations are given by replacements of a - by a +. Its minimal and maximal elements are the sequences $-^{n-1}$ and $+^{n-1}$, respectively.

Remark 1. Observe that we denote by \mathfrak{B}_n the boolean lattice on $n-1$ elements, not n . We take this convention for their relation to permutations of n elements and binary trees with n internal nodes, where our use of n is standard. Our sets \mathfrak{S}_n , \mathfrak{T}_n , and \mathfrak{B}_n are respectively denoted by S_n , Y_n and Q_n in J.-L.-Loday’s paper [Lod04].

These three lattices are related by the following commutative diagram of lattice morphisms:

$$\begin{array}{ccccc} \mathfrak{S}_n & \xrightarrow{\text{rec}} & \mathfrak{B}_n \\ \text{bt} \searrow & & \swarrow \text{can} \\ & \mathfrak{T}_n & \end{array}$$

where

- the *binary tree map* (or *Tonks projection*) sends a permutation $\sigma := \sigma_1 \dots \sigma_n \in \mathfrak{S}_n$ to the binary tree $\text{bt}(\sigma) \in \mathfrak{T}_n$ obtained by successive insertions of $\sigma_n, \dots, \sigma_1$ in a binary search tree. That is, σ_n is the root, and its left and right subtrees are constructed recursively from the permutations obtained by restriction of σ to $\{i \mid i < \sigma_n\}$ and to $\{i \mid i > \sigma_n\}$, respectively. Equivalently, the **bt** fiber of a tree T is precisely the set of linear extensions of T , *i.e.* all permutations σ such that for any $i, j \in [n]$, if i is a descendant of j in T , then i appears before j in σ . This map was considered in [Ton97, Lod04, HNT05, Rea06].
- the *canopy map* (or *Loday–Ronco projection*) sends a binary tree $T \in \mathfrak{T}_n$ to the binary sequence $\text{can}(T) \in \mathfrak{B}_n$ where at position $i \in [n-1]$ there is a $-$ if i appears below $i+1$ and a $+$ if i appears above $i+1$ in T . This map was first used by J.-L. Loday and M. Ronco in [LR98], but the name “canopy” was coined by X. Viennot [Vie07].
- the *recoil map* is the composition of the previous two. It sends a permutation $\sigma \in \mathfrak{S}_n$ to the binary sequence $\text{rec}(\sigma) \in \mathfrak{B}_n$ where at position $i \in [n-1]$ there is a $-$ if i appears before $i+1$ and a $+$ if $i+1$ appears after i in σ .

Note that the cover graphs of these lattices are classical combinatorial graphs, namely the simple transposition graph on \mathfrak{S}_n , the rotation graph on \mathfrak{T}_n , and the bit change graph on \mathfrak{B}_n . Their combinatorial properties have been extensively investigated; in particular:

- they all admit Hamiltonian cycles. This is classical for \mathfrak{S}_n by the Steinhaus–Johnson–Trotter algorithm [Ste64, Joh63, Tro62], and for \mathfrak{B}_n by the seminal work of F. Gray [Knu11]. For \mathfrak{T}_n , it was shown in [Luc87] and later revisited in [HN99, HM21].
- their exact diameter is known. It is obviously $\binom{n}{2}$ for \mathfrak{S}_n and $n-1$ for \mathfrak{B}_n . For \mathfrak{T}_n , there is an immediate lower bound of n and, via the bijection between binary trees and triangulations of the $(n+2)$ -gon, it is relatively easy to show an upper bound of $2n-6 + \lfloor \frac{12}{n+2} \rfloor$ (see *e.g.* [DRS10, Proposition 1.1.5]). In particular, the diameter is at most $2n-6$ for all $n > 10$. Using volumetric arguments in hyperbolic geometry it was shown in [STT88] that the diameter indeed equals $2n-6$ for n sufficiently large, but it took another 25 years until a (purely combinatorial, yet sophisticated) proof for all $n > 10$ was given in [Pou14]. Along the way, a lower bound of $2n - 2\sqrt{2n}$ valid for all n was obtained in [Deh10] using Thompson’s groups.

1.2. Polytopes. We denote by $(e_i)_{i \in [n]}$ the canonical basis of \mathbb{R}^n and $\mathbf{1} := \sum_{i \in [n]} e_i$. All our polytopal constructions will lie in the affine subspace $\mathbb{H} := \{\mathbf{x} \in \mathbb{R}^n \mid \langle \mathbf{1} | \mathbf{x} \rangle = \sum_{i \in [n]} x_i = \binom{n+1}{2}\}$, and their normal fans will lie in the vector subspace $\mathbf{1}^\perp := \{\mathbf{x} \in \mathbb{R}^n \mid \langle \mathbf{1} | \mathbf{x} \rangle = 0\}$. The main objects of J.-L. Loday’s paper [Lod04] are the following three polytopes illustrated in Figure 2:




FIGURE 2. Permutahedron $\text{Perm}(4)$ (left), Loday’s associahedron $\text{Asso}(4)$ (middle), parallelepiped $\text{Para}(4)$ (right). Shaded facets are preserved to get the next polytope. [CP17, Fig. 1].

Permutahedron. The *permutohedron* $\text{Perm}(n)$ is the polytope in \mathbb{R}^n obtained equivalently as:

- the convex hull of the points $\sum_{i \in [n]} i \mathbf{e}_{\sigma(i)}$ for all permutations $\sigma \in \mathfrak{S}_n$, see [Sch11],
- the intersection of the hyperplane \mathbb{H} with the halfspaces $\{\mathbf{x} \in \mathbb{R}^n \mid \sum_{i \in I} x_i \geq \binom{|I|+1}{2}\}$ for all $\emptyset \neq I \subsetneq [n]$, see [Rad52],
- (a translate of) the Minkowski sum of all segments $[\mathbf{e}_i, \mathbf{e}_j]$ for all $(i, j) \in \binom{[n]}{2}$ (where the Minkowski sum of two polytopes $\mathbb{P}, \mathbb{Q} \subseteq \mathbb{R}^n$ is the polytope $\mathbb{P} + \mathbb{Q} := \{p + q \mid p \in \mathbb{P}, q \in \mathbb{Q}\}$).

The face lattice of the permutohedron $\text{Perm}(n)$ is the refinement lattice on ordered partitions of $[n]$. The normal fan of the permutohedron $\text{Perm}(n)$ is the *braid fan* $\mathcal{F}(n)$ defined by the (type A) Coxeter arrangement formed by the hyperplanes $\{\mathbf{x} \in \mathbf{1}^\perp \mid x_i = x_j\}$ for all $1 \leq i < j \leq n$. Namely, each permutation $\sigma \in \mathfrak{S}_n$ corresponds to the maximal cone $\mathbf{C}^\circ(\sigma) := \{\mathbf{x} \in \mathbf{1}^\perp \mid x_{\sigma(1)} \leq \dots \leq x_{\sigma(n)}\}$ of the braid fan $\mathcal{F}(n)$, consisting of all points whose coordinates are ordered by the permutation σ .

Associahedron. The *associahedron* $\text{Asso}(n)$ is the polytope in \mathbb{R}^n obtained equivalently as:

- the convex hull of the points $\sum_{i \in [n]} \ell(T, i) r(T, i) \mathbf{e}_i$ for all binary trees $T \in \mathfrak{T}_n$, where $\ell(T, i)$ and $r(T, i)$ respectively denote the numbers of leaves in the left and right subtrees of the i th node of T in infix labeling, see [Lod04],
- the intersection of the hyperplane \mathbb{H} with the halfspaces $\{\mathbf{x} \in \mathbb{R}^n \mid \sum_{i \leq \ell \leq j} x_\ell \geq \binom{j-i+2}{2}\}$ for all $1 \leq i \leq j \leq n$, see [SS93],
- (a translate of) the Minkowski sum of the faces $\Delta_{[i,j]}$ of the standard simplex $\Delta_{[n]}$ for all $1 \leq i \leq j \leq n$, where $\Delta_X := \text{conv}\{\mathbf{e}_x \mid x \in X\}$ for $X \subseteq [n]$, see [Pos09].

The face lattice of the associahedron $\text{Asso}(n)$ is the edge contraction lattice on Schröder trees with $n+1$ leaves (rooted plane trees where each node has at least 2 children). The normal fan of the associahedron is the *sylvester fan*. Each binary tree $T \in \mathfrak{T}_n$ corresponds to a normal cone $\mathbf{C}^\circ(T) := \{\mathbf{x} \in \mathbf{1}^\perp \mid x_i \leq x_j \text{ for } i \text{ child of } j \text{ in } T\}$.

Parallelepiped. Finally, we consider the *parallelepiped* $\text{Para}(n)$ in \mathbb{R}^n obtained equivalently as:

- the convex hull of the points $\frac{n+1}{2} \mathbf{1} + \frac{n-1}{2} \sum_{i \in [n-1]} \chi_i (\mathbf{e}_i - \mathbf{e}_{i+1})$ for all binary sequences $\chi \in \mathfrak{B}_n$,
- the intersection of the hyperplane \mathbb{H} with the halfspaces $\{\mathbf{x} \in \mathbb{R}^n \mid \sum_{1 \leq \ell \leq i} x_\ell \geq \binom{i+1}{2}\}$ and $\{\mathbf{x} \in \mathbb{R}^n \mid \sum_{i < \ell \leq n} x_\ell \geq \binom{n-i+1}{2}\}$ for all $i \in [n-1]$,
- (a translate of) the Minkowski sum of the segments $(n-1) \cdot [\mathbf{e}_i, \mathbf{e}_{i+1}]$ for $i \in [n-1]$.

The face lattice of the parallelepiped $\text{Para}(n)$ is the lattice on ternary words on $\{-, 0, +\}$ given by the componentwise order, where the order on $\{-, 0, +\}$ is defined by $0 \leq -$ and $0 \leq +$. The normal fan of the parallelepiped $\text{Para}(n)$ is the *coordinate fan*. Each binary sequence $\chi \in \mathfrak{B}_n$ corresponds to a normal cone $\mathbf{C}^\circ(\chi) := \{\mathbf{x} \in \mathbf{1}^\perp \mid \chi_i (x_i - x_{i+1}) \geq 0 \text{ for all } i \in [n-1]\}$.

Remark 2. Observe that we stick to our convention of Remark 1 and depart from Loday's notation; the polytopes $\text{Perm}(n)$, $\text{Asso}(n)$ and $\text{Para}(n)$ (which are of dimension $n-1$ but are naturally embedded in a hyperplane in \mathbb{R}^n) are denoted by \mathcal{P}^{n-1} , \mathcal{K}^{n-1} , and \mathcal{C}^{n-1} in [Lod04].

These three polytopes and fans refine each other:

- the facet description of $\text{Perm}(n)$ contains the facet description of $\text{Asso}(n)$, which contains the facet description of $\text{Para}(n)$ (see Section 1.4.1 for more details).
- the braid fan refines the sylvester fan, which refines the coordinate fan. Namely,

$$\mathbf{C}^\circ(T) = \bigcup_{\substack{\sigma \in \mathfrak{S}_n \\ \text{bt}(\sigma) = T}} \mathbf{C}^\circ(\sigma) \quad \text{and} \quad \mathbf{C}^\circ(\chi) = \bigcup_{\substack{\sigma \in \mathfrak{S}_n \\ \text{rec}(\sigma) = \chi}} \mathbf{C}^\circ(\sigma) = \bigcup_{\substack{T \in \mathfrak{T}_n \\ \text{can}(T) = \chi}} \mathbf{C}^\circ(T).$$

Moreover, these polytopes geometrically realize the lattices of Section 1.1. Indeed, oriented in the direction $\omega := (n, \dots, 1) - (1, \dots, n) = \sum_{i \in [n]} (n+1-2i) \mathbf{e}_i$, the graph of the permutohedron $\text{Perm}(n)$ (resp. of the associahedron $\text{Asso}(n)$, resp. of the parallelepiped $\text{Para}(n)$) is the Hasse diagram of the weak order on \mathfrak{S}_n (resp. of the Tamari lattice on \mathfrak{T}_n , resp. of the boolean lattice on \mathfrak{B}_n).

1.3. Hopf algebras. Recall that a combinatorial Hopf algebra is a combinatorial vector space \mathfrak{A} endowed with an associative product $\cdot : \mathfrak{A} \otimes \mathfrak{A} \rightarrow \mathfrak{A}$ and a coassociative coproduct $\Delta : \mathfrak{A} \rightarrow \mathfrak{A} \otimes \mathfrak{A}$, subject to the compatibility relation $\Delta(a \cdot b) = \Delta(a) \cdot \Delta(b)$, where the right hand side product has to be understood componentwise. As discussed in [Lod05], the construction of J.-L. Loday's paper [Lod04] was fundamentally motivated by the following three Hopf algebras:

Malvenuto–Reutenauer Hopf algebra. For $\rho \in \mathfrak{S}_m$ and $\sigma \in \mathfrak{S}_n$, the *shuffle* $\rho \sqcup \sigma$ (resp. the *convolution* $\rho \star \sigma$) denotes the set of permutations of \mathfrak{S}_{m+n} where the order of the first m and last n values (resp. positions) is given by ρ and σ respectively. For instance

$$\begin{aligned} 12 \sqcup 231 &= \{12453, 14253, 14523, 14532, 41253, 41523, 41532, 45123, 45132, 45312\}, \\ 12 \star 231 &= \{12453, 13452, 14352, 15342, 23451, 24351, 25341, 34251, 35241, 45231\}. \end{aligned}$$

Let $\mathfrak{S} := \bigsqcup_{n \in \mathbb{N}} \mathfrak{S}_n$ be the set of all finitary permutations (any size) and $\mathbf{k}\mathfrak{S}$ denote its \mathbf{k} -vector span with basis $(\mathbb{F}_\tau)_{\tau \in \mathfrak{S}}$. The *Malvenuto–Reutenauer Hopf algebra* [MR95] is the Hopf algebra on $\mathbf{k}\mathfrak{S}$ where the product \cdot and coproduct Δ are defined by

$$\mathbb{F}_\rho \cdot \mathbb{F}_\sigma = \sum_{\tau \in \rho \sqcup \sigma} \mathbb{F}_\tau \quad \text{and} \quad \Delta \mathbb{F}_\tau = \sum_{\tau \in \rho \star \sigma} \mathbb{F}_\rho \otimes \mathbb{F}_\sigma.$$

Loday–Ronco Hopf algebra. The *Loday–Ronco Hopf algebra* [LR98] is the Hopf subalgebra $\mathbf{k}\mathfrak{T}$ of $\mathbf{k}\mathfrak{S}$ generated by the elements

$$\mathbb{P}_T := \sum_{\substack{\tau \in \mathfrak{S} \\ \mathbf{bt}(\tau) = T}} \mathbb{F}_\tau,$$

for all binary trees $T \in \mathfrak{T}$ of any size.

Recoil Hopf algebra. The *recoil Hopf algebra* [GKL⁺95] is the Hopf subalgebra $\mathbf{k}\mathfrak{B}$ of $\mathbf{k}\mathfrak{S}$ generated by the elements

$$\mathbb{X}_\chi := \sum_{\substack{\tau \in \mathfrak{S} \\ \mathbf{rec}(\tau) = \chi}} \mathbb{F}_\tau = \sum_{\substack{T \in \mathfrak{T} \\ \mathbf{can}(T) = \chi}} \mathbb{P}_T,$$

for all binary sequences $\chi \in \mathfrak{B}$ of any size.

By definition, these three Hopf algebras are closely related: the recoil algebra is a Hopf subalgebra of the Loday–Ronco algebra, which is a subalgebra of the Malvenuto–Reutenauer algebra. Moreover, there are close connections between these Hopf algebras, the lattices of Section 1.1 and the polytopes of Section 1.2. Namely,

- the product of two elements in $\mathbf{k}\mathfrak{S}$ (resp. in $\mathbf{k}\mathfrak{T}$, resp. in $\mathbf{k}\mathfrak{B}$) is a sum over an interval in a weak order (resp. a Tamari lattice, resp. a boolean lattice) [LR02].
- the product in $\mathbf{k}\mathfrak{S}$ (resp. in $\mathbf{k}\mathfrak{T}$, resp. in $\mathbf{k}\mathfrak{B}$) can be interpreted as the product (as formal power series) of the non-negative integer point enumerators of the normal cones of the permutohedra (resp. associahedra, resp. parallelepipeds) [Cha07, CHNT08, BFLR12]. The non-negative integer point enumerator of the cone $C^\circ(P)$ of a poset P on $[n]$ whose Hasse diagram is a tree is given by

$$\sum_{\mathbf{x} \in \mathbb{R}_{\geq 0}^n \cap C^\circ(P)} \mathbf{y}^{\mathbf{x}} = \prod_{i < j \in P} \frac{\mathbf{y}^{\delta_{i>j} \cdot \text{sc}(i,j,P)}}{1 - \mathbf{y}^{\text{sc}(i,j,P)}},$$

where $\text{sc}(i,j,P)$ is the connected component of i in the Hasse diagram of P minus (i,j) .

We note that these three Hopf algebras can be extended to Hopf algebras on all faces of the permutohedra, associahedra and cubes [Cha00].

We refer to [AA17, MR95, AS05, LR98, AS06, HNT05, GKL⁺95, Cha00] for more details on the topic of this section, but we want to highlight the work [AA17]. In this paper, M. Aguiar and F. Ardila use Loday's realization to explain the role of the face structure of the associahedron in the compositional inverse of power series, which was a question asked by J.-L. Loday in [Lod05]. Quoting from [AA17]: *there are many other realizations of the associahedron as a generalized permutohedron. [...] Surprisingly, to answer Loday's question within this algebro-polytopal context, Loday's realization of the associahedron is precisely the one that we need!*

1.4. Three surprising geometric properties. To close this section, we present three surprising geometric properties connecting the permutohedron $\text{Perm}(n)$ to the associahedron $\text{Asso}(n)$. The first two were already partially discussed in [Lod04], while the last one is more recent.

1.4.1. Removahedra. A *removahedron* of a polytope \mathbb{P} is a polytope obtained by removing some inequalities from the facet description of \mathbb{P} . As already mentioned, $\text{Para}(n)$ is a removahedron of $\text{Asso}(n)$, which is a removahedron of $\text{Perm}(n)$. This is illustrated in Figure 2, where the facets which are not deleted to pass to the next polytope are shaded. This property motivates the following observations:

- (1) $\text{Asso}(n)$ has precisely n pairs of parallel facets, which are the pairs of parallel facets of $\text{Para}(n)$.
- (2) $\text{Perm}(n)$ and $\text{Asso}(n)$ have precisely 2^{n-2} common vertices, which correspond to the permutations where all values before i or all values after i are larger than i for all $i \in [n]$, and to the binary trees where every node has at most one (internal) child.
- (3) $\text{Asso}(n)$ and $\text{Para}(n)$ have precisely n common vertices, which correspond to the binary trees where the left subtree of the root is a left comb and the right subtree of the root is a right comb, and to the binary sequences of the form $-^i +^{n-1-i}$.
- (4) $\text{Perm}(n)$, $\text{Asso}(n)$ and $\text{Para}(n)$ have precisely 2 common vertices, which correspond to the permutations $12\dots n$ and $n\dots 21$, to the left and right combs, and to the binary sequences $-^{n-1}$ and $+^{n-1}$.

1.4.2. Vertex barycenter. The *vertex barycenter* of a polytope \mathbb{P} is the isobarycenter of the vertices of \mathbb{P} (note that it is in general not the center of mass of \mathbb{P}). Surprisingly, the vertex barycenters of $\text{Perm}(n)$, of $\text{Asso}(n)$, and of $\text{Para}(n)$ all coincide. This was observed by F. Chapoton and reported in [Lod04, Lod05]. However, note that the argument given in [Lod04, Lod05] is wrong: the contribution of the vertex of $\text{Asso}(n)$ corresponding to a binary tree T is not the sum of the contributions of the vertices of $\text{Perm}(n)$ corresponding to the linear extensions of T . A correct argument, involving averaging over orbits of the dihedral rotation on triangulations, appeared later in [HLR10] and was simplified in [LP18]. An alternative approach based on brick polytopes (see Section 3.2) appeared in [PS15b]. Finally, an argument based on the universal associahedron (see Section 3.1) appeared in [HPS18].

1.4.3. Deformation cone. A *deformation* of a polytope \mathbb{P} can be equivalently described as (i) a polytope whose normal fan coarsens the normal fan of \mathbb{P} [McM73], (ii) a Minkowski summand of a dilate of \mathbb{P} [Mey74, She63], (iii) a polytope obtained from \mathbb{P} by perturbing the vertices so that the directions of all edges are preserved [Pos09, PRW08], (iv) a polytope obtained from \mathbb{P} by gliding its facets in the direction of their normal vectors without passing a vertex [Pos09, PRW08]. The deformations of \mathbb{P} form a cone under positive dilations and Minkowski sums, called the *deformation cone* of \mathbb{P} [McM73, PRW08]. Polytopes in the deformation cone of \mathbb{P} are parametrized by the values of the support functions in the directions normal to the facets of \mathbb{P} . Under this parametrization, this cone is defined by the *wall-crossing inequalities*, corresponding to the pairs of maximal adjacent cones of the normal fan of \mathbb{P} .

For instance, the deformations of the permutohedron $\text{Perm}(n)$ have been studied extensively (historically as *polymatroids* [Edm70], or more recently as *generalized permutohedra* [Pos09, PRW08, AA17]). They are parametrized by the cone of submodular functions. While the facets of this cone correspond to (minimal) submodular inequalities, its rays are still poorly understood (it is still an open question to determine its number of rays).

Surprisingly, the deformation cone of the associahedron $\text{Asso}(n)$ is a simplicial cone [BMCLD⁺18, PPPP19, PPR22]. Its facets are given by the wall-crossing inequalities corresponding to the exchanges of pairs of rays given by two intervals of the form $[i, j]$ and $[i+1, j+1]$. Its rays are (all positive dilations of) the faces $\Delta_{[i,j]}$ of the standard simplex corresponding to intervals of $[n]$. Hence, the sylvester fan is the normal fan of any Minkowski sum of positive dilates of the faces $\Delta_{[i,j]}$. It is however arguable that Loday's associahedron is the most natural realization of the sylvester fan, as it is the isobarycenter of the rays of its simplicial deformation cone.

2. LATTICE CONGRUENCES AND QUOTIENTOPES

In this section, we survey the deep influence of Loday's associahedron in the construction of polytopal realizations of lattice quotients of the weak order, in particular (type A) *Cambrian* and *permutoree* lattices [Rea06, PP18]. Alternative detailed sources include the original papers [Rea04, Rea06, Rea05, Rea15] and the survey articles [Rea12, Rea16a, Rea16b] for lattice quotients of the weak order, and the original papers [HL07, PP18, PS19, PPR22] for their polytopal realizations.

2.1. Sylvester congruence. A *lattice congruence* of a lattice (L, \leq, \wedge, \vee) is an equivalence relation on L that respects the meet and the join, *i.e.* such that $x \equiv x'$ and $y \equiv y'$ implies $x \wedge y \equiv x' \wedge y'$ and $x \vee y \equiv x' \vee y'$. A lattice congruence \equiv automatically defines a *lattice quotient* L/\equiv on the congruence classes of \equiv where the order relation is given by $X \leq Y$ if and only if there exists $x \in X$ and $y \in Y$ such that $x \leq y$. The meet $X \wedge Y$ (resp. the join $X \vee Y$) of two congruence classes X and Y is the congruence class of $x \wedge y$ (resp. of $x \vee y$) for arbitrary representatives $x \in X$ and $y \in Y$.

Our prototypical example of lattice congruence is the *sylvester congruence* of the weak order (the term “sylvester”, coined in [HNT05], is an adjective meaning “woody” and is not referring to the mathematician James Joseph Sylvester). Its equivalence classes are the sets of linear extensions of the binary trees of \mathfrak{B}_n , or equivalently, the fibers of the binary tree map **bt** of Section 1.1. It can be defined equivalently as the transitive closure of the rewriting rule $UacVbW \equiv^{\text{sylv}} UcaVbW$ where $a < b < c$ are letters and U, V, W are words on $[n]$. It was studied in particular in [Ton97, HNT05, Rea04, Rea06, CP17, PP18]. The quotient $\leq_w / \equiv^{\text{sylv}}$ of the weak order by the sylvester congruence is the Tamari lattice [Tam51]. In other words, for any $T, T' \in \mathfrak{T}_n$, we have $T \leq_T T'$ if and only if there exist $\sigma, \sigma' \in \mathfrak{S}_n$ such that $\mathbf{bt}(\sigma) = T$, $\mathbf{bt}(\sigma') = T'$ and $\sigma \leq_w \sigma'$.

2.2. Cambrian and permutoree congruences. A *permutoree* [PP18] is a tree whose nodes are labeled bijectively by $[n]$ and whose edges are oriented with the following local rules around each node:

- each node may have either one or two parents and either one or two children,
- if a node j has two parents (resp. children), all nodes in the left parent (resp. child) of j are smaller than j , while all nodes in the right parent (resp. child) of j are larger than j .

We decorate each node with the symbols \oplus , \otimes , \circlearrowleft , \circlearrowright depending on their number of parents and children, and the sequence $\delta(T)$ of these symbols is called the *decoration* of the permutoree T . As illustrated in Figure 3, the permutorees extend and interpolate between permutations when $\delta(T) = \oplus^n$, binary trees when $\delta(T) = \otimes^n$, and binary sequences when $\delta(T) = \circlearrowleft^n$. The permutorees with $\delta(T) \in \{\otimes, \circlearrowleft\}$ are called *Cambrian trees* [LP18, CP17].

Generalizing Section 1, it was shown in [PP18] that for each $\delta \in \{\oplus, \otimes, \circlearrowleft, \circlearrowright\}^n$,

- (1) the sets of linear extensions of the δ -permutorees define a lattice congruence of the weak order, the *δ -permutoree congruence* \equiv_δ . These sets are also the fiber of the *δ -permutoree map* sending permutations to δ -permutorees. The δ -permutoree congruence \equiv_δ is also the transitive closure of the rewriting rules

$$\begin{aligned} UacVbW &\equiv_\delta UcaVbW & \text{if } a < b < c \text{ and } \delta_b = \otimes \text{ or } \circlearrowright, \\ UbVacW &\equiv_\delta UbVcaW & \text{if } a < b < c \text{ and } \delta_b = \circlearrowleft \text{ or } \otimes, \end{aligned}$$

where a, b, c are letters while U, V, W are words on $[n]$. See [PP18, Sect. 2.3]. For instance, \equiv_δ is the trivial congruence when $\delta = \oplus^n$, the sylvester congruence [HNT05] when $\delta = \otimes^n$,




FIGURE 3. Four examples of permutorees. The first is generic, and the last three use specific decorations corresponding to permutations, binary trees, and binary sequences. [PP18, Fig. 2 & 3].




FIGURE 4. The permutabletree congruences \equiv_δ (left), δ -permutabletree lattice (middle left), the permutabletree fan $\mathcal{F}(\delta)$ (middle right), and the permutabletreehedron $\text{PT}(\delta)$ (right) for $\delta = \textcircled{1} \otimes \textcircled{Y} \otimes \textcircled{1}$. Adapted from [PP18, Figs. 11, 14 & 15] and [APR21, Fig. 9].

the (type A) Cambrian congruences [Rea04, Rea06, CP17] when $\delta \in \{\textcircled{1}, \textcircled{Y}\}^n$, and the hypoplactic congruence [KT97, Nov00] when $\delta = \textcircled{X}^n$. See Figure 4 (left) for a generic example.

- (2) the quotient \leq_W / \equiv_δ is the **δ -permutabletree lattice**, whose cover relations are right rotations on permutabletrees. See [PP18, Sect. 2.6]. For instance, the δ -permutabletree lattice is the weak order when $\delta = \textcircled{1}^n$, the Tamari lattice [Tam51] when $\delta = \textcircled{Y}^n$, the (type A) Cambrian lattices [Rea06] when $\delta \in \{\textcircled{1}, \textcircled{Y}\}^n$ and the boolean lattice when $\delta = \textcircled{X}^n$. See Figure 4 (midddle left) for a generic example.

- (3) the cones $C^\circ(T) := \{x \in \mathbf{1}^\perp \mid x_i \leq x_j, \forall i \rightarrow j \text{ in } T\}$ for all δ -permutabletrees form a complete simplicial fan, the **δ -permutabletree fan** $\mathcal{F}(\delta)$. In other words, the maximal cones of $\mathcal{F}(\delta)$ are obtained by glueing together the maximal cones of the braid fan $\mathcal{F}(n)$ which correspond to permutations in the same equivalence class for \equiv_δ . See [PP18, Sect. 3.1]. For instance, $\mathcal{F}(\delta)$ is the braid fan when $\delta = \textcircled{1}^n$, the sylvester fan when $\delta = \textcircled{Y}^n$, the (type A) Cambrian fans of [RS09] when $\delta \in \{\textcircled{1}, \textcircled{Y}\}^n$, and the coordinate fan when $\delta = \textcircled{X}^n$. See Figure 4 (middle right) for a generic example, drawn in stereographic projection.
- (4) the fan $\mathcal{F}(\delta)$ is the normal fan of the **δ -permutabletreehedron** $\text{PT}(\delta)$, which can be equivalently defined as

- the convex hull of the points $\mathbf{p}(T)$ for all δ -permutabletrees T , whose i th coordinate is defined by $\mathbf{p}(T)_i = 1 + d + \underline{\ell}r - \bar{\ell}\bar{r}$, where d denotes the number of descendants of i in T , $\underline{\ell}$ and \underline{r} denote the sizes of the left and right descendant subtrees of i in T when $\delta_i \in \{\textcircled{1}, \textcircled{X}\}$ (otherwise, $\underline{\ell} = \underline{r} = 0$), and $\bar{\ell}$ and \bar{r} denote the sizes of the left and right ancestor subtrees of i in T when $\delta_i \in \{\textcircled{Y}, \textcircled{X}\}$ (otherwise, $\bar{\ell} = \bar{r} = 0$).
- the intersection of the hyperplane \mathbb{H} with the halfspaces $\{x \in \mathbb{R}^n \mid \sum_{i \in B} x_i \geq \binom{|B|+1}{2}\}$ for each subset $B \subseteq [n]$ which is an edge cut in some δ -permutabletree (equivalently, $a, c \in B$ implies $b \in B$ or $\delta_b \in \{\textcircled{1}, \textcircled{Y}\}$, and $a, c \notin B$ implies $b \notin B$ or $\delta_b \in \{\textcircled{1}, \textcircled{X}\}$, for all $a < b < c$).

See [PP18, Sect. 3.2]. For instance, $\text{PT}(\delta)$ is the permutablehedron $\text{Perm}(n)$ when $\delta = \textcircled{1}^n$, Loday's associahedron $\text{Asso}(n)$ [SS93, Lod04] when $\delta = \textcircled{Y}^n$, Hohlweg–Lange's associahedra $\text{Asso}(\delta)$ [HL07, LP18] when $\delta \in \{\textcircled{1}, \textcircled{Y}\}^n$, and the parallelepiped $\text{Para}(n)$ when $\delta = \textcircled{X}^n$. See Figure 4 (right) for a generic example. The face lattice of the δ -permutabletreehedron $\text{PT}(\delta)$ can be described in terms of Schröder permutabletrees, see [PP18, Sect. 5]. We note that the very simple expression of the vertex coordinates of $\text{Asso}(n)$ given in [Lod04] was particularly influencial in the definition of Hohlweg–Lange's associahedra [HL07], which in turn motivated the definition of the permutablehedra of [PP18].

- (5) the Hasse diagram of the δ -permutabletree lattice is the graph of the δ -permutabletreehedron $\text{PT}(\delta)$ oriented in the direction $\omega := (n, \dots, 1) - (1, \dots, n) = \sum_{i \in [n]} (n+1-2i) e_i$.




FIGURE 5. The δ -permutohedra, for all decorations $\delta \in \Pi \cdot \{\Pi, \otimes, \circlearrowleft, \otimes\}^2 \cdot \Pi$. [PP18, Fig. 16].

Generalizing Section 1, there are natural refinements between the permutotree objects. For two decorations $\delta, \delta' \in \{\Pi, \otimes, \circlearrowleft, \otimes\}^n$ with $\delta_i \preccurlyeq \delta'_i$ for all $i \in [n]$, for the order $\Pi \preccurlyeq \{\otimes, \circlearrowleft\} \preccurlyeq \otimes$,

- the δ -permutotree congruence refines the δ' -permutotree congruence,
- the δ' -permutotree lattice is a lattice quotient of the δ -permutotree lattice,
- the δ -permutotree fan $\mathcal{F}(\delta)$ refines the δ' -permutotree fan $\mathcal{F}(\delta')$,
- the permutohedron $\text{PT}(\delta)$ is obtained by deleting inequalities in the facet description of the permutohedron $\text{PT}(\delta')$. In particular, $\text{PT}(\delta) \subseteq \text{PT}(\delta')$. See Figure 5.

The pairs of parallel facets, the common vertices of $\text{PT}(\delta)$ and $\text{PT}(\delta')$, and the isometries between permutohedra are discussed in [PP18, Sect. 3.3], with motivation from [BHLT09].

Finally, we briefly mention that the combinatorial Hopf algebras of Section 1.3 extend to a big combinatorial Hopf algebra on all permutotrees (all sizes, all decorations), see [CP17] and [PP18, Sect. 4]. Similarly, the Hopf algebras of [Cha00] extend to a Hopf algebra on all Schröder permutotrees, see [PP18, Sect. 5].

2.3. All congruences. We now consider all lattice quotients of the weak order on \mathfrak{S}_n . To keep this section short, we refrain from presenting the wonderful combinatorics of these lattice quotients in terms of non-crossing arc diagrams [Rea15], and focus on their geometric realizations.

Any lattice congruence \equiv of the weak order on \mathfrak{S}_n defines a *quotient fan* $\mathcal{F}(\equiv)$ [Rea05], whose maximal cones are obtained by glueing together the maximal cones of the braid fan $\mathcal{F}(n)$ which correspond to permutations in the same equivalence class for \equiv . This fan $\mathcal{F}(\equiv)$ is complete but not necessarily simplicial (the congruences for which the quotient fan is simplicial are characterized in [HM21, Sect. 4.4], see also [DIR⁺23, Thm. 1.13] for a representation theoretic approach, and [BNP23] for a shorter combinatorial proof and the connection to the permutotrees of Section 2.2). This fan $\mathcal{F}(\equiv)$ is the normal fan of a *quotientope* $\text{QT}(\equiv)$ which was constructed

- in [PS19] by a direct but quite intricate facet description,
- in [PPR22] as a Minkowski sum of certain simple pieces called *shard polytopes*.

See Figure 6. The graph of the quotientope, oriented in a linear direction, is the Hasse diagram of the quotient of the weak order by \equiv . We note that these quotientopes somehow simultaneously reach the limit and use the full power of Loday's simple construction of the associahedron:

- there is no simple formula for the vertex coordinates of the quotientopes similar to [Lod04].
- the only lattice congruences of the weak order which can be realized by a removahedron of $\text{Perm}(n)$ are the permutope congruences [APR21].
- however, as observed in [PPR22], any quotient fan can be realized as the normal fan of a Minkowski sum of well-chosen associahedra $\text{Asso}(\delta)$ of [HL07], which admit a simple vertex description inspired from [Lod04] and are removahedra of $\text{Perm}(n)$, as discussed in Section 2.2. For instance, the diagonal rectangulation polytope [LR12] (orange in Figure 6) is the Minkowski sum of two opposite associahedra of [Lod04] (blue and purple in Figure 6).

To complete our comparison between associahedra and arbitrary quotientopes from the perspective of Section 1, we observe that

- the graphs of the quotientopes all admit Hamiltonian paths [HM21] (it is open to prove that they all admit a Hamiltonian cycle).
- the vertex barycenters of the quotientopes do not coincide (not even for permutopeheda),
- the deformation cone of a quotientope $\text{QT}(\equiv)$ is simplicial if and only if the congruence \equiv is refined by a Cambrian congruence [APR21, PP23].
- generalizing the Hopf algebras of [MR95, LR98, GKL⁺95] described in Section 1.3, various Hopf algebra structures have been investigated, either on specific congruences [Rea05, LR12, Gir12, CP17, PP18, Pil18, Law14, NRT11], or on all lattice quotients [Pil19].

2.4. Beyond the braid arrangement. Consider now a central hyperplane arrangement \mathcal{H} defining a fan \mathcal{F} , and a distinguished base region B of \mathcal{F} . The *poset of regions* $\text{R}(\mathcal{H}, B)$ is the set of regions of \mathcal{F} ordered by inclusion of their separating sets (the set of hyperplanes of \mathcal{H} that separate the given region from the base region B). For instance, the poset of region is the weak order on W when \mathcal{H} is the Coxeter arrangement of a finite Coxeter group W [Hum90, BB05]. In general, the poset of regions $\text{R}(\mathcal{H}, B)$ is always a lattice when the fan \mathcal{F} is simplicial, and never a lattice when the chamber B is not simplicial [BEZ90]. See also the survey of N. Reading [Rea16b] for further conditions, in particular a discussion on tight arrangements.

Assume now that $\text{R}(\mathcal{H}, B)$ is a lattice, and consider a lattice congruence \equiv of $\text{R}(\mathcal{H}, B)$. It was proved in [Rea05] that the lattice congruence \equiv defines a complete fan $\mathcal{F}(\equiv)$ obtained by glueing together the cones of the fan \mathcal{F} that belong to the same congruence class of \equiv . It remains an open question whether these quotient fans are polytopal. The answer is known to be positive in for:

- the braid arrangement (whose poset of regions is the weak order on permutations) by [PS19, PPR22] as discussed in Section 2.3,
- graphical arrangements of skeletal graphs (whose poset of regions are precisely the acyclic reorientation lattices) by [Pil21],
- the hyperoctahedral arrangement (or type B Coxeter arrangement) by [PPR22].

Using quiver representation theory (see Section 3.3), [DHT23] also recently proposed candidates for shard polytopes, which should lead to quotientopes for arbitrary finite Coxeter arrangements. We note also that quotientopes for the braid arrangement and for graphical arrangements can be constructed as Minkowski sums of (lower dimensional) associahedra of [HL07], showing again the longlasting influence of Loday's associahedron [Lod04].




FIGURE 6. The quotientope lattice for $n = 4$: all quotientopes ordered by inclusion (which corresponds to refinement of the lattice congruences). We only consider lattice congruences whose fan is essential. We have highlighted the cube (green), Loday's associahedron [Lod04] (blue), another one of Hohlweg–Lange's associahedra [HL07] (purple), the diagonal rectangulation polytope [LR12] (orange), and the permutohedron (red). Adapted from [PS19, Fig. 9].

3. CLUSTER ALGEBRAS, BRICK POLYTOPES, AND QUIVER REPRESENTATION THEORY

In this section, we present some polytope constructions in the theory of cluster algebras, of subword complexes, and of quiver representation theory, in which Loday's associahedra were instrumental. Alternative surveys on this section include [FWZ, FR07, Hoh12, Tho12, Tho21].

3.1. Cluster algebras and generalized associahedra. Cluster algebras were introduced in the series of papers [FZ02, FZ03a, BFZ05, FZ07]. Their motivations came from total positivity and canonical bases, but cluster algebras quickly appeared to be a fundamental structure in many areas of mathematics (representation theory of quivers, Poisson geometry, integrable systems, etc). See the cluster algebra portal [Fom], or the surveys [FR07, FWZ].

A *cluster algebra* is a commutative ring generated by a set of *cluster variables* grouped into overlapping *clusters*. One can choose an initial cluster as a *seed* from which all other clusters are obtained by a *mutation process* controlled by a combinatorial object (a skew-symmetrizable matrix, or a weighted quiver). During a mutation, a single variable in the cluster is perturbed and the new variable is computed by an *exchange relation*. One fundamental aspect of this process is the Laurent phenomenon [FZ02]: all cluster variables are Laurent polynomials with respect to the cluster variables of the initial seed.

An important combinatorial and geometric object associated to a cluster algebra is its cluster complex: the simplicial complex with cluster variables as vertices and clusters as facets. A cluster algebra is of *finite type* if its cluster complex is finite. Finite type cluster algebras are classified by the same Cartan–Killing classification of finite root systems [FZ03a], and there are combinatorial models for the cluster variables and clusters of the cluster algebras of non-exceptional finite types. In particular, the cluster complex of the cluster algebra of type A_{n-1} is isomorphic to the simplicial associahedron: cluster variables correspond to internal diagonals of an $(n+2)$ -gon, clusters correspond to triangulations of this polygon, a mutation between clusters corresponds to a flip between triangulations, and the exchange relation can even be interpreted as a Ptolemy relation in a quadrilateral.

A finite type cluster algebra (together with a choice of initial seed) naturally defines two complete simplicial fans, both combinatorially isomorphic to the cluster complex but geometrically different: the \mathbf{d} -vector fan [FZ03b] and the \mathbf{g} -vector fan [FZ07]. The information needed to construct these fans is encoded in the algebra as follows: the \mathbf{d} -vectors are given by the denominators of the cluster variables, while the \mathbf{g} -vectors are given by exponents of coefficients in the cluster algebra with principal coefficients. In type A_{n-1} , the \mathbf{d} -vector fan is the compatibility fan (the ray corresponding to a diagonal δ of the $(n+2)$ -gon consists of all positive multiples of the characteristic vector of the diagonals of the initial triangulation of the $(n+2)$ -gon crossed by δ), while the \mathbf{g} -vector fan is the sylvester fan introduced in Section 1.2. In fact, the \mathbf{g} -vector fans of finite type cluster algebras with respect to acyclic initial seeds are precisely the *Cambrian fans* of [RS09] realizing the *Cambrian lattices* of [Rea06] in crystallographic types. The Cambrian lattices are particular lattice quotients of the weak orders on Coxeter groups [Hum90, BB05]. (Note that \mathbf{g} -vector fans are not limited to acyclic initial seeds but are restricted to Weyl groups, while Cambrian lattices and fans are limited to acyclic initial seeds, but are defined for arbitrary finite Coxeter groups). See Figure 8 for illustrations of \mathbf{g} -vector fans.

The *generalized associahedron* of a finite type cluster algebra is a simple polytope whose polar realizes the cluster complex. Generalized associahedra were first constructed in [CFZ02] using the \mathbf{d} -vector fans, and alternative realizations were obtained in [HLT11] using the \mathbf{g} -vector fans with respect to acyclic initial seeds (in fact, the Cambrian fans of [RS09] for arbitrary finite Coxeter groups). The latter realizations are direct descendants of Loday's associahedra. Namely, generalizing the construction of the associahedra of [HL07] (directly inspired by [Lod04] as discussed in Section 2.2), they are obtained by deleting inequalities in the facet description of the *Coxeter permutohedron*, the convex hull of the orbit of a generic point under the action of the reflection group. The remaining facets are those that contain at least one singleton, *i.e.* a point of the Coxeter permutohedron corresponding to a singleton class of the Cambrian congruence (thus a common point with the resulting generalized associahedron). The isometry classes of generalized

associahedra of [HLT11] were described in [BHLT09]. See Figure 7 and the first two columns of Figure 8 for illustrations. We refer to [Hoh12] for a very instructive presentation.

The construction of [HLT11] was revisited in [Ste13] with an approach similar to the original one of [CFZ02], and in [PS15a] via brick polytopes (see Section 3.2). Later, it was proved in [HPS18] that all \mathbf{g} -vector fans with respect to any initial seed (acyclic or not) are actually polytopal (this construction yields the same generalized associahedron as [HLT11] when it starts from an acyclic initial seed). See the last two columns of Figure 8 for illustrations. This construction also led to the definition of a *universal associahedron* [HPS18], a polytope whose normal fan simultaneously contains all \mathbf{g} -vector fans of a given finite type cluster algebra.

To complete our connection between Loday's associahedra [Lod04] and the generalized associahedra of [HLT11, HPS18] from the perspective of Section 1, we observe that

- the diameters of the mutation graphs of the finite type cluster algebras have been determined in [Pou14, CP16, Pou17], see [CP16, Tab. 2]. Moreover, all generalized associahedra have the non-leaving face property [CP16, Wil17].
- the realizations of [HPS18] are removalahedra of the underlying zonotope only when the initial seed is acyclic (hence the resulting generalized associahedron coincides with that of [HLT11]) or when the cluster algebra is of type A . See Figure 8.
- the vertex barycenters of all generalized associahedra of [HLT11, HPS18] coincide with that of the Coxeter permutohedron [PS15b, HPS18].
- the deformation cone of a generalized associahedron of [HLT11, HPS18] is always simplicial [BMCLD⁺18, PPPP19]. The rays of this deformation cone are (positive dilations of) the Newton polytopes of the F -polynomials [FZ07] of the cluster variables of the cluster algebra [BMCLD⁺18].

Finally, let us mention that cluster algebras and in particular the associahedron appear in an extremely promising recent line of research in high energy physics. In quantum field theory, the scattering amplitudes (the probabilities that particular interactions occur among particles) are traditionally expressed as sums over all possible Feynman diagrams for the interaction. In [AHT14], N. Arkani-Hamed and J. Trnka introduce *amplituhedra*, geometric objects that greatly simplify these computations. Although the theory is still under construction, in the case treated in [AHT14] (Super Yang-Mills theory with $N = 4$), the amplituhedron is a linear image of the positive Grassmannian and the scattering amplitudes are computed by evaluating a certain form in it, conjecturally the volume of a “dual amplituhedron” which should exist.

In [AHBHY18], this approach is applied to the so-called “bi-adjoint ϕ^3 scalar theory”, for which the amplituhedron turns out to be exactly Loday's associahedron. The relation is as follows: in this theory n cyclically ordered particles in the plane interact via ternary trees. Hence, there is one Feynman diagram corresponding to each vertex of $\text{Asso}(n - 2)$, that is, to each facet F of the polar $\text{Asso}(n - 2)^\circ$. It turns out that the summand of each ternary tree in the expression for the scattering amplitude equals the volume of the simplex obtained coning the corresponding facet F to the origin, so indeed the total sum equals the volume of $\text{Asso}(n - 2)^\circ$. There is a deformation of $\text{Asso}(n - 2)$ involved in the process. Before computing the polar, each facet of $\text{Asso}(n - 2)$ is translated by an amount depending in the momenta of the two particles defining that facet. In fact, the treatment of deformed associahedra in [AHBHY18] inspired the works [BMCLD⁺18, PPPP19] mentioned above.

The relation between associahedron-like polytopes and scattering amplitudes has been explored intensively in the past few years, see *e.g.* [BLR19, Ram19, ABJ⁺20, Koj20, JKM20, JL21, AJK19, Kal20c, Kal20a, Kal20b, KJ20, Chh19, Sal19, SS21].

3.2. Subword complexes and brick polytopes. Subword complexes were introduced in [KM05] in the context of Gröbner geometry of Schubert varieties, and extended to all finite Coxeter groups in [KM04]. Given a finite Coxeter system (W, S) [Hum90, BB05], a word Q of S^m , and an element w of W , the *subword complex* $\mathcal{SC}(Q, w)$ is the simplicial complex of subwords of Q whose complements contain a reduced expression of w . In other words, its ground set is the set $[m]$ of positions in Q , and its facets are the complements of the reduced expressions of w in Q . Here,




FIGURE 7. Some Coxeter permutohedra (red) and generalized associahedra of [HLT11] (blue) in type A (top), B (middle), and D (bottom). The singletons (common vertices of the permutohedron and associahedron) are marked (purple). Adapted from [PS15a, Figs. 14, 15 & 16].




FIGURE 8. Some g -vector fans and generalized associahedra of finite type cluster algebras. Top: all type A_3 and the cyclic type C_3 initial exchange matrices. Middle: The corresponding dual c -vector fans (thin red) and g -vector fans (bold blue). Bottom: The corresponding zonotopes (thin red) and generalized associahedra (bold blue). In type A , the generalized associahedra are removalhedra of the corresponding zonotope. In cyclic type C_3 , the shaded facets of the zonotope and of the associahedron are parallel but do not coincide. Adapted from [HPS18, Figs. 2, 3, 5 & 6].




FIGURE 9. The sorting network \mathcal{N}_Q corresponding to the word $Q = \tau_2\tau_3\tau_1\tau_3\tau_2\tau_1\tau_2\tau_3\tau_1$ (left) and the pseudoline arrangements corresponding to the facets $\{2, 3, 5\}$ (middle) and $\{2, 3, 9\}$ (right). Adapted from [PS12, Figs. 3 & 4].

we only consider the case where $w = w_\circ$ is the longest element of W , and Q contains a subword which is a reduced expression for w_\circ , and we just write $\mathcal{SC}(Q)$ for $\mathcal{SC}(Q, w_\circ)$.

The subword complex $\mathcal{SC}(Q)$ is known to be a vertex-decomposable simplicial sphere [KM04]. The question of whether these simplicial spheres are polytopal is a longstanding open problem [KM04, PS09, PP12, PS12, CLS14, PS15a, CZ12, CS22, Cre23]. Largely inspired from Loday's associahedron [Lod04], the brick polytope of [PS12, PS15a] was designed as an attempt to solve this question.

To a facet I of $\mathcal{SC}(Q)$ and a position $k \in [m]$, we associate a root $\mathbf{r}(I, k) := \Pi Q_{[k-1] \setminus I}(\alpha_{q_k})$ [CLS14] and a weight $\mathbf{w}(I, k) := \Pi Q_{[k-1] \setminus I}(\alpha_{q_k})$ [PS15a], where ΠQ_X denotes the product of the reflections $q_x \in Q$, for $x \in X$, in the order given by Q . The *root configuration* of I is the set $\mathbf{R}(I) := \{\mathbf{r}(I, i) \mid i \in I\}$ and the *brick vector* is the vector $\mathbf{b}(I) := \sum_{k \in [l]} \mathbf{w}(I, k)$. The *brick polytope* $\text{BP}(Q)$ of the word Q is the convex hull of the brick vectors of all facets of $\mathcal{SC}(Q)$ [PS12, PS15a]. It was shown in [PS12, PS15a, JS21] that the vertices of $\text{BP}(Q)$ correspond to the facets I of $\mathcal{SC}(Q)$ whose root configuration $\mathbf{R}(I)$ is acyclic (*i.e.* form a pointed cone). It follows that the brick polytope $\text{BP}(Q)$ realizes the subword complex $\mathcal{SC}(Q)$ if and only if the root configurations of all (or equivalently, of one of) the facets of $\mathcal{SC}(Q)$ is linearly independant.

Subword complexes have a simple visual interpretation in type A_{n-1} , *i.e.* when $W = \mathfrak{S}_n$ and $S = \{\tau_p \mid p \in [n-1]\}$ where τ_p is the simple transposition $(p \ p+1)$. A (primitive) *sorting network* \mathcal{N} is formed by n horizontal lines (its *levels*, labeled from bottom to top) together with m vertical segments (its *commutators*, labeled from left to right) joining two consecutive levels. A *pseudoline arrangement* on \mathcal{N} is a collection of n x -monotone paths supported by \mathcal{N} which cross pairwise precisely once at a commutator and have no other intersection. The commutators where two pseudolines cross are called *crossings* while the others are called *contacts* of the pseudoline arrangement. A word $Q := q_1 q_2 \cdots q_m$ on S is represented by the sorting network \mathcal{N}_Q whose k th commutator lies between the p th and $(p+1)$ th levels if $q_k = \tau_p$. A facet I of $\mathcal{SC}(Q)$ is then represented by the pseudoline arrangements whose contacts lie at the positions given by I . See Figure 9. As pointed out in [PP12], type A subword complexes can be used to provide a combinatorial model for many relevant families of geometric graphs (see Figure 10 for illustrations):

- triangulations of a convex polygon [Woo04, PP12, Stu11, SS12].
- k -triangulations of a convex polygon [PS09, PP12, Stu11, SS12]. A *k -triangulation* of a convex $(n+2k)$ -gon is a maximal set of diagonals such that no $k+1$ of them are pairwise crossing [CP92, Jon05, Nak00, DKM02, PS09]. Some of the corresponding brick polytopes are illustrated in Figure 11. We note that these particular subword complexes have various connections with lattice quotients and Hopf algebras discussed in Sections 1 and 2 [Pil18].
- pseudotriangulations of a point set P in general position (no line contains three points). A *pseudotriangulations* of P is a maximal pointed crossing-free set of edges between points of P [PV96, RSS08]. Pseudotriangulations correspond to the vertices of the *pseudotriangulation polytope* of [RSS03].
- pseudotriangulations of a set of disjoint convex bodies in general position (no line is tangent to three convex bodies).




FIGURE 10. Sorting networks interpretations of certain geometric graphs: a triangulation of the convex octagon, a 2-triangulation of the convex octagon, a pseudotriangulation of a point set, and a pseudotriangulation of a set of disjoint convex bodies. [PS15a, Fig. 6].




FIGURE 11. The brick polytopes $\text{BP}(c^k w_o(c))$ for $c = \tau_1 \tau_2 \tau_3$ in type A_3 and $k \in [3]$. Adapted from [Pil18, Fig. 11].

In particular, for subword complexes corresponding to triangulations of a convex polygon, the brick polytopes precisely recover the associahedra of [Lod04, HL07] (up to a translation).

More generally, the subword complex interpretation of [PP12, Stu11, SS12] for triangulations and multitriangulations of convex polygons was extended to arbitrary finite Coxeter groups in [CLS14]. In particular, for a Coxeter element c of a finite Coxeter group, the c -cluster complex is isomorphic to the subword complex $\mathcal{SC}(cw_o(c))$ where $w_o(c)$ denotes the c -sorting word of w_o (meaning the lexicographic minimal reduced word in $c^\infty := ccc\cdots$). The corresponding brick polytope $\text{BP}(cw_o(c))$ then coincides (up to translation) with the c -associahedra $\text{Asso}(c)$ of [HLT11] discussed in Section 3.1 [PS15a]. This alternative interpretation provides an explicit vertex description of $\text{Asso}(c)$, and thus enables to easily derive some of their geometric properties, for instance that their barycenter all coincide with that of the Coxeter permutohedron [PS15b].

For completeness, let us mention that non-spherical subword complexes also have their brick polyhedra [JS21], whose oriented skeleta are sometimes lattice quotients of intervals of the weak order [BCCP23].

3.3. Quiver representation theory and gentle associahedra. To illustrate the far reaching influence of Loday's associahedra, we conclude with some of its apparitions and generalizations in the quiver representation theory.

There are several connections between representation theory and the associahedron. For instance, the introductory paper [Tho12] connects the lattice of torsion classes of the linear A_{n-1} quiver to the Tamari lattice. Here, we prefer to follow [Tho23] to go straight from representation theory to Loday's associahedron.

A *quiver* $Q := (Q_0, Q_1)$ is a directed graph with vertices Q_0 and arrows Q_1 . A *representation* V of Q is an assignment of a finite dimensional vector space V_i (over a fixed algebraic closed field) for each vertex i of Q_0 , and of a linear map $V_\alpha : V_i \rightarrow V_j$ for each arrow $i \xrightarrow{\alpha} j$ of Q_1 . Representations are considered up to isomorphisms, meaning basis changes in each of the vector spaces V_i .

Note that the representations of Q correspond to the modules over the path algebra of Q (the algebra of oriented paths on Q , where the product of two paths π, π' is the concatenation $\pi\pi'$ when π' starts where π ends, and 0 otherwise). As any (basic) finite dimensional algebra is the path algebra of a quiver with relations, the theory of quiver representations (with relations) thus corresponds to the theory of modules over finite dimensional algebras. We refer to the textbooks [ASS06, Sch14].

The *dimension vector* of a representation V is $\dim V := (\dim V_i)_{i \in Q_0}$. The *direct sum* of two representations V, W of Q is the representation $V \oplus W$ of Q defined by $(V \oplus W)_i = V_i \oplus W_i$ for each vertex i of Q_0 , and $(V \oplus W)_\alpha = V_\alpha \oplus W_\alpha$ for each arrow α of Q_1 . A representation is *indecomposable* if it cannot be decomposed as the direct sum of two non-zero representations. A *subrepresentation* of a representation V is a representation W such that W_i is a vector subspace of V_i for each vertex i of Q_0 , and $V_\alpha(W_i) \subseteq W_j$ for each arrow $i \xrightarrow{\alpha} j$ of Q_1 . The *Harder–Narasimhan polytope* $\text{HN}(V)$ of a representation V is the convex hull of the dimension vectors of the subrepresentations of V . It transports direct sums to Minkowski sums: $\text{HN}(V \oplus W) = \text{HN}(V) + \text{HN}(W)$. In particular, if Q admits only finitely many indecomposable representations (up to isomorphism), it is natural to consider $\text{HN}(Q) := \text{HN}(\bigoplus_V V) = \sum_V \text{HN}(V)$, where the (direct and Minkowski) sums range over all (representatives of the isomorphism classes of) indecomposable representations of Q .

Consider now the linear A_{n-1} quiver Q , with $Q_0 = [n-1]$ and $Q_1 = \{i \xrightarrow{\alpha_i} i+1 \mid i \in [n-2]\}$. Up to isomorphism, its indecomposable representations are precisely the representations E^{ij} for $1 \leq i \leq j \leq n-1$, consisting in a one-dimensional vector space at each vertex between i and j (included) with identity maps between them, and zero vector spaces and maps elsewhere. One can check the Harder–Narasimhan polytope $\text{HN}(E^{ij})$ is (up to a change of basis) the face $\Delta_{[i,j+1]}$ of the standard simplex corresponding to the interval $[i, j+1]$. Hence, $\text{HN}(Q)$ coincides with Loday's associahedron $\text{Asso}(n)$. Moreover, as already mentioned, the Tamari lattice given by the linear orientation of the graph of $\text{Asso}(n)$ corresponds to the lattice of torsion classes of Q [DIR⁺23].

Note that starting from any orientation Q of a type $A/D/E$ Dynkin quiver, the polytopes $\text{HN}(V)$ of the indecomposable representations V would be the Newton polytopes of the F -polynomials of the corresponding cluster algebra [BMCLD⁺18], the polytope $\text{HN}(Q)$ would coincide with another associahedron of [HLT11], and the lattice of torsion classes of Q' would be a Cambrian lattice [Rea06] of type $A/D/E$ [IT09].

Even more powerful statements arise in the situation of quivers with relations (see [ASS06, Sch14] for definitions). Two families of examples deserve a particular mention here as they are closely connected to Loday's associahedron and the material discussed here.

The first family is that of *preprojective quivers* of type $A/D/E$. The polytope $\text{HN}(Q)$ is the corresponding Coxeter permutohedron [AHI⁺22], and the lattice of torsion classes is the weak order of the corresponding finite Coxeter group [Miz14]. Moreover, the Harder–Narasimhan polytopes $\text{HN}(V)$ of the indecomposable brick representations of \tilde{Q} are natural candidates for shard polytopes [DHT23], which would enable to realize all lattice quotients of these weak orders, as mentioned in Section 2.4. The same construction should extend to arbitrary finite Weyl groups, working with quiver representations on non algebraically closed fields.

The second family is that of *gentle quivers* [AS87, BR87]. The lattice of torsion classes of a gentle quiver was interpreted combinatorially in terms of non-kissing walks of a blossoming quiver,




FIGURE 12. Some gentle quivers (top), their gentle fans (middle) and their gentle associahedra (bottom). Adapted from [PPP21, Figs. 40 & 41] and [PPP19, Fig. 30].

or equivalently in terms of non-crossing accordions on a dissection of a surface [BDM⁺20, PPP21, PPP19]. Moreover, gentle quivers with finitely many indecomposables naturally define a gentle fan (generalizing the sylvester fan) and a gentle associahedron (generalizing Loday's associahedron), whose constructions were directly inspired from that of Section 1, see [PPP21, PPPP19]. See Figure 12 for illustrations. These gentle associahedra specialize to and uniformize *grid associahedra* previously considered in [PPS10, SSW17, McC17, GM17], and *accordiohedra* previously considered in [Bar01, Cha16, GM18, MP19].

4. GRAPH ASSOCIAHEDRA AND NESTOHEDRA

In this section, we briefly present the families of graph associahedra and hypergraph associahedra (or nestohedra). They were constructed in [CD06, Dev09, Pos09, FS05, DP11] in connection to wonderful compactifications of hyperplane arrangements [DCP95]. The associahedron of [Lod04] served again as the prototype in these constructions.

4.1. Graph associahedra. Consider a simple graph G with vertex set V . A *tube* of G is a non-empty subset of V which induces a connected subgraph of G . Two tubes are *compatible* if they are either nested, or disjoint and non-adjacent (their union is not a tube). A *tubing* of G is a set of pairwise compatible tubes, which contains the connected components of G . The *nested complex* of G is the simplicial complex of tubings of G . It can be geometrically realized by the *nested fan*, with a ray $\mathbf{g}(t)$ for each tube t of G , given by (the projection of) the characteristic vector of t . Moreover, the nested fan is the normal fan of a polytope, called *graph associahedron* $\text{Asso}(G)$, and first constructed in [CD06] (see also [DJS03]). This polytope can be obtained

- by truncating some faces of the standard simplex [CD06],
- as the intersection of \mathbb{H} with the hyperplanes $\langle \mathbf{g}(t) | \mathbf{x} \rangle \leq -3^{|t|}$ for all tubes t of G [Dev09],
- as the Minkowski sum $\sum_t \Delta_t$ of the faces of the standard simplex given by all tubes t of G [Pos09].

See the first two columns of Figure 14 for two generic examples. As illustrated in Figure 13, the graph associahedra of certain special families of graphs coincide with well-known families of




FIGURE 13. The permutohedron (left), the associahedron (middle), and the cyclohedron (right) as graph associahedra. Adapted from [MPP22].

polytopes: complete graph associahedra are permutohedra, path associahedra are classical associahedra, cycle associahedra are cyclohedra, star associahedra are stellohedra, and parallelotopes are empty graph associahedra (meaning graphs with no edges).

We restrict ourselves to a few observations motivated by Section 1:

- graph properties of graph associahedra have been investigated in [MP15, CMM22, CPVP22]: their graphs are Hamiltonian, and their diameters are partially understood.
- the oriented graphs of graph associahedra are not always Hasse diagrams of lattices. The ones which are lattice quotients of the weak order have been characterized in [BM21].
- the nested fan of G is realized as a removahedron of $\text{Perm}(n)$ if and only if every cycle of G induces a clique [Pil17]. In particular, the cyclohedron is not a removahedron of $\text{Perm}(n)$.
- the deformation cone of the graph associahedra are described in [PPP23]. In particular, the Cartesian products of associahedra are the only graph associahedra whose deformation cone is simplicial.
- Hopf algebraic structures on graph associahedra were investigated in [FS10, Ron12, BM21].

4.2. Hypergraph associahedra and nestohedra. *Hypergraph associahedra* [DP11] are obtained exactly as graph associahedra by replacing the initial graph data by a hypergraph [Ber89]. They were described independently as *nestohedra* or *building sets* [Pos09, FS05, Zel06]. They are constructed as the Minkowski sums of the faces of the standard simplex corresponding to their tubes [Pos09], and thus belong to the family of hypergraphical polytopes [BBM19] inside the rich family of generalized permutohedra [Pos09]. See the last two columns of Figure 14 for two generic examples.

A particularly interesting family of hypergraph associahedra is obtained from interval hypergraphs (hypergraphs all of whose hyperedges are intervals of $[n]$): they contain

- the classical associahedron of [SS93, Lod04] for the building set with all intervals of $[n]$,
- the Pitman–Stanley polytope of [SP02] for the building set with all singletons $\{i\}$ and all initial intervals $[i]$ for $i \in [n]$,
- the freehedron of [San09] for the building set with all singletons $\{i\}$, all initial intervals $[i]$ for $i \in [n]$, and all final intervals $[n] \setminus [i]$ for $i \in [n - 1]$,
- the fertilotopes of [Def21] for the binary building sets defined as the interval building sets where any two intervals are either nested or disjoint.

For completeness, we just briefly mention that

- the hypergraph associahedra which are removahedra of $\text{Perm}(n)$ were characterized in [Pil17].
- the deformation cones of hypergraph associahedra were described in [PPP23]. In particular, all interval hypergraph associahedra have a simplicial deformation cone.

Various generalizations of nestohedra were studied *e.g.* in [DFRS15, Gal21, MPP22].




FIGURE 14. Some hypergraphs (top), nested fans (middle) and hypergraph associahedra (bottom). The first two are graphical, the last two are hypergraphical. Adapted from [PPP23, Fig. 4, 5, 7 & 8].

5. OPERADS AND DIAGONALS

In this section, we come to the relation of the associahedron to loop spaces. Although this was the original motivation for J. Stasheff [Sta63], we have delayed this topic until the end since in Stasheff's original work only the combinatorics of the associahedron is important. In fact, he defined it as a CW-complex with no particular embedding as a polytope. Only recently Loday's realization has acquired importance in this area, for the construction of diagonals of associahedra.

We briefly introduce loop spaces and the notion of A_∞ -operad, and then focus on the diagonal of the associahedron where (a weighted version of) Loday's construction was fundamentally exploited [MTTV21]. Expository texts for this section include [Sta04, LV12, Val14].

5.1. Loop spaces and operads. In a pointed topological space $(X, *)$, a *loop* is a continuous map $f : [0, 1] \rightarrow X$ with $f(0) = f(1) = *$. The *concatenation product* fg of two loops f and g is defined by $fg(t) = f(2t)$ if $t \leq 1/2$ and $fg(t) = g(2t - 1)$ if $t \geq 1/2$. This product fails to be associative: for three loops f, g, h , the images of $f(gh)$ and $(fg)h$ coincide, but their parametrizations differ. However, one can easily find a homotopy that deforms $f(gh)$ to $(fg)h$.

If we now consider four loops f, g, h, k , then their five possible concatenations (parenthesizations of the word $fghk$) are homotopic, but also the two possible ways to compose homotopies between the loops $f(g(hk))$ and $((fg)h)k$ are homotopic. Continuing this process, J. Stasheff [Sta63] was led to the definition of an *A_∞ -algebra*, or *homotopy associative algebra*, that is, a topological monoid together with an infinite tower of homotopies correcting coherently the defect of associativity of the product. Such a structure defines a homotopy invariant characterization of loop spaces: a space is a loop space if and only if it possesses an A_∞ -algebra structure, encoded by the associahedra.

These associahedra in turn form an A_∞ operad, equivalent to the little intervals operads. An *operad* is an algebraic structure encoding a type of algebras (for instance, associative, commutative, Lie algebras, or A_∞ -algebras in this case). See [Sta04, MSS02, LV12, Val14, Gir18] for introductions and references. The notion of operad first appeared in the seminal work of P. May [May72]




FIGURE 15. The diagonal of the associahedron in dimension 2 (left) and 3 (right). The 3-dimensional picture is a courtesy of G. Laplante-Anfossi. [LA22, Fig. 13 (left)].




FIGURE 16. The diagonal of the 2-dimensional associahedron, where faces are labeled by pairs of Schröder trees (left) and attached to their max-min pair of binary trees given by the magical formula (right). Adapted from [BCP23, Fig. 2].

on iterated loop spaces, where he proved the *recognition principle*, generalizing J. Stasheff’s result: a space is a k -fold loop space (space of functions from the k -dimensional sphere to a pointed topological space, sending the north pole to the base point) if and only if it is an algebra over the little k -cubes operad. Today, operads are ubiquitous in mathematics: in addition to algebraic topology, they are used in differential geometry, algebraic geometry, non-commutative geometry, mathematical physics and probability, among other fields [MSS02, LV12]. Many of these fields interact in the work [DSV19] where the toric varieties associated with Loday’s associahedra are used to define a non-symmetric analogue of the little 2-cubes operad, leading to a non-commutative notion of cohomological field theory with Givental-type symmetries.

5.2. Cellular diagonal of the associahedron. Taking cellular chains on associahedra one arrives to the algebraic version of an A_∞ -algebra. This algebraic structure plays a prominent role in many fields, for instance in symplectic topology where it describes the fine structure of the Fukaya category of a symplectic manifold [Sei08]. In this context, but also in others such as the study of the homology of fibered spaces [Pro86], one is led to the problem of defining a universal tensor product of A_∞ -algebras, which can be achieved by the construction of a cellular approximation of the diagonal of the associahedra.

The diagonal of a polytope P is the map $\delta : P \rightarrow P \times P$ defined by $x \mapsto (x, x)$. A *cellular diagonal* of P is a map $\tilde{\delta} : P \rightarrow P \times P$ homotopic to δ , which agrees with δ on the vertices of P , and whose image is a union of faces of $P \times P$. For face-coherent families of polytopes (*i.e.* when faces are products of polytopes in the family, like simplices, cubes, permutohedra or associahedra), some algebraic purposes additionally require the cellular diagonal to be compatible with the face structure. Finding cellular diagonals in such families of polytopes is an important challenge at

the crossroad of operad theory, homotopical algebra, combinatorics and discrete geometry, see [SU04, MS06, Lod11, MTTV21, LA22].

For the family of associahedra, algebraic diagonals were described in [SU04] and later in [MS06, Lod11]. However, there were no known topological diagonals as defined above, until the recent work of [MTTV21] defining a cellular diagonal Δ_n for (a weighted version of) the associahedron $\text{Asso}(n)$ of [Lod04, SS93] (and recovering, at the cellular level, all the previous formulas [SU22, DOJVLA⁺23]). We note that the method of [MTTV21] essentially relies on the theory of fiber polytopes of [BS92], and enables to see the cellular diagonal of the associahedron as a polytopal complex refining the associahedron, see Figure 15.

The face structure of the cellular diagonal Δ_n is given by the *magical formula* [MTTV21]. Namely, the k -dimensional faces correspond to the pairs (F, G) of faces of the associahedron $\text{Asso}(n)$ with $\dim(F) + \dim(G) = k$ and $\max(F) \leq \min(G)$ (where \leq , \max and \min refer to the order given by the Tamari lattice). See Figure 16. In particular, the vertices of Δ_n correspond to intervals of the Tamari lattice, which are counted by

$$\frac{2}{(3n+1)(3n+2)} \binom{4n+1}{n+1}$$

as proved in [Cha07, Cha18]. This formula also counts the rooted 3-connected planar triangulations with $2n+2$ faces, and explicit bijections were given in [BB09, FFN23]. More generally, the f -vector of Δ_n is given by

$$f_k(\Delta_n) = \frac{2}{(3n+1)(3n+2)} \binom{n-1}{k} \binom{4n+1-k}{n+1}.$$

as proved in [BCP23, FFN23].

Note that the construction of the diagonal Δ_n , its face description by the magical formula, and the product formulas for its f -vector all rely essentially on the fact that the normal fan of the associahedron is the sylvester fan and behaves nicely with the Tamari lattice (Section 1).

6. FURTHER GENERALIZATIONS

We believe that the main impact of Loday's description of the associahedron was to break the psychological barrier of realizing this "mythical polytope" [Hai84] by showing a natural and well-behaved realization. Consequently, this construction was the first seed of a teeming forest of polytopal realizations of combinatorial structures. Even if it is impossible to cite all its descendants, we conclude with a partial list of polytopes obtained from Loday's associahedron, with pointers to the literature for the interested readers.

- (1) Cartesian products and Minkowski sums:
 - quotientopes [PS19, PPR22] (presented in Section 2),
 - multiplihedron, graph multiplihedron, and multinestohedron [For08, DF08, AD13, CP22],
 - biassociahedron [SU11, Mar15, CP22],
 - constrainahedron [Pol21, BP22, CP22],
- (2) sections and projections:
 - accordiohedra [MP19] (and their extensions to gentle algebras [PPP21, PPPP19]),
 - poset associahedra and acyclonestohedra [Gal21, Sac23, MPP22],
- (3) polyhedral decompositions:
 - ν -associahedra [CPS19], in connection to multivariate diagonal harmonics [Ber13, BPR12, PRV17],
 - diagonals of the associahedron [MTTV21, BCP23] (presented in Section 5.2),
- (4) further descendants:
 - pebble tree associahedra [PT18, Pil22],
 - categorical k -associahedra [Bot19, BP22, BBP23],
 - permutoassociahedra [Kap93, RZ94, FKS17, BIP19, Iva20, CL23].

We hope that this survey invites the reader to develop further the family of these associahedra.

ACKNOWLEDGEMENTS

Our presentation in Section 3.3 closely follows a talk by Hugh Thomas [Tho23] and our Section 5 is shaped from personal communications with Guillaume Laplante-Anfossi who is also the author of Figure 15, reproduced from [LA22, Fig. 13 (left)]. We are grateful to both of them for letting us use that material. We also thank both of them, together with Spencer Backman, Nate Bottman, Cesar Ceballos, Satyan Devadoss, Stefan Forcey, Christophe Hohlweg, Torsten Mütze, and an anonymous referee for comments and suggestions on our original text. We also deeply benefited from reports and corrections by Frédéric Chapoton, Nathan Reading and Hugh Thomas on the Habilitation thesis of V. Pilaud [Pil20], on which this survey is largely based.

REFERENCES

- [AA17] Marcelo Aguiar and Federico Ardila. Hopf monoids and generalized permutohedra. Preprint, [arXiv:1709.07504](https://arxiv.org/abs/1709.07504). To appear in *Mem. Amer. Math. Soc.*, 2017.
- [ABJ⁺20] P. B. Aneesh, Pinaki Banerjee, Mrunmay Jagadale, Renjan Rajan John, Alok Laddha, and Sujoy Mahato. On positive geometries of quartic interactions: Stokes polytopes, lower forms on associahedra and world-sheet forms. *J. High Energy Phys.*, 2020(4):149, 46, 2020.
- [AD13] Federico Ardila and Jeffrey Doker. Lifted generalized permutohedra and composition polynomials. *Adv. in Appl. Math.*, 50(4):607–633, 2013.
- [AHBHY18] Nima Arkani-Hamed, Yuntao Bai, Song He, and Gongwang Yan. Scattering forms and the positive geometry of kinematics, color and the worldsheet. *J. High Energy Phys.*, 2018(5):096, 75, 2018.
- [AHI⁺22] Toshitaka Aoki, Akihiro Higashitani, Osamu Iyama, Ryoichi Kase, and Yuya Mizuno. Fans and polytopes in tilting theory I: Foundations. Preprint, [arXiv:2203.15213](https://arxiv.org/abs/2203.15213), 2022.
- [AHT14] Nima Arkani-Hamed and Jaroslav Trnka. The amplituhedron. *J. High Energy Phys.*, 2014(10):30, 2014.
- [AJK19] P. B. Aneesh, Mrunmay Jagadale, and Nikhil Kalyanapuram. Accordiohedra as positive geometries for generic scalar field theories. *Phys. Rev. D*, 100(10):106013, 12, 2019.
- [APR21] Doriann Albertin, Vincent Pilaud, and Julian Ritter. Removahedral congruences versus permutree congruences. *Electron. J. Combin.*, 28(4):Paper No. 4.8, 38, 2021.
- [AS87] Ibrahim Assem and Andrzej Skowroński. Iterated tilted algebras of type $\tilde{\mathbf{A}}_n$. *Math. Z.*, 195(2):269–290, 1987.
- [AS05] Marcelo Aguiar and Frank Sottile. Structure of the Malvenuto-Reutenauer Hopf algebra of permutations. *Adv. Math.*, 191(2):225–275, 2005.
- [AS06] Marcelo Aguiar and Frank Sottile. Structure of the Loday-Ronco Hopf algebra of trees. *J. Algebra*, 295(2):473–511, 2006.
- [ASS06] Ibrahim Assem, Daniel Simson, and Andrzej Skowroński. *Elements of the representation theory of associative algebras. Vol. 1*, volume 65 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 2006. Techniques of representation theory.
- [Bar01] Yuliy Baryshnikov. On Stokes sets. In *New developments in singularity theory (Cambridge, 2000)*, volume 21 of *NATO Sci. Ser. II Math. Phys. Chem.*, pages 65–86. Kluwer Acad. Publ., Dordrecht, 2001.
- [BB05] Anders Björner and Francesco Brenti. *Combinatorics of Coxeter groups*, volume 231 of *Graduate Texts in Mathematics*. Springer, New York, 2005.
- [BB09] Olivier Bernardi and Nicolas Bonichon. Intervals in Catalan lattices and realizers of triangulations. *J. Combin. Theory Ser. A*, 116(1):55–75, 2009.
- [BBM19] Carolina Benedetti, Nantel Bergeron, and John Machacek. Hypergraphic polytopes: combinatorial properties and antipode. *J. Comb.*, 10(3):515–544, 2019.
- [BBP23] Spencer Backman, Nathaniel Bottman, and Daria Poliakova. Higher categorical associahedra. In preparation, 2023.
- [BCCP23] Nantel Bergeron, Noémie Cartier, Cesar Ceballos, and Vincent Pilaud. Lattices of acyclic pipe dreams. Preprint, [arXiv:2303.11025](https://arxiv.org/abs/2303.11025), 2023.
- [BCP23] Alin Bostan, Frédéric Chyzak, and Vincent Pilaud. Refined product formulas for Tamari intervals. Preprint, [arXiv:2303.10986](https://arxiv.org/abs/2303.10986), 2023.
- [BDM⁺20] Thomas Brüstle, Guillaume Douville, Kaveh Mousavand, Hugh Thomas, and Emine Yıldırım. On the combinatorics of gentle algebras. *Canad. J. Math.*, 72(6):1551–1580, 2020.
- [Ber89] Claude Berge. *Hypergraphs*, volume 45 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam, 1989. Combinatorics of finite sets, Translated from the French.
- [Ber13] François Bergeron. Multivariate diagonal coinvariant spaces for complex reflection groups. *Adv. Math.*, 239:97–108, 2013.
- [BEZ90] Anders Björner, Paul H. Edelman, and Günter M. Ziegler. Hyperplane arrangements with a lattice of regions. *Discrete Comput. Geom.*, 5(3):263–288, 1990.

- [BFLR12] Adrien Boussicault, Valentin Féray, Alain Lascoux, and Victor Reiner. Linear extension sums as valuations on cones. *J. Algebraic Combin.*, 35(4):573–610, 2012.
- [BFS90] Louis J. Billera, Paul Filliman, and Bernd Sturmfels. Constructions and complexity of secondary polytopes. *Adv. Math.*, 83(2):155–179, 1990.
- [BFZ05] Arkady Berenstein, Sergey Fomin, and Andrei Zelevinsky. Cluster algebras. III. Upper bounds and double Bruhat cells. *Duke Math. J.*, 126(1):1–52, 2005.
- [BHLT09] Nantel Bergeron, Christophe Hohlweg, Carsten Lange, and Hugh Thomas. Isometry classes of generalized associahedra. *Sém. Lothar. Combin.*, 61A:Art. B61Aa, 13, 2009.
- [BIP19] Djordje Baralić, Jelena Ivanović, and Zoran Petrić. A simple permutoassociahedron. *Discrete Math.*, 342(12):111591, 18, 2019.
- [Bjö84] Anders Björner. Orderings of Coxeter groups. In *Combinatorics and algebra (Boulder, Colo., 1983)*, volume 34 of *Contemp. Math.*, pages 175–195. Amer. Math. Soc., Providence, RI, 1984.
- [BKT14] Pierre Baumann, Joel Kamnitzer, and Peter Tingley. Affine Mirković-Vilonen polytopes. *Publ. Math. Inst. Hautes Études Sci.*, 120:113–205, 2014.
- [BLR19] Pinaki Banerjee, Alok Laddha, and Prashanth Raman. Stokes polytopes: the positive geometry for ϕ^4 interactions. *J. High Energy Phys.*, 2019(8):067, 34, 2019.
- [BM21] Emily Barnard and Thomas McConvile. Lattices from graph associahedra and subalgebras of the Malvenuto-Reutenauer algebra. *Algebra Universalis*, 82(1):Paper No. 2, 53, 2021.
- [BMCLD⁺18] Véronique Bazier-Matte, Nathan Chapelier-Laget, Guillaume Douville, Kaveh Mousavand, Hugh Thomas, and Emine Yildirim. ABHY Associahedra and Newton polytopes of F -polynomials for finite type cluster algebras. Preprint, [arXiv:1808.09986](https://arxiv.org/abs/1808.09986), 2018.
- [BNP23] Emily Barnard, Jean-Christophe Novelli, and Vincent Pilaud. On simple congruences of the weak order. In preparation, 2023.
- [Bot19] Nathaniel Bottman. 2-associahedra. *Algebr. Geom. Topol.*, 19(2):743–806, 2019.
- [BP22] Nathaniel Bottman and Daria Poliakova. Constraintahedra. Preprint, [arXiv:2208.14529](https://arxiv.org/abs/2208.14529), 2022.
- [BPR12] François Bergeron and Louis-François Préville-Ratelle. Higher trivariate diagonal harmonics via generalized Tamari posets. *Journal of Combinatorics*, 3(3):317–341, 2012.
- [BR87] Michael C. R. Butler and Claus Michael Ringel. Auslander-Reiten sequences with few middle terms and applications to string algebras. *Comm. Algebra*, 15(1-2):145–179, 1987.
- [BS92] Louis J. Billera and Bernd Sturmfels. Fiber polytopes. *Ann. of Math. (2)*, 135(3):527–549, 1992.
- [CD06] Michael P. Carr and Satyan L. Devadoss. Coxeter complexes and graph-associahedra. *Topology Appl.*, 153(12):2155–2168, 2006.
- [CFZ02] Frédéric Chapoton, Sergey Fomin, and Andrei Zelevinsky. Polytopal realizations of generalized associahedra. *Canad. Math. Bull.*, 45(4):537–566, 2002.
- [Cha07] Frédéric Chapoton. Sur le nombre d'intervalles dans les treillis de Tamari. *Sém. Lothar. Combin.*, 55:Art. B55f, 18, 2005/07.
- [Cha00] Frédéric Chapoton. Algèbres de Hopf des permutohèdres, associahèdres et hypercubes. *Adv. Math.*, 150(2):264–275, 2000.
- [Cha07] Frédéric Chapoton. The anticyclic operad of moulds. *Int. Math. Res. Not. IMRN*, 2007(20):Art. ID rnm078, 36, 2007.
- [Cha16] Frédéric Chapoton. Stokes posets and serpent nests. *Discrete Math. Theor. Comput. Sci.*, 18(3), 2016.
- [Cha18] Frédéric Chapoton. Une note sur les intervalles de Tamari. *Ann. Math. Blaise Pascal*, 25(2):299–314, 2018.
- [Chh19] Saroj Chhatoi. A note on convex realization of haloedron. Preprint, [arXiv:1910.13786](https://arxiv.org/abs/1910.13786), 2019.
- [CHNT08] Frédéric Chapoton, Florent Hivert, Jean-Christophe Novelli, and Jean-Yves Thibon. An operational calculus for the mould operad. *Int. Math. Res. Not. IMRN*, 2008(9):Art. ID rnm018, 22, 2008.
- [CL23] Federico Castillo and Fu Liu. The permuto-associahedron revisited. *European J. Combin.*, 110:Paper No. 103706, 30, 2023.
- [CLS14] Cesar Ceballos, Jean-Philippe Labb  , and Christian Stump. Subword complexes, cluster complexes, and generalized multi-associahedra. *J. Algebraic Combin.*, 39(1):17–51, 2014.
- [CMM22] Jean Cardinal, Arturo Merino, and Torsten Mütze. Efficient generation of elimination trees and graph associahedra. In *Proceedings of the 2022 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 2128–2140, 2022. Extended abstract of [arXiv:2106.16204](https://arxiv.org/abs/2106.16204).
- [CP92] Vasilis Capoyleas and J  nos Pach. A Tur  n-type theorem on chords of a convex polygon. *J. Combin. Theory Ser. B*, 56(1):9–15, 1992.
- [CP16] Cesar Ceballos and Vincent Pilaud. The diameter of type D associahedra and the non-leaving-face property. *European J. Combin.*, 51:109–124, 2016.
- [CP17] Gr  gory Chatel and Vincent Pilaud. Cambrian Hopf Algebras. *Adv. Math.*, 311:598–633, 2017.
- [CP22] Fr  d  ric Chapoton and Vincent Pilaud. Shuffles of deformed permutohedra, multiplihedra, constraintahedra, and biassociahedra. Preprint, [arXiv:2201.06896](https://arxiv.org/abs/2201.06896), 2022.
- [CPS19] Cesar Ceballos, Arnau Padrol, and Camilo Sarmiento. Geometry of ν -Tamari lattices in types A and B . *Trans. Amer. Math. Soc.*, 371(4):2575–2622, 2019.

- [CPVP22] Jean Cardinal, Lionel Pournin, and Mario Valencia-Pabon. Diameter Estimates for Graph Associahedra. *Ann. Comb.*, 26(4):873–902, 2022.
- [Cre23] Luis Crespo Ruiz. Realizations of multiassociahedra via bipartite rigidity. Preprint, [arXiv:2303.15776](https://arxiv.org/abs/2303.15776), 2023.
- [CS22] Luis Crespo Ruiz and Francisco Santos. Realizations of multiassociahedra via rigidity. Preprint, [arXiv:2212.14265](https://arxiv.org/abs/2212.14265), 2022.
- [CSZ15] Cesar Ceballos, Francisco Santos, and Günter M. Ziegler. Many non-equivalent realizations of the associahedron. *Combinatorica*, 35(5):513–551, 2015.
- [CZ12] Cesar Ceballos and Günter M. Ziegler. Realizing the Associahedron: Mysteries and Questions. In *Associahedra, Tamari lattices and related structures*, volume 299 of *Prog. Math.*, pages 129–159. Birkhäuser/Springer, Basel, 2012.
- [DCP95] Conrado De Concini and Claudio Procesi. Wonderful models of subspace arrangements. *Selecta Math. (N.S.)*, 1(3):459–494, 1995.
- [Def21] Colin Defant. Fertilopes. Preprint, [arXiv:2102.11836](https://arxiv.org/abs/2102.11836). To appear in *Discr. & Comput. Geom.*, 2021.
- [Deh10] Patrick Dehornoy. On the rotation distance between binary trees. *Adv. Math.*, 223(4):1316–1355, 2010.
- [Dev09] Satyan L. Devadoss. A realization of graph associahedra. *Discrete Math.*, 309(1):271–276, 2009.
- [DF08] Satyan Devadoss and Stefan Forcey. Marked tubes and the graph multiplihedron. *Algebr. Geom. Topol.*, 8(4):2081–2108, 2008.
- [DFRS15] Satyan L. Devadoss, Stefan Forcey, Stephen Reisdorf, and Patrick Showers. Convex polytopes from nested posets. *European J. Combin.*, 43:229–248, 2015.
- [DHT23] Will Dana, Eric Hanson, and Hugh Thomas. Shard polytopes via representation theory. In preparation, 2023.
- [DIR⁺23] Laurent Demonet, Osamu Iyama, Nathan Reading, Idun Reiten, and Hugh Thomas. Lattice theory of torsion classes: Beyond τ -tilting theory. *Trans. Amer. Math. Soc. Ser. B*, 10:542–612, 2023.
- [DJS03] Michael Davis, Tadeuz Januszkiwicz, and Richard Allan Scott. Fundamental groups of blow-ups. *Adv. Math.*, 177(1):115–179, 2003.
- [DKM02] Andreas Dress, Jack H. Koolen, and Vincent L. Moulton. On line arrangements in the hyperbolic plane. *European J. Combin.*, 23(5):549–557, 2002.
- [DOJVLA⁺23] Bérénice Delcroix-Oger, Mathieu Josuat-Vergès, Guillaume Laplante-Anfossi, Vincent Pilaud, and Kurt Stoeckl. The combinatorics of the permutohedron diagonals. In preparation, 2023.
- [DP11] Kosta Došen and Zoran Petrić. Hypergraph polytopes. *Topology Appl.*, 158(12):1405–1444, 2011.
- [DRS10] Jesus A. De Loera, Jörg Rambau, and Francisco Santos. *Triangulations: Structures for Algorithms and Applications*, volume 25 of *Algorithms and Computation in Mathematics*. Springer Verlag, 2010.
- [DSV19] Vladimir Dotsenko, Sergey Shadrin, and Bruno Vallette. Toric varieties of Loday’s associahedra and noncommutative cohomological field theories. *J. Topol.*, 12(2):463–535, 2019.
- [Edm70] Jack Edmonds. Submodular functions, matroids, and certain polyhedra. In *Combinatorial Structures and their Applications (Proc. Calgary Internat. Conf., Calgary, Alta., 1969)*, pages 69–87. Gordon and Breach, New York, 1970.
- [FFN23] Wenjie Fang, Éric Fusy, and Philippe Nadeau. Bijections between Tamari intervals and blossoming trees. In preparation, 2023.
- [FKS17] Stefan Forcey, Logan Keefe, and William Sands. Split-facets for balanced minimal evolution polytopes and the permutoassociahedron. *Bull. Math. Biol.*, 79(5):975–994, 2017.
- [Fom] Sergey Fomin. Cluster algebras portal. <http://www.math.lsa.umich.edu/~fomin/cluster.html>.
- [For08] Stefan Forcey. Convex hull realizations of the multiplihedra. *Topology Appl.*, 156(2):326–347, 2008.
- [FR07] Sergey Fomin and Nathan Reading. Root systems and generalized associahedra. In *Geometric combinatorics*, volume 13 of *IAS/Park City Math. Ser.*, pages 63–131. Amer. Math. Soc., Providence, RI, 2007.
- [FS05] Eva Maria Feichtner and Bernd Sturmfels. Matroid polytopes, nested sets and Bergman fans. *Port. Math. (N.S.)*, 62(4):437–468, 2005.
- [FS10] Stefan Forcey and Derriell Springfield. Geometric combinatorial algebras: cyclohedron and simplex. *J. Algebraic Combin.*, 32(4):597–627, 2010.
- [FWZ] Sergey Fomin, Lauren Williams, and Andrei Zelevinsky. Introduction to cluster algebras. In preparation. First chapters available as [arXiv:1608.05735](https://arxiv.org/abs/1608.05735), [arXiv:1707.07190](https://arxiv.org/abs/1707.07190), [arXiv:2008.09189](https://arxiv.org/abs/2008.09189), and [arXiv:2106.02160](https://arxiv.org/abs/2106.02160).
- [FZ02] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. I. Foundations. *J. Amer. Math. Soc.*, 15(2):497–529, 2002.
- [FZ03a] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. II. Finite type classification. *Invent. Math.*, 154(1):63–121, 2003.
- [FZ03b] Sergey Fomin and Andrei Zelevinsky. Y -systems and generalized associahedra. *Ann. of Math. (2)*, 158(3):977–1018, 2003.

- [FZ07] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. IV. Coefficients. *Compos. Math.*, 143(1):112–164, 2007.
- [Gal21] Pavel Galashin. Poset associahedra. Preprint, [arXiv:2110.07257](https://arxiv.org/abs/2110.07257), 2021.
- [Gir12] Samuele Giraudo. Algebraic and combinatorial structures on pairs of twin binary trees. *J. Algebra*, 360:115–157, 2012.
- [Gir18] Samuele Giraudo. *Nonsymmetric operads in combinatorics*. Springer, Cham, 2018.
- [GKL⁺95] Israel M. Gelfand, Daniel Krob, Alain Lascoux, Bernard Leclerc, Vladimir S. Retakh, and Jean-Yves Thibon. Noncommutative symmetric functions. *Adv. Math.*, 112(2):218–348, 1995.
- [GKZ08] Israel Gelfand, Mikhail Kapranov, and Andrei Zelevinsky. *Discriminants, resultants and multidimensional determinants*. Modern Birkhäuser Classics. Birkhäuser Boston Inc., Boston, MA, 2008. Reprint of the 1994 edition.
- [GM17] Alexander Garver and Thomas McConville. Enumerative properties of Grid-Associahedra. Preprint, [arXiv:1705.04901](https://arxiv.org/abs/1705.04901), 2017.
- [GM18] Alexander Garver and Thomas McConville. Oriented flip graphs of polygonal subdivisions and noncrossing tree partitions. *J. Combin. Theory Ser. A*, 158:126–175, 2018.
- [GR63] Georges Th. Guilbaud and Pierre Rosenstiehl. Analyse algébrique d'un scrutin. *Math. Inform. Sci. Humaines*, (4):9–33, 1963.
- [Hai84] Mark Haiman. Constructing the associahedron. Unpublished manuscript, 11 pages, available at <http://www.math.berkeley.edu/~mchaiman/ftp/assoc/manuscript.pdf>, 1984.
- [HL07] Christophe Hohlweg and Carsten Lange. Realizations of the associahedron and cyclohedron. *Discrete Comput. Geom.*, 37(4):517–543, 2007.
- [HLR10] Christophe Hohlweg, Jonathan Lortie, and Annie Raymond. The centers of gravity of the associahedron and of the permutohedron are the same. *Electron. J. Combin.*, 17(1):Research Paper 72, 14, 2010.
- [HLT11] Christophe Hohlweg, Carsten Lange, and Hugh Thomas. Permutahedra and generalized associahedra. *Adv. Math.*, 226(1):608–640, 2011.
- [HM21] Hung P. Hoang and Torsten Mütze. Combinatorial generation via permutation languages. II. Lattice congruences. *Israel J. Math.*, 244(1):359–417, 2021.
- [HN99] Ferran Hurtado and Marc Noy. Graph of triangulations of a convex polygon and tree of triangulations. *Comput. Geom.*, 13(3):179–188, 1999.
- [HNT05] Florent Hivert, Jean-Christophe Novelli, and Jean-Yves Thibon. The algebra of binary search trees. *Theoret. Comput. Sci.*, 339(1):129–165, 2005.
- [Hoh12] Christophe Hohlweg. Permutahedra and associahedra: generalized associahedra from the geometry of finite reflection groups. In *Associahedra, Tamari lattices and related structures*, volume 299 of *Prog. Math.*, pages 129–159. Birkhäuser/Springer, Basel, 2012.
- [HPS18] Christophe Hohlweg, Vincent Pilaud, and Salvatore Stella. Polytopal realizations of finite type g-vector fans. *Adv. Math.*, 328:713–749, 2018.
- [Hum90] James E. Humphreys. *Reflection groups and Coxeter groups*, volume 29 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1990.
- [IT09] Colin Ingalls and Hugh Thomas. Noncrossing partitions and representations of quivers. *Compos. Math.*, 145(6):1533–1562, 2009.
- [Iva20] Jelena Ivanović. Geometrical realisations of the simple permutoassociahedron by Minkowski sums. *Appl. Anal. Discrete Math.*, 14(1):55–93, 2020.
- [JKM20] Renjan Rajan John, Ryota Kojima, and Sujoy Mahato. Weights, recursion relations and projective triangulations for positive geometry of scalar theories. *J. High Energy Phys.*, 2020, 2020.
- [JL21] Mrunmay Jagadale and Alok Laddha. On the positive geometry of quartic interactions III : one loop integrands from polytopes. *J. High Energy Phys.*, 2021(7), 2021.
- [Joh63] Selmer M. Johnson. Generation of permutations by adjacent transposition. *Math. Comp.*, 17:282–285, 1963.
- [Jon05] Jakob Jonsson. Generalized triangulations and diagonal-free subsets of stack polyominoes. *J. Combin. Theory Ser. A*, 112(1):117–142, 2005.
- [JS21] Dennis Jahn and Christian Stump. Bruhat intervals, subword complexes and brick polyhedra for finite Coxeter groups. Preprint, [arXiv:2103.03715](https://arxiv.org/abs/2103.03715). To appear in *Math. Z.*, 2021.
- [Kal20a] Nikhil Kalyanapuram. Geometric recursion from polytope triangulations and twisted homology. *Phys. Rev. D*, 102(12):125027, 8, 2020.
- [Kal20b] Nikhil Kalyanapuram. On polytopes and generalizations of the KLT relations. *J. High Energy Phys.*, 2020(12):057, 31, 2020.
- [Kal20c] Nikhil Kalyanapuram. Stokes polytopes and intersection theory. *Phys. Rev. D*, 101(10):105010, 16, 2020.
- [Kap93] Mikhail M. Kapranov. The permutoassociahedron, Mac Lane's coherence theorem and asymptotic zones for the KZ equation. *J. Pure Appl. Algebra*, 85(2):119–142, 1993.
- [Kel01] Bernhard Keller. Introduction to A-infinity algebras and modules. *Homology Homotopy Appl.*, 3(1):1–35., 2001.

- [KJ20] Nikhil Kalyanapuram and Raghav G. Jha. Positive geometries for all scalar theories from twisted intersection theory. *Phys. Rev. Research*, 2:033119, 2020.
- [KM04] Allen Knutson and Ezra Miller. Subword complexes in Coxeter groups. *Adv. Math.*, 184(1):161–176, 2004.
- [KM05] Allen Knutson and Ezra Miller. Gröbner geometry of Schubert polynomials. *Ann. of Math. (2)*, 161(3):1245–1318, 2005.
- [Knu11] Donald E. Knuth. *The art of computer programming. Vol. 4A. Combinatorial algorithms. Part 1*. Addison-Wesley, Upper Saddle River, NJ, 2011.
- [Koj20] Ryota Kojima. Weights and recursion relations for ϕ^p tree amplitudes from the positive geometry. *J. High Energy Phys.*, 2020:54, 2020.
- [KT97] Daniel Krob and Jean-Yves Thibon. Noncommutative symmetric functions. IV. Quantum linear groups and Hecke algebras at $q = 0$. *J. Algebraic Combin.*, 6(4):339–376, 1997.
- [LA22] Guillaume Laplante-Anfossi. The diagonal of the operahedra. *Adv. Math.*, 405:Paper No. 108494, 50, 2022.
- [Law14] Shirley Law. Combinatorial realization of the Hopf algebra of sashes. Preprint, [arXiv:1407.4073](https://arxiv.org/abs/1407.4073), 2014.
- [Lee89] Carl W. Lee. The associahedron and triangulations of the n -gon. *European J. Combin.*, 10(6):551–560, 1989.
- [Lod04] Jean-Louis Loday. Realization of the Stasheff polytope. *Arch. Math. (Basel)*, 83(3):267–278, 2004.
- [Lod05] Jean-Louis Loday. The multiple facets of the associahedron. Available at <https://www.claymath.org/library/academy/LectureNotes05/Lodaypaper.pdf>, 2005.
- [Lod11] Jean-Louis Loday. The diagonal of the Stasheff polytope. In *Higher structures in geometry and physics*, volume 287 of *Progr. Math.*, pages 269–292. Birkhäuser/Springer, New York, 2011.
- [LP18] Carsten Lange and Vincent Pilaud. Associahedra via spines. *Combinatorica*, 38(2):443–486, 2018.
- [LR98] Jean-Louis Loday and María O. Ronco. Hopf algebra of the planar binary trees. *Adv. Math.*, 139(2):293–309, 1998.
- [LR02] Jean-Louis Loday and María O. Ronco. Order structure on the algebra of permutations and of planar binary trees. *J. Algebraic Combin.*, 15(3):253–270, 2002.
- [LR12] Shirley Law and Nathan Reading. The Hopf algebra of diagonal rectangulations. *J. Combin. Theory Ser. A*, 119(3):788–824, 2012.
- [Luc87] Joan M. Lucas. The rotation graph of binary trees is Hamiltonian. *J. Algorithms*, 8(4):503–535, 1987.
- [LV12] Jean-Louis Loday and Bruno Vallette. *Algebraic operads*, volume 346 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Heidelberg, 2012.
- [Mar15] Martin Markl. Bipermutahedron and biassociahedron. *J. Homotopy Relat. Struct.*, 10(2):205–238, 2015.
- [May72] J. Peter May. *The geometry of iterated loop spaces*. Lecture Notes in Mathematics, Vol. 271. Springer-Verlag, Berlin-New York, 1972.
- [McC17] Thomas McConville. Lattice structure of Grid-Tamari orders. *J. Combin. Theory Ser. A*, 148:27–56, 2017.
- [McM73] Peter McMullen. Representations of polytopes and polyhedral sets. *Geometriae Dedicata*, 2:83–99, 1973.
- [Mey74] Walter Meyer. Indecomposable polytopes. *Trans. Amer. Math. Soc.*, 190:77–86, 1974.
- [MHPS12] Folkert Müller-Hoissen, Jean Marcel Pallo, and Jim Stasheff, editors. *Associahedra, Tamari Lattices and Related Structures. Tamari Memorial Festschrift*, volume 299 of *Progress in Mathematics*. Springer, New York, 2012.
- [Miz14] Yuya Mizuno. Classifying τ -tilting modules over preprojective algebras of Dynkin type. *Math. Z.*, 277(3-4):665–690, 2014.
- [MP15] Thibault Manneville and Vincent Pilaud. Graph properties of graph associahedra. *Sém. Lothar. Combin.*, pages Art. B73d, 31 pp, 2015.
- [MP17] Thibault Manneville and Vincent Pilaud. Compatibility fans for graphical nested complexes. *J. Combin. Theory Ser. A*, 150:36–107, 2017.
- [MP19] Thibault Manneville and Vincent Pilaud. Geometric realizations of the accordion complex of a dissection. *Discrete Comput. Geom.*, 61(3):507–540, 2019.
- [MPP22] Chiara Mantovani, Arnau Padrol, and Vincent Pilaud. Poset associahedra as sections of graph associahedra. In preparation, 2022.
- [MR95] Claudia Malvenuto and Christophe Reutenauer. Duality between quasi-symmetric functions and the Solomon descent algebra. *J. Algebra*, 177(3):967–982, 1995.
- [MS06] Martin Markl and Steve Shnider. Associahedra, cellular W -construction and products of A_∞ -algebras. *Trans. Amer. Math. Soc.*, 358(6):2353–2372, 2006.
- [MSS02] Martin Markl, Steve Shnider, and Jim Stasheff. *Operads in algebra, topology and physics*, volume 96 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2002.

- [MTTV21] Naruki Masuda, Hugh Thomas, Andy Tonks, and Bruno Vallette. The diagonal of the associahedra. *J. Éc. polytech. Math.*, 8:121–146, 2021.
- [Nak00] Tomoki Nakamigawa. A generalization of diagonal flips in a convex polygon. *Theoret. Comput. Sci.*, 235(2):271–282, 2000. Combinatorics and optimization (Okinawa, 1996).
- [Nov00] Jean-Christophe Novelli. On the hypoplactic monoid. *Discrete Math.*, 217(1-3):315–336, 2000. Formal power series and algebraic combinatorics (Vienna, 1997).
- [NRT11] Jean-Christophe Novelli, Christophe Reutenauer, and Jean-Yves Thibon. Generalized descent patterns in permutations and associated Hopf algebras. *European J. Combin.*, 32(4):618–627, 2011.
- [Pil17] Vincent Pilaud. Which nestohedra are removahedra? *Rev. Colombiana Mat.*, 51(1):21–42, 2017.
- [Pil18] Vincent Pilaud. Brick polytopes, lattice quotients, and Hopf algebras. *J. Combin. Theory Ser. A*, 155:418–457, 2018.
- [Pil19] Vincent Pilaud. Hopf algebras on decorated noncrossing arc diagrams. *J. Combin. Theory Ser. A*, 161:486–507, 2019.
- [Pil20] Vincent Pilaud. *From permutohedra to associahedra, a walk through geometric and algebraic combinatorics*. Habilitation à Diriger des Recherches, Université Paris-Saclay, 2020. Available at <http://www.lix.polytechnique.fr/~pilaud/documents/reports/habilitationVincentPilaud.pdf>.
- [Pil21] Vincent Pilaud. Acyclic reorientation lattices and their lattice quotients. Preprint, [arXiv:2111.12387](https://arxiv.org/abs/2111.12387), 2021.
- [Pil22] Vincent Pilaud. Pebble trees. Preprint, [arXiv:2205.06686](https://arxiv.org/abs/2205.06686), 2022.
- [Pol21] Daria Poliakova. *Homotopical algebra and combinatorics of polytopes*. PhD thesis, University of Copenhagen, 2021.
- [Pos09] Alexander Postnikov. Permutohedra, associahedra, and beyond. *Int. Math. Res. Not. IMRN*, 2009(6):1026–1106, 2009.
- [Pou14] Lionel Pournin. The diameter of associahedra. *Adv. Math.*, 259:13–42, 2014.
- [Pou17] Lionel Pournin. The asymptotic diameter of cyclohedra. *Israel J. Math.*, 219(2):609–635, 2017.
- [PP12] Vincent Pilaud and Michel Pocchiola. Multitriangulations, pseudotriangulations and primitive sorting networks. *Discrete Comput. Geom.*, 48(1):142–191, 2012.
- [PP18] Vincent Pilaud and Viviane Pons. Permutrees. *Algebraic Combinatorics*, 1(2):173–224, 2018.
- [PP23] Vincent Pilaud and Germain Poullot. Deformation cones of quotientopes. In preparation, 2023.
- [PPP19] Yann Palu, Vincent Pilaud, and Pierre-Guy Plamondon. Non-kissing and non-crossing complexes for locally gentle algebras. *J. Comb. Algebra*, 3(4):401–438, 2019.
- [PPP21] Yann Palu, Vincent Pilaud, and Pierre-Guy Plamondon. Non-kissing complexes and tau-tilting for gentle algebras. *Mem. Amer. Math. Soc.*, 274(1343), 2021.
- [PPP23] Arnau Padrol, Vincent Pilaud, and Germain Poullot. Deformation cones of graph associahedra and nestohedra. *European J. Combin.*, 107:103594, 2023.
- [PPPP19] Arnau Padrol, Yann Palu, Vincent Pilaud, and Pierre-Guy Plamondon. Associahedra for finite type cluster algebras and minimal relations between g -vectors. Preprint, [arXiv:1906.06861](https://arxiv.org/abs/1906.06861), 2019.
- [PPR22] Arnau Padrol, Vincent Pilaud, and Julian Ritter. Shard polytopes. *Int. Math. Res. Not. IMRN*, 2022, 2022. online first.
- [PPS10] T. Kyle Petersen, Pavlo Pylyavskyy, and David E. Speyer. A non-crossing standard monomial theory. *J. Algebra*, 324(5):951–969, 2010.
- [Pro86] Alain Prouté. A_∞ -structures. *Modèles minimaux de Baues-Lemaire et Kadeishvili et homologie des fibrations*. PhD thesis, Université Paris 7, 1986.
- [PRV17] Louis-François Préville-Ratelle and Xavier Viennot. The enumeration of generalized Tamari intervals. *Trans. Amer. Math. Soc.*, 369(7):5219–5239, 2017.
- [PRW08] Alexander Postnikov, Victor Reiner, and Lauren K. Williams. Faces of generalized permutohedra. *Doc. Math.*, 13:207–273, 2008.
- [PS09] Vincent Pilaud and Francisco Santos. Multitriangulations as complexes of star polygons. *Discrete Comput. Geom.*, 41(2):284–317, 2009.
- [PS12] Vincent Pilaud and Francisco Santos. The brick polytope of a sorting network. *European J. Combin.*, 33(4):632–662, 2012.
- [PS15a] Vincent Pilaud and Christian Stump. Brick polytopes of spherical subword complexes and generalized associahedra. *Adv. Math.*, 276:1–61, 2015.
- [PS15b] Vincent Pilaud and Christian Stump. Vertex barycenter of generalized associahedra. *Proc. Amer. Math. Soc.*, 143(6):2623–2636, 2015.
- [PS19] Vincent Pilaud and Francisco Santos. Quotientopes. *Bull. Lond. Math. Soc.*, 51(3):406–420, 2019.
- [PT18] Kate Poirier and Thomas Tradler. The combinatorics of directed planar trees. *J. Combin. Theory Ser. A*, 160:31–61, 2018.
- [PV96] Michel Pocchiola and Gert Vegter. Topologically sweeping visibility complexes via pseudotriangulations. *Discrete Comput. Geom.*, 16(4):419–453, 1996.
- [Rad52] Richard Rado. An inequality. *J. London Math. Soc.*, 27:1–6, 1952.
- [Ram19] Prashanth Raman. The positive geometry for ϕ^p interactions. *J. High Energy Phys.*, 2019(10):271, 33, 2019.
- [Rea04] Nathan Reading. Lattice congruences of the weak order. *Order*, 21(4):315–344, 2004.

- [Rea05] Nathan Reading. Lattice congruences, fans and Hopf algebras. *J. Combin. Theory Ser. A*, 110(2):237–273, 2005.
- [Rea06] Nathan Reading. Cambrian lattices. *Adv. Math.*, 205(2):313–353, 2006.
- [Rea12] Nathan Reading. From the tamari lattice to cambrian lattices and beyond. In Folkert Müller-Hoissen, Jean Pallo, and Jim Stasheff, editors, *Associahedra, Tamari Lattices and Related Structures – Tamari Memorial Festschrift*, volume 299 of *Progress in Mathematics*, pages 293–322. Birkhäuser, 2012.
- [Rea15] Nathan Reading. Noncrossing arc diagrams and canonical join representations. *SIAM J. Discrete Math.*, 29(2):736–750, 2015.
- [Rea16a] Nathan Reading. Finite Coxeter groups and the weak order. In *Lattice theory: special topics and applications. Vol. 2*, pages 489–561. Birkhäuser/Springer, Cham, 2016.
- [Rea16b] Nathan Reading. Lattice theory of the poset of regions. In *Lattice theory: special topics and applications. Vol. 2*, pages 399–487. Birkhäuser/Springer, Cham, 2016.
- [Ron12] María Ronco. Generalized Tamari order. In *Associahedra, Tamari lattices and related structures*, volume 299 of *Progr. Math.*, pages 339–350. Birkhäuser/Springer, Basel, 2012.
- [RS09] Nathan Reading and David E. Speyer. Cambrian fans. *J. Eur. Math. Soc.*, 11(2):407–447, 2009.
- [RSS03] Günter Rote, Francisco Santos, and Ileana Streinu. Expansive motions and the polytope of pointed pseudo-triangulations. In *Discrete and computational geometry*, volume 25 of *Algorithms Combin.*, pages 699–736. Springer, Berlin, 2003.
- [RSS08] Günter Rote, Francisco Santos, and Ileana Streinu. Pseudo-triangulations—a survey. In *Surveys on discrete and computational geometry*, volume 453 of *Contemp. Math.*, pages 343–410. Amer. Math. Soc., Providence, RI, 2008.
- [RZ94] Victor Reiner and Günter M. Ziegler. Coxeter-associahedra. *Mathematika*, 41(2):364–393, 1994.
- [Sac23] Andrew Sack. A realization of poset associahedra. Preprint, [arXiv:2301.11449](https://arxiv.org/abs/2301.11449), 2023.
- [Sal19] Giulio Salvatori. 1-loop amplitudes from the Halohedron. *J. High Energy Phys.*, 2019(12):074, 16, 2019.
- [San09] Samson Saneblidze. The bitwisted Cartesian model for the free loop fibration. *Topology Appl.*, 156(5):897–910, 2009.
- [Sch11] Pieter Hendrik Schout. *Analytical treatment of the polytopes regularly derived from the regular polytopes. Section I: The simplex.*, volume 11. 1911.
- [Sch14] Ralf Schiffler. *Quiver representations*. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC. Springer, Cham, 2014.
- [Sei08] Paul Seidel. *Fukaya categories and Picard-Lefschetz theory*. Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2008.
- [She63] Geoffrey C. Shephard. Decomposable convex polyhedra. *Mathematika*, 10:89–95, 1963.
- [SP02] Richard P. Stanley and Jim Pitman. A polytope related to empirical distributions, plane trees, parking functions, and the associahedron. *Discrete Comput. Geom.*, 27(4):pp. 603–634, 2002.
- [SS93] Steve Shnider and Shlomo Sternberg. *Quantum groups: From coalgebras to Drinfeld algebras*. Series in Mathematical Physics. International Press, Cambridge, MA, 1993.
- [SS12] Luis Serrano and Christian Stump. Maximal fillings of moon polyominoes, simplicial complexes, and Schubert polynomials. *Electron. J. Combin.*, 19, 2012.
- [SS21] Giulio Salvatori and Stefan Stanojevic. Scattering amplitudes and simple canonical forms for simple polytopes. *J. High Energy Phys.*, 2021(3):067, 24, 2021.
- [SSW17] Francisco Santos, Christian Stump, and Volkmar Welker. Noncrossing sets and a Grassmann associahedron. *Forum Math. Sigma*, 5:e5, 49, 2017.
- [Sta63] James D. Stasheff. Homotopy associativity of H-spaces I & II. *Trans. Amer. Math. Soc.*, 108(2):275–312, 1963.
- [Sta97] Jim Stasheff. From operads to “physically” inspired theories. In *Operads: Proceedings of Renaissance Conferences (Hartfort, CT/Luminy, 1995)*, volume 202 of *Contemporary Mathematics*, pages 53–81, Cambridge, MA, 1997. American Mathematical Society. Appendix B by Steve Shnider and Jim Stasheff for a corrected polytope construction.
- [Sta04] Jim Stasheff. What is . . . an operad? *Notices Amer. Math. Soc.*, 51(6):630–631, 2004.
- [Ste64] Hugo Steinhaus. *One hundred problems in elementary mathematics*. Basic Books Inc. Publishers, New York, 1964.
- [Ste13] Salvatore Stella. Polyhedral models for generalized associahedra via Coxeter elements. *J. Algebraic Combin.*, 38(1):121–158, 2013.
- [Str12] Ross Street. Parenthetical remarks. In *Associahedra, Tamari lattices and related structures*, volume 299 of *Prog. Math.*, pages 251–268. Birkhäuser/Springer, Basel, 2012.
- [STT88] Daniel D. Sleator, Robert E. Tarjan, and William P. Thurston. Rotation distance, triangulations, and hyperbolic geometry. *J. Amer. Math. Soc.*, 1(3):647–681, 1988.
- [Stu11] Christian Stump. A new perspective on k -triangulations. *J. Combin. Theory Ser. A*, 118(6):1794–1800, 2011.
- [SU04] Samson Saneblidze and Ronald Umble. Diagonals on the permutohedra, multiplihedra and associahedra. *Homology Homotopy Appl.*, 6(1):363–411, 2004.

- [SU11] Samson Saneblidze and Ronald Umble. Matrads, biassociahedra, and A_∞ -bialgebras. *Homology Homotopy Appl.*, 13(1):1–57, 2011.
- [SU22] Samson Saneblidze and Ronald Umble. Comparing diagonals on the associahedra. Preprint, [arXiv:2207.08543](https://arxiv.org/abs/2207.08543), 2022.
- [Tam51] Dov Tamari. *Monoides préordonnés et chaînes de Malcev*. PhD thesis, Université Paris Sorbonne, 1951.
- [Tho12] Hugh Thomas. The Tamari lattice as it arises in quiver representations. In *Associahedra, Tamari lattices and related structures*, volume 299 of *Progr. Math.*, pages 281–291. Birkhäuser/Springer, Basel, 2012.
- [Tho21] Hugh Thomas. An introduction to the lattice of torsion classes. *Bull. Iranian Math. Soc.*, 47(suppl. 1):S35–S55, 2021.
- [Tho23] Hugh Thomas. Harder-Narasimhan polytopes, 2023. Talk at the Simons Center Workshop on Combinatorics and Geometry of Convex Polyhedra, Stony Brook University, USA. Available at <https://scgp.stonybrook.edu/video/video.php?id=5805>.
- [Ton97] Andy Tonks. Relating the associahedron and the permutohedron. In *Operads: Proceedings of Renaissance Conferences (Hartford, CT/Luminy, 1995)*, volume 202 of *Contemp. Math.*, pages 33–36. Amer. Math. Soc., Providence, RI, 1997.
- [Tro62] H. F. Trotter. Algorithm 115: Perm. *Commun. ACM*, 5(8):434–435, 1962.
- [Val14] Bruno Vallette. Algebra + homotopy = operad. In *Symplectic, Poisson, and noncommutative geometry*, volume 62 of *Math. Sci. Res. Inst. Publ.*, pages 229–290. Cambridge Univ. Press, New York, 2014.
- [Vie07] Xavier Viennot. Catalan tableaux and the asymmetric exclusion process. In *19th International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2007)*. 2007.
- [Wil17] Nathan Williams. W -associahedra have the non-leaving-face property. *European J. Combin.*, 62:272–285, 2017.
- [Woo04] Alexander Woo. Catalan numbers and Schubert polynomials for $w = 1(n+1)\dots 2$. Preprint, [arXiv:math/0407160](https://arxiv.org/abs/math/0407160), 2004.
- [Zel06] Andrei Zelevinsky. Nested complexes and their polyhedral realizations. *Pure Appl. Math. Q.*, 2(3):655–671, 2006.

(Vincent Pilaud) CNRS & LIX, ÉCOLE POLYTECHNIQUE, PALAISEAU
Email address: vincent.pilaud@lix.polytechnique.fr
URL: <http://www.lix.polytechnique.fr/~pilaud/>

(Francisco Santos) FACULTAD DE CIENCIAS, UNIVERSIDAD DE CANTABRIA, AV. DE LOS CASTROS s/n, E-39005 SANTANDER, SPAIN.

Email address: francisco.santos@unican.es
URL: <https://personales.unican.es/santosf/>

(Günter M. Ziegler) INST. MATHEMATICS, FU BERLIN, ARNIMALLEE 2, 14195 BERLIN, GERMANY.
Email address: ziegler@math.fu-berlin.de
URL: <http://www.mi.fu-berlin.de/math/groups/discgeom/ziegler>