



Non-Hausdorff manifolds over locally ordered spaces via sheaf theory

Yorgo Chamoun



LIX, Ecole polytechnique

- 1 Motivations
- 2 Some sheaf theory
- 3 The blowup construction
- 4 Precubical sets



Geometric semantics





Geometric semantics





Geometric semantics





Geometric semantics





Geometric semantics





Geometric semantics





Geometric semantics





Geometric semantics





Geometric semantics




Geometric semantics





Geometric semantics




Why blowing up?




Why blowing up?




Graph blowup




Graph blowup




Graph blowup




Graph blowup




Graph blowup




Graph blowup



An example which is not a graph product




An example which is not a graph product




$$\{z = 0\}$$

An example which is not a graph product




$$\{y = 0\}$$

An example which is not a graph product




$$\{x = 0\}$$

An example which is not a graph product




$$\{yz = 0 \wedge y \geq 0 \wedge z \leq 0\}$$

An example which is not a graph product




$$\{yz = 0 \wedge y \leq 0 \wedge z \geq 0\}$$

An example which is not a graph product




$$\{xz = 0 \wedge x \leq 0 \wedge z \geq 0\}$$

An example which is not a graph product




$$\{xz = 0 \wedge x \geq 0 \wedge z \leq 0\}$$

An example which is not a graph product



$$\{xy = 0 \wedge x \leq 0 \wedge y \geq 0\}$$

An example which is not a graph product



$$\{xy = 0 \wedge x \geq 0 \wedge y \leq 0\}$$

Outline

- 1 Motivations
- 2 Some sheaf theory
- 3 The blowup construction
- 4 Precubical sets

Some sheaf theory

Let X be a topological space.

Definition

- $O(X)$ is the poset category of open sets of X .
- A *presheaf* on X is a functor $O(X)^{op} \rightarrow \mathbf{Set}$.
- A presheaf F is a *sheaf* if the following diagram is an equalizer:

$$F(U) \longrightarrow \prod_{i \in I} F(U_i) \rightrightarrows \prod_{i,j \in I} F(U_i \cap U_j)$$

Some sheaf theory

Let X be a topological space.

Definition

- $O(X)$ is the poset category of open sets of X .
- A *presheaf* on X is a functor $O(X)^{op} \rightarrow \mathbf{Set}$.
- A presheaf F is a *sheaf* if the following diagram is an equalizer:

$$F(U) \longrightarrow \prod_{i \in I} F(U_i) \rightrightarrows \prod_{i,j \in I} F(U_i \cap U_j)$$

Example

$F : U \mapsto \mathcal{C}^0(U, \mathbb{R})$ with restriction defines a sheaf on X . In this case, the sheaf condition corresponds to the fact that a family $f_i : U_i \rightarrow \mathbb{R}$ satisfying

$$(f_i)|_{U_i \cap U_j} = (f_j)|_{U_i \cap U_j}$$

can be amalgamated into a unique $f : \bigcup_i U_i \rightarrow \mathbb{R}$.

The fundamental correspondance

Theorem

There is an adjunction between $Psh(X)$ and Top/X that restricts to an equivalence

$$\phi : Sh(X) \cong Et(X)$$

Definition

Let F be a sheaf on X . Let $x \in X$. The *set of germs of F at x* is given by

$$F_x := \operatorname{colim}_{U \ni x} F(U)$$

For $s \in F(U)$, the *germ of s at x* , noted s_x , is the equivalence class of s in F_x .

To build a bundle out of F , just take the projection $p : \bigsqcup_{x \in X} F_x \rightarrow X$.
The open sets are generated by

$$U_s := \{s_x \mid x \in U\} \quad (s \in F(U))$$

Outline

- 1 Motivations
- 2 Some sheaf theory
- 3 The blowup construction**
- 4 Precubical sets

The blowup construction

Let X be a locally ordered space.

Definition

The *sheaf of n -traversals* $O_X^{(n)}$ is defined by:

$$O_X^{(n)}(U) := \{A \subseteq U \mid A \cong E \text{ with } E \text{ } n\text{-euclidean local order}\}$$

for an open set $U \subseteq X$, the restriction map $O_X^{(n)}(V) \rightarrow O_X^{(n)}(U)$ for $U \subseteq V$ being given by intersection with U .

Definition

The *n -blowup* of X is defined by

$$\tilde{X} := \sqcup_{x \in X} \{A_x \in O_{X,x}^{(n)} \mid x \in A\} \subseteq \phi(O_X^{(n)})$$

(with the induced topology and local order). The *n -blowup map* is the projection $\beta_X : \tilde{X} \rightarrow X$.


Back to the example





The ghost space,
i.e. the elements $\emptyset_x, x \in X$



Back to the example




$$U_A$$




$$A \in O_n^{(X)}(U)$$

Back to the example





$$U_A$$




$$A \in O_n^{(X)}(U)$$



Back to the example


 U_A

 $A \in O_n^{(X)}(U)$

Back to the example



$$U_A$$



$$A \in O_n^{(X)}(U)$$

The universal property

Theorem

The n -blowup \tilde{X} satisfies the following universal property: for any n -euclidean local order E , for any local embedding $f : E \rightarrow X$, there is a unique continuous map $\tilde{f} : E \rightarrow \tilde{X}$ making the following diagram commute

$$\begin{array}{ccc} & & \tilde{X} \\ & \nearrow \tilde{f} & \downarrow \beta_X \\ E & \xrightarrow{f} & X \end{array}$$

Moreover, \tilde{f} is a local embedding.

Corollary

The category of n -euclidean local orders and local embeddings is a coreflective subcategory of the category of local orders and weakly euclidean morphisms (or local embeddings).

Outline

- 1 Motivations
- 2 Some sheaf theory
- 3 The blowup construction
- 4 Precubical sets

The concrete case

Definition

- A *graph* is a presheaf on the category $0 \rightrightarrows 1$.
- \square is the free monoidal category on $0 \rightrightarrows 1$ with unit 0 .
- A *precubical set* is a presheaf on \square .

Informally, a precubical set is a set of n -cubes for every $n \in \mathbb{N}$, glued together on their boundaries. A precubical set can be realized as a locally ordered space.

Proposition

The blowup of the locally ordered realization of a precubical set admits a combinatorial expression.


This description is encapsulated in a presheaf over the opposite of the category of elements of the precubical set.

- Let P be a precubical set.
- $\int P$ is the category of cubes of P and face inclusions.
- $\frac{1}{2}P$ is the 2-subdivision of P . In particular, every cube c of P has a central vertex $m(c)$ in $\frac{1}{2}P$.
- We define $Comb_P : \int P \rightarrow \mathbf{Set}$ as follows :


$$Comb_P(k, c) := \{e : (-1 \rightarrow 0 \rightarrow 1)^n \rightarrowtail \frac{1}{6}P \mid e(0) = m(c)\} \cup \{\emptyset_c\}$$

- This can also be seen as a relational presheaf over P .

Back to the (other) example...



The set $\text{Comb}_P(0, v)$



- The tools developped can be used to prove rigidity results on locally ordered realizations.
- We can see this contruction as an instance of an abstract construction on *geometric contexts à la Toën*.
- We conjecture the existence of a smooth atlas on the blowup of a precubical set.

- [1] Lisbeth Fajstrup, Eric Goubault, Emmanuel Haucourt, Samuel Mimram, and Martin Raussen. *Directed algebraic topology and concurrency*, 2016.
- [2] Pierre-Yves Coursole and Emmanuel Haucourt. *Non-existing and ill-behaved coequalizers of locally ordered spaces*, 2024.
- [3] Saunders MacLane and Ieke Moerdijk. *Sheaves in geometry and logic: A first introduction to topos theory*, 1992.
- [4] Emmanuel Haucourt. *Non-hausdorff parallelized manifolds over geometric models of conservative programs*, 2024.
- [5] Yorgo Chamoun and Emmanuel Haucourt. *Non-Hausdorff manifolds over locally ordered spaces via sheaf theory*, 2025.